



The effects of ethylene on the HCl-extractability of trace elements during soybean seed germination



Chunming Liu, Yixin Zhao, Jianfeng Liu, Wanting Gen, Yunqing Cheng *

College of Life Sciences, Jilin Normal University, Siping, Jilin 136000, China

ARTICLE INFO

Article history:

Received 2 April 2015

Accepted 26 May 2015

Available online 26 July 2015

Keywords:

ACC synthase

Bioavailability

Ethylene

Metallic elements

Seed germination

Soybean

ABSTRACT

Background: Ethylene is capable of promoting seed germination in some plant species. Mobilization of metals such as Fe, Cu, Mn, and Zn in mature seeds takes place when seeds are germinating. However, whether ethylene is involved in the regulation of soybean seed germination and metal element mobilization during early seed germination stage remains unknown. In the present study, seeds were treated with ethylene synthesis inhibitor aminoethoxyvinylglycine (AVG) and ethylene precursor 1-aminocyclopropane-1-carboxylic acid (ACC), and double distilled H₂O (ddH₂O) treatment was used as control. Ethylene emission, ACC synthase (ACS) expression, ACS enzyme activity and Ca, Zn, Mn, Cu and Fe content in hypocotyls were qualified to analyze the relationship between ethylene and mobilization of these elements.

Results: The results showed that ACS expression, ACS enzyme activity and ethylene emission peaked at 1 and 7 d after sowing. AVG inhibited ethylene production, promoted the hypocotyls length, ACS expression and its activity, concentrations of total and HCl-extractable Zn, and HCl-extractable Fe in hypocotyls, while ACC caused opposite effects. AVG and ACC treatment had no significantly effects on total and HCl-extractable Ca, Cu and HCl-extractable Mn. Total Mn concentration was promoted by AVG at 1, 3, and 5 d significantly, while ACC treatment tended to have no significantly effects on Mn concentration.

Conclusion: These findings suggested that ethylene is at least partly involved in the regulation of soybean seed germination. Remobilization of Zn and Fe may be negatively regulated by ethylene.

© 2015 Pontificia Universidad Católica de Valparaíso. Production and hosting by Elsevier B.V. All rights reserved.

1. Introduction

Seed germination involves regulation of a series of metabolic processes by plant hormones [1]. In some plant species, including soybean (*Glycine max* (L.) Merr.), it has been shown that detectable ethylene production begins with the onset of germination, i.e., with radicle emergence [2]. Moreover, many reports indicate that ethylene promotes seed germination [3,4]. The triple response, as described by Neljubov [5], indicates the plant sensitivity to ethylene. Understanding the mechanisms underlying the control of seed germination and its regulation by ethylene is not only of academic interest, but is also important for improving crop production and yield [6].

Minerals accumulated during seed development constitute less than 3% of the seed dry mass, yet they form an important pool of essential nutrients [7]. These mineral reserves are mobilized during germination and are a source of cofactors for enzymes, which are required for rapid growth [8]. The cotyledon of a soybean seed is very important for seed germination, seedling establishment, as well

as growth and survival because it serves as the main nutrient resource for young seedlings [1]. Large seed reserves of mineral nutrients may be of importance in order to support plant adaptation in micronutrient-deficient soils. Mobilization of metals such as Fe, Cu, Mn, and Zn in mature seeds takes place during germination and early seedling development. However, there is limited knowledge regarding the transport mechanisms of these immobile metal elements during the process of seed germination.

A large number of solutions have been used for metal extraction in order to assess the bioavailability of trace elements [8]. Single extractants may broadly be divided into three main classes: (1) weak replacement ion salts: MgCl₂, CaCl₂ and NH₄NO₃, (2) dilute solutions of either weak acids: acetic acid, or strong acids: HCl and HNO₃; and (3) chelating agents: pentetic acid (DTPA) and ethylenediaminetetraacetic acid (EDTA) [9]. The chelating agents DTPA and EDTA reduce the activity of the free metal ions in solution by forming complexes with the free metal ions. The second type, acid extractants are able to release into solution and can be considered as bioavailability [10,11]. Input of total and extractable metals in the growing hypocotyls is vital for enhancing seed germination and seedling growth. The transport and bioavailability of immobile metal element are dependent on a number of factors in addition to plant

* Corresponding author.

E-mail address: 1762369884@qq.com (Y. Cheng).

Peer review under responsibility of Pontificia Universidad Católica de Valparaíso.

Table 1

The primers used for RT-PCR and qRT-PCR amplification of GMACS gene.

Target gene	Primer	Primer sequence (5'–3')	Length (bp)	GC %	Amplification length (bp)
Actin	Actin-F	5'-ACCTCGACATACTGGTGTATGGTT-3'	25	44.00	81
	Actin-R	5'-ATACCTCTTTTGGATTGGGCTTC-3'	23	43.40	
ACS	ACC-F	5'-CACCTCAAATCCCGGTCAA-3'	19	54.55	105
	ACC-R	5'-AGCAACTGGAGCACACGAAG-3'	20	40.00	

growth regulators [12,13]. Little information is available regarding the ethylene regulation of immobile metal element transport during soybean seed imbibition and germination by far. The current study was carried out to examine ethylene metabolism, and its influence on the mobilization and bioavailability of immobile metal element during soybean seed germination, as well as to determine the dynamics of Fe, Mn, Zn, Ca, and Cu mobilization in hypocotyls.

2. Materials and methods

2.1. Materials and treatments

Soybean seeds of Tiefeng-31 were imbibed in 100 mg/L of aminoethoxyvinylglycine (AVG), 75 μ mol/L of 1-aminocyclopropane-1-carboxylic acid (ACC) and double-distilled H₂O (ddH₂O), respectively for 4 h and sowed in vermiculite. After thorough watering, the seeds were incubated in an illumination incubator at 25°C and 65% relative humidity for 7 d. Each treatment was performed in triplicates (100 seeds). At 1, 3, 5 and 7 d after sowing, hypocotyls of three treatments were sampled for further analysis.

2.2. Ethylene qualification

Fresh hypocotyls (0.5 g fresh weight) were sealed in an 8 mL vial and incubated for 4 h in darkness at 30°C. Then, the accumulated ethylene was quantified by gas chromatography with a glass column (2 mm \times 2 m) of Porapak N (Waters, Milford, MA, USA) at 80°C [14]. Peak areas were determined with a Chromatopak C-R6A system (Shimadzu, Kyoto, Japan).

2.3. Quantification of ACC synthase (ACS) activity

ACS was extracted and assayed using high performance liquid chromatography (HPLC) as previously described [15].

2.4. Quantification of ACS expression by RT-PCR and qRT-PCR

Total RNA was extracted from 100 mg hypocotyls using TRIZOL Reagent method (Invitrogen, Carlsbad, CA). The first strand cDNA synthesis and RT-PCR were performed using One Step RNA PCR kit (Takara Biochemicals, Kyoto, Japan) using oligo-dT and random

oligonucleotide primers, followed by amplification of the resulting DNA using polymerase chain reaction. The reaction was performed in an Eppendorf master cycler, which began with an initial denaturation step at 95°C for 3 min, followed by 35 cycles of 15 s at 94°C, 30 s at 45°C, 1.5 min at 72°C, and a final 7 min extension at 72°C. qRT-PCR analysis was performed following the method described by Cheng [14]. Primers used for RT-PCR and real-time amplification of GMACS cDNA were designed according to the sequences of soybean ACC (GMACS) gene. The primer sets are listed in Table 1.

2.5. Quantification of total and HCl-extracted Ca, Zn, Mn, Cu and Fe

Ca, Zn, Mn, Cu and Fe in hypocotyls were quantified using dry ashing and atomic absorption spectrometry method described by Altundag [16]. Ca, Zn, Mn, Cu and Fe were extracted with 0.03 M HCl solution described by Maki [17]. Then, their concentration was determined with atomic absorption spectrometry [16].

2.6. Data collection and statistical analysis

All experiments were conducted in triplicates. Statistical analysis was carried out with ANOVA process of SAS version 8.01 (SAS Institute, Inc., Cary, NC, USA). Means were compared using least significant difference (LSD) *t*-test at the 5% level of significance.

3. Results

3.1. Effects of ACC and AVG treatment on hypocotyls growth

Dry weights and lengths of hypocotyls in germination seeds were measured during the various stages of seed germination (Table 2). In the control, the hypocotyls length increased from 0.22 ± 0.02 cm at 1 d to 4.50 ± 0.24 cm at 7 d, and their dry weight increased from 0.0076 ± 0.0005 g at 1 d to 0.0208 ± 0.0015 g at 7 d. The results obtained from AVG and ACC treatments indicated that: a) AVG treatment strongly promoted hypocotyl length; b) the growth of hypocotyl length appeared to be more sensitive to AVG treatment after 5 to 7 d of incubation (the hypocotyl elongation stage); c) the effects of ACC on hypocotyls length were opposite to that caused by AVG, and d) in contrast to hypocotyl lengths, hypocotyl dry weight appeared to be insensitive to AVG or ACC treatments.

Table 2

Changes of hypocotyls growth, ACS expression by qRT-PCR, ACS activity, ethylene production during seed germination stage.

Treatment	Time (d)	Hypocotyls length (cm)	Hypocotyls dry weight (g)	ACS expression	ACS activity (nmol/g·h)	Ethylene (ppm/h·g)
AVG	1	0.22 ± 0.01 g	0.0075 ± 0.0002 d	0.180 ± 0.042 g	206.88 ± 10.20 fg	4.21 ± 0.23 e
	3	1.06 ± 0.03 f	0.0097 ± 0.0003 cd	0.120 ± 0.004 g	144.41 ± 5.11 hg	1.90 ± 0.15 f
	5	4.57 ± 0.176 b	0.0142 ± 0.0019 b	0.112 ± 0.022 g	35.59 ± 2.15 i	0.85 ± 0.07 f
	7	8.49 ± 0.15 a	0.0211 ± 0.0013 a	0.890 ± 0.039 e	254.60 ± 2.67 ef	11.85 ± 0.28 d
Control	1	0.22 ± 0.02 g	0.0076 ± 0.0005 cd	1.003 ± 0.006 d	355.20 ± 27.43 e	12.41 ± 0.57 d
	3	1.02 ± 0.02 f	0.0095 ± 0.0004 cd	0.333 ± 0.007 f	53.46 ± 1.11 hi	2.74 ± 0.28 ef
	5	2.57 ± 0.19 d	0.0141 ± 0.0033 b	0.147 ± 0.009 g	34.72 ± 1.80 i	1.86 ± 0.13 f
	7	4.50 ± 0.24 b	0.0208 ± 0.0015 a	1.523 ± 0.058 c	785.23 ± 7.40 d	20.75 ± 0.91 b
ACC	1	0.22 ± 0.02 g	0.0081 ± 0.0003 cd	1.776 ± 0.106 b	1234.68 ± 195.38 b	21.59 ± 0.66 b
	3	1.01 ± 0.02 f	0.0099 ± 0.0012 c	0.943 ± 0.022 de	871.27 ± 22.06 cd	11.60 ± 0.75 d
	5	1.75 ± 0.25 e	0.0149 ± 0.0007 b	1.458 ± 0.028 c	946.31 ± 29.13 c	16.95 ± 2.30 c
	7	2.850 ± 0.19 c	0.0211 ± 0.001 a	2.807 ± 0.055 a	1436.97 ± 71.45 a	39.12 ± 3.08 a

Data represent the mean \pm SD. Different letters within the same column indicated significant difference at 5% level by LSD.

Download English Version:

<https://daneshyari.com/en/article/200537>

Download Persian Version:

<https://daneshyari.com/article/200537>

[Daneshyari.com](https://daneshyari.com)