



Research article

Chromium-induced tropane alkaloid production and H6H gene expression in *Atropa belladonna* L. (Solanaceae) *in vitro*-propagated plantletsBahareh Vakili^a, Farah Karimi^{a,*}, Mozafar Sharifi^b, Mehrdad Behmanesh^c^a Dep. of Biology, Faculty of Basic Sciences, Shahed University, P.O.Box: 3319118651, Tehran, Iran^b Dep. of Plant Biology, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran^c Dep. of Genetics, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran

ARTICLE INFO

Article history:

Received 12 September 2011

Accepted 6 December 2011

Available online 13 December 2011

Keywords:

Atropa belladonna L.

h6h

Hyoscyamine

Scopolamine

Tropane alkaloids

ABSTRACT

Hyoscyamine and scopolamine tropane alkaloids found in several solanaceous plants are anticholinergic drugs. Hyoscyamine 6 β -hydroxylase (H6H) catalyzes two consecutive oxidation reactions. The first reaction is the hydroxylation of hyoscyamine to 6 β -hydroxyhyoscyamine and the second is epoxidation of 6 β -hydroxyhyoscyamine yielding scopolamine that is the final metabolite in the tropane alkaloid biosynthetic pathway. The effects of trivalent chromium as KCr (SO₄)₂ on the production of tropane alkaloids and the expression of hyoscyamine 6 β -hydroxylase gene (h6h) were studied in micro-propagated *Atropa belladonna* L. plantlets. The results showed that chromium treatment decreased the growth parameters (weights and lengths of the plantlets) and chlorophyll contents and increased proline contents. Moreover, semiquantitative RT-PCR analysis showed that the transcript level of H6H increased under chromium treatment. This treatment also increased hyoscyamine and scopolamine contents as shown by HPLC analysis. Changes of scopolamine contents correlate with the expression levels of h6h gene under different concentrations of chromium.

© 2011 Elsevier Masson SAS. All rights reserved.

1. Introduction

Solanaceaeous plants such as *Atropa*, *Datura*, *Duboisia*, *Hyoscyamus* and *Scopolia* exclusively produce the hyoscyamine and scopolamine tropane alkaloids and are widely used as anticholinergic drugs, which act on the parasympathetic nervous system [1]. Hyoscyamine is usually the main alkaloid in many of these plants while scopolamine is often produced in small amounts. Scopolamine is more valuable and is preferable to hyoscyamine in the pharmaceutical market because of its higher physiological activity and fewer side effects. There is currently a tenfold higher commercial demand for scopolamine (in the N butylbromide form) than for both hyoscyamine and atropine (the racemic mixture of D and L Hyoscyamine) [2]. In *Atropa belladonna*, these two tropane alkaloids mostly synthesize in young root cells and translocate to aerial parts of the plant [3].

The biosynthesis of scopolamine is a complex process requiring many distinct enzymatic steps and several genes of these enzymes have now been cloned. The last step of scopolamine biosynthesis is catalyzed by Hyoscyamine 6 β -hydroxylase (H6H, EC 1.14.11.11). H6H is a member of the 2-oxoglutarate-dependent dioxygenase family. The enzyme catalyzes two consecutive oxidation reactions, i.e. the hydroxylation of hyoscyamine to 6 β -hydroxyhyoscyamine and the epoxidation of 6 β -hydroxyhyoscyamine yielding scopolamine which is the final biosynthetic metabolite [4–8]. In addition to the alkaloid substrate H6H requires Fe²⁺, 2-oxoglutarate, O₂, and ascorbate for catalysis [4,8].

Among the strategies employed to enhance plant secondary metabolite production an outstanding one is the use of biotic and abiotic elicitors. An elicitor is a compound that not only induces accumulation of antimicrobial phytoalexins in plants but also stimulates any type of defense responses [9]. Vernay et al. [10] found that exogenous chromium induces an increase in the scopolamine synthesis from hyoscyamine in *Datura innoxia*. An increase in the expression of h6h gene can considerably enhance the production of scopolamine in organ cultures of *A. belladonna*, *H. niger* and *A. baetica* [2], which proves that metabolic engineering may be a feasible approach to improve scopolamine production.

In this study, we have investigated the effects of trivalent chromium [Cr (III)] on growth, chlorophyll and proline contents

Abbreviations: H6H, Hyoscyamine 6 β -Hydroxylase; IAA, Indol Acetic Acid; BA, Benzyl Adenine; PMT, Putrescine N-methyltransferase; TR (I and II), Tropinon Reductase (I and II).

* Corresponding author. Tel.: +98 21 51212624; fax: +98 21 51212601.

E-mail address: fkarimi@shahed.ac.ir (F. Karimi).

and the production of tropane alkaloids from *in vitro*-propagated *A. belladonna* L. plantlets while comparing scopolamine contents with the expression levels of h6h.

2. Results

2.1. Growth changes

The lengths and weights of shoots and roots significantly declined by increasing the chromium concentrations in treated plantlets compared to the control. The largest decrease was obtained from 1 mM chromium concentration in roots and shoots of the plantlets (Fig. 1).

2.2. Chlorophyll contents

The contents of chlorophylls a and b decreased in all the applied concentrations of the chromium. Chlorophyll a decreased more than Chlorophyll b compared to the control. The highest contents of chlorophylls a and b were observed in the control plantlets (Fig. 2).

2.3. Proline contents

Proline contents increased significantly in response to the increased chromium concentrations. The highest proline contents were observed in shoots and roots of the plantlets in 1 mM chromium treatment, and the lowest ones were seen in shoots and roots of the control plantlets (Fig. 3).

2.4. Alkaloid contents

The retention times for scopolamine and hyoscyamine were obtained 4.5 ± 0.5 and 12 ± 0.5 by HPLC analysis respectively (Fig. 4). Total alkaloid, hyoscyamine and scopolamine contents rose




Fig. 1. Changes in the lengths and weights of *in vitro*-propagated *A. belladonna* plantlets, under different concentrations of chromium. All the values are the means of three biological replicates. Values without a common letter are statistically different ($P < 0.05$).




Fig. 2. Changes in chlorophyll a and chlorophyll b contents obtained from aerial parts of *in vitro*-propagated *A. belladonna* plantlets, in response to different concentrations of chromium. All the values are the means of three biological replicates. Values without a common letter are statistically different ($P < 0.05$).

significantly as the Cr concentration increased (Fig. 5). The highest level of total alkaloids for shoots ($0.38 \text{ mg g}^{-1} \text{ DW}$) and roots ($0.33 \text{ mg g}^{-1} \text{ DW}$) were observed in 1 mM and 0.5 mM of Cr treatment respectively. The highest level of scopolamine content ($0.27 \text{ mg g}^{-1} \text{ DW}$) was observed in roots of the plantlets treated with 1 mM of chromium and the highest level of hyoscyamine content ($0.31 \text{ mg g}^{-1} \text{ DW}$) was seen in roots of the plantlets treated with 0.5 mM chromium. The ratios of hyoscyamine/scopolamine were 0.96, 1.2, 0.88 and 0.9 in shoots; and 1.35, 1.17, 1.38 and 0.9 in roots; for 0, 0.2, 0.5 and 1 mM chromium concentrations respectively. The comparison of total alkaloids and scopolamine contents in shoots and roots showed no significant difference for the control plantlets and the plantlets which were treated by 0.5 mM of the chromium. However, in these two Cr concentrations hyoscyamine contents in roots were significantly more than shoots. No significant differences were observed between the hyoscyamine and scopolamine contents of shoots compared to the roots in other Cr treated plantlets. In 1 mM of Cr treatment, total alkaloids content of shoots was significantly more than roots.

2.5. Expression of h6h

We observed the highest level of h6h expression, in 1 mM of chromium, for both shoots and roots of the *in vitro*-propagated




Fig. 3. Changes of proline contents in shoots and roots of *in vitro*-propagated *A. belladonna* plantlets in response to different concentrations of chromium. All the values are the means of three biological replicates. Values without a common letter are statistically different ($P < 0.05$).

Download English Version:

<https://daneshyari.com/en/article/2016278>

Download Persian Version:

<https://daneshyari.com/article/2016278>

[Daneshyari.com](https://daneshyari.com)