

## Short communication

# Winter–spring transition induces changes in nutrients and microbial biomass in mid-alpine forest soils



Michele Freppaz <sup>a, b, \*</sup>, Daniel Said-Pullicino <sup>a</sup>, Gianluca Filippa <sup>b</sup>, Fabienne Curtaz <sup>a, b</sup>, Luisella Celi <sup>a</sup>, Ermanno Zanini <sup>a, b</sup>

<sup>a</sup> Department of Agricultural, Forest and Food Sciences, University of Torino, Largo Paolo Braccini 2, 10095 Grugliasco, Italy

<sup>b</sup> NATRISK – Research Centre on Natural Risks in Mountain and Hilly Environments, University of Torino, Italy

## ARTICLE INFO

## Article history:

Received 6 February 2014

Received in revised form

7 July 2014

Accepted 10 July 2014

Available online 17 July 2014

## Keywords:

Snow-covered alpine soils

Seasonal variation

Resource availability

Microbial biomass dynamics

Amino sugars

## ABSTRACT

Active microbial biomass beneath seasonal snowpacks may influence biogeochemical processes in mountain soils. Here we show that soil microbial biomass in temperate forest ecosystems reaches its annual peak under snow, and declines rapidly as snow melts in the spring. This decline was associated with changes in temperature regime and resource availability. However, in contrast to alpine ecosystems at higher elevations, fungal/bacterial ratios, evaluated by amino sugar analysis, were lower in snow-covered winter soils and increased during snowmelt. We associated this shift to a faster decline in bacterial with respect to fungal biomass during snowmelt.

© 2014 Elsevier Ltd. All rights reserved.


Seasonally snow-covered ecosystems play a critical role in global carbon (C) and nitrogen (N) biogeochemical cycles. The development of a persistent snow cover in mountain forest soils which decouples soil from atmospheric temperatures may represent a key factor influencing the dynamics of the microbial biomass and related seasonal variations in C and N cycling (Brooks et al., 1998; Freppaz et al., 2008; Viglietti et al., 2013). These ecosystems are especially susceptible to climate change as small changes in temperature or precipitation may result in large variations in the amount and timing of snow cover (Barnett et al., 2005; Edwards et al., 2007). This could have important implications on organic matter (OM) turnover and nutrient cycling which in turn affect C source/sink functions and plant nutrition (Haei et al., 2013). The objectives of our study were to investigate temporal patterns in fungal and bacterial microbial biomass in seasonally snow-covered mountain forest soils, and to link these dynamics to the variations in available C and N under different forest stands.

The study was conducted in three forest sites (Navette, Secae and Valcona) located in the Ligurian mid-alpine area (Ormea, NW Italy, ~30 km North from the Mediterranean sea), under a maritime

snow climate, at elevations of 1609, 1312 and 1321 m a.s.l. respectively. Tree vegetation is dominated by Silver fir (*Abies alba* Mill.), beech (*Fagus sylvatica* L.) and Scots pine (*Pinus sylvestris* L.), respectively, while the soils are classified as Typic Udorthent, fine-loamy, mixed, active, nonacid, frigid (Navette), Typic Udorthent, loamy-skeletal, mixed, frigid (Secae) and Ultic Hapludalf, clayey-skeletal, mixed, frigid (Valcona) (Soil Survey Staff, 2010). Annual precipitation during the experimental period (2010/11) accumulated to 1282 mm and the mean annual air temperature was +5.4 °C. In 2010, snowpack accumulation started in the middle of November reaching a maximum of 216 cm by the middle of March 2011, while snowmelt lasted throughout April 2011 (Fig. 1a). At each of the experimental sites, triplicate soil samples (200 cm<sup>3</sup> core) were collected from the A horizon (0–10 cm depth) on the 21st October 2010 and 26th January, 8th March, 5th May and 13th July 2011. Within 24 h from collection, fresh samples were homogenized by sieving at 2 mm, and analysed for microbial biomass C and N, inorganic N and water-extractable organic carbon (WEOC) and nitrogen (WEON). Fungal and bacterial microbial biomass were also evaluated by analysing for amino sugars as described by Zhang and Amelung (1996). These biomarkers are important constituents of the cell wall of bacteria and fungi and can thus be used as an indicator for the origin of different microbial C and N residues in soils (Amelung, 2001; Glaser et al., 2004). To test for difference between sampling dates and sites we used a linear mixed effect

\* Corresponding author. Department of Agricultural, Forest and Food Sciences, University of Torino, Largo Paolo Braccini 2, 10095 Grugliasco, Italy.

E-mail address: [michele.freppaz@unito.it](mailto:michele.freppaz@unito.it) (M. Freppaz).



**Fig. 1.** Variations in (a) daily average air temperatures, snowpack depth (shaded area), and (b) daily average soil temperatures measured at each of the three sites over the experimental period.


model (Bates et al., 2014), incorporating two fixed effects (sampling date and site) and two random effects (site and replicate) to account for autocorrelation data. Significant differences between sampling dates and sites were then identified at  $p < 0.05$  by the Tukey Honest significant difference test. Pearson correlation analyses were run with soil C and N forms.

Mean air temperatures during the snow-covered winter months fluctuated between  $-11.4$  and  $+8.0$  °C, while daily mean soil temperatures in all sites were rather constant with mean values of  $+0.5$ ,  $+1.1$  and  $+1.2$  °C for Valcona, Secae and Navette, respectively (Fig. 1). Although it is generally assumed that microbial activity is related to temperature (Pietikäinen et al., 2005), long-term soil warming may actually reduce biomass and activity of the microbial community (Rousk et al., 2012). Recent studies have shown that soil conditions under alpine snowpacks can be very favourable for microbial growth and activity (Brooks et al., 2011), with important impacts on element cycling and annual fluxes. In all sites, greater content of microbial biomass C was observed in January and March with respect to the other sampling dates (Fig. 2a), as reported for alpine tundra (Lipson et al., 2000) and temperate beech forest sites (Kaiser et al., 2011). Moreover, seasonal variations were greater than the differences that could be attributed to vegetation cover (Björk et al., 2008). Accordingly, mixed effect models for these two variables showed  $F$  values that were 1–2 orders of magnitude greater for factor time than for factor site, indicating a much higher variance explained by seasonal variation compared to vegetation cover.

Microbial biomass in forest soils is not only controlled by temperature and moisture status, but also by the supply of nutrients and labile organic C required to sustain microbial growth. In fact, relatively high biomass C content observed during the winter months coincided with highest concentrations of inorganic N ( $r = 0.314$ ;  $p < 0.05$ ), and labile WEOC ( $r = 0.496$ ;  $p < 0.05$ ) and WEOC ( $r = 0.441$ ;  $p < 0.05$ ) pools (Fig. 2b and c). Snow-covered soil temperature and moisture regimes may strongly influence nutrient release and labile C supply through their effects on litter decomposition as well as nutrient and dissolved organic C fluxes (Celi et al., 2010; Brooks et al., 2011). During the snow-covered winter months, the suppression of nitrification and denitrification at low temperatures (Frolking et al., 1998), together with limited plant uptake could have contributed to increasing  $\text{NH}_4^+$  concentrations (Fig. 2b). Similarly, slower rates of mineralization during litter degradation could have resulted in a net accumulation of labile WEOC and N pools (Fig. 2c). This was most evident at the Navette

site that also showed lowest mean winter soil temperatures. Similarly, Koranda et al. (2013) found that the winter community in a temperate beech forest soil showed a higher capacity for degrading complex C substrates (cellulose, plant cell walls) but a lower utilization of labile C sources than the summer community. As temperatures rise, the microbial biomass and growing vegetation compete for available N that, together with significant nutrient leaching during snowmelt, were probably responsible for the decrease in inorganic N contents with time (Fig. 2b). Moreover, the rapid consumption of labile OM with warmer temperatures may result in a decrease in C availability that coincides with the increase in microbial C demand causing a steep decline in biomass (Brooks et al., 1998; Lipson et al., 2000).

Variations in total amino sugars reflected the changes in microbial biomass with greatest biomarker contents during the winter months (Fig. 3). We assume that the observed seasonal variations in amino sugars were due to fresh biomass in winter and accelerated microbial turnover during the warmer months. We generally observed a higher abundance of fungal biomarkers (GluN) during the winter that decreased during snowmelt ( $p < 0.05$ ). This is consistent with other studies that report a higher



**Fig. 2.** Seasonal variations in (a) microbial biomass C and N, (b) inorganic N, and (c) water extractable organic C and N. Error bars represent the standard error. Different letters indicate significant differences ( $p < 0.05$ ) between sampling dates, as determined by the Tukey HSD test.

Download English Version:

<https://daneshyari.com/en/article/2024569>

Download Persian Version:

<https://daneshyari.com/article/2024569>

[Daneshyari.com](https://daneshyari.com)