

Short communication

Spatial heterogeneity provides organic matter refuges for soil microbial activity in the Patagonian steppe, Argentina

Marina Gonzalez-Polo*, Amy T. Austin

Instituto de Investigaciones Fisiológicas y Ecológicas Vinculadas a la Agricultura (IFEVA), Facultad de Agronomía, CONICET and Universidad de Buenos Aires, Avenida San Martín 4453, Buenos Aires C1417DSE, Argentina

ARTICLE INFO

Article history:

Received 28 August 2008

Received in revised form

2 March 2009

Accepted 11 March 2009

Available online 3 April 2009

Keywords:

Microbial biomass

β -Glucosidase activity

Mosses

Arid ecosystems

Extracellular enzymes

Biogeochemistry

Carbon cycling

Islands of fertility

Semi-arid steppe

Patagonia

Argentina

ABSTRACT

In arid and semi-arid ecosystems that are frequently classified as water limited, it is unclear how spatial and temporal variability of vegetation and climate could affect microbially-mediated soil processes. Our objective was to determine how aboveground spatial heterogeneity creates characteristic soil conditions that modulate microbial growth and activity in a semi-arid Patagonian steppe. In particular, we explored how micro-environmental and biogeochemical soil characteristics generated by the native vegetation could control soil β -glucosidase activity. Both life-form (shrubs, grasses, mosses and bare soil) and season exerted strong controls on all measured abiotic (soil temperature and gravimetric soil water content, inorganic nitrogen, pH and total C and N) and biotic (microbial biomass C and β -glucosidase activity) soil characteristics. Partial correlation between β -glucosidase activity and extracellular organic C (EOC) was high across seasons ($r = 0.5$; $P < 0.001$) while soil water content did not correlate with soil enzymatic activity ($r = 0.09$; $P > 0.05$). We postulate that labile soil carbon rather than water availability functions as a principal limitation of microbial activity in this semi-arid ecosystem, and the distribution of this carbon is, in large part, determined by the patchy distribution of vegetation.

© 2009 Elsevier Ltd. All rights reserved.

Above- and belowground communities are inextricably linked in terrestrial ecosystems; belowground biological processes are modulated by both the quantity and quality of organic material that enters the soil pool from net primary production (Wardle, 2002). In arid and semi-arid ecosystems, vegetation is typically dispersed in a matrix of bare soil creating high resource microsites associated with vegetated patches known as “islands of fertility” (Charley and West, 1975), which have been well-documented in deserts (Schade and Hobbie, 2005; Schaeffer and Evans, 2005). In addition, greater C-degrading and nutrient mineralization enzyme activity under vegetated patches and showing seasonal fluctuations have been reported in a number of low rainfall ecosystems (Pavel et al., 2004; Thompson et al., 2006; Goberna et al., 2007).

Water availability is clearly one of the major limitations of net primary production in semi-arid ecosystems (Austin and Sala, 2002; Muldavin et al., 2008). However, the magnitude of the

water pulse that stimulates microbially-mediated process greatly differs from the ones for vascular plants. Small and short pulses could stimulate soil microbial activity but have no effect on germination of many desert plants (Schwinning and Sala, 2004). The temporal distribution of rainfall, both seasonal variation and size of rainfall events, can regulate microbial processes (Austin et al., 2004). The other dominant regulator of soil microbial processes is temperature, which also has a strong temporal dimension in deserts affecting microbial activity (Fierer et al., 2003) and microbial community composition (Cookson et al., 2007). It is unclear how the constraints of water availability and temperature interact with vegetation patches to determine soil ecosystem processes.

Our objective in this study was to determine how aboveground spatial and temporal heterogeneity generated by the presence of vegetation patches could create characteristic conditions that control soil β -glucosidase activity. We hypothesized that indirect effects of low water availability (low organic matter inputs) may be more important than direct regulation of biological processes as a control on soil microbial activity.

* Corresponding author. Tel.: +54 11 45248070; fax: +54 11 45148730.

E-mail address: maripolo@agro.uba.ar (M. Gonzalez-Polo).

The study was conducted in a semi-arid steppe (1972 grazing enclosure), in the Río Mayo experimental station of INTA (Instituto Nacional de Tecnología Agropecuaria) (45°41'S and 70°16'W). Mean monthly temperature ranges from 1 °C (July) to 15 °C (January). Precipitation (150 mm) falls primarily in winter (Paruelo and Sala, 1995). Soils are coarse-textured aridisols (Valle, 1998). We established 4 microsites: shrubs ($n = 18$) (*Mulinum spinosum* (Cav.) Pers., *Adesmia volckmanni* Phil. and *Senecio filaginoides*), grasses ($n = 18$) (*Poa ligularis*, *Stipa speciosa*, and *Bromus pictus*), below moss patches ($n = 6$) and in bare soil ($n = 6$). We collected soil (0–5 cm) under each of these microsites with the exception of the moss microsite (0–2 cm) in September 2003 (spring), January 2004 (summer) and May 2004 (fall). We installed 12 plots (5 × 5 m) with four transects of 5 m for vegetative cover measurements. Percent vegetative cover in the study area was calculated using a line-intercept technique (Canfield, 1941).

Soils extracts were analyzed for NH_4^+ and NO_3^- using an Alpkem Flow IV® autoanalyzer (O-I Corporation, College Station, TX, USA) (Robertson et al., 1999). Total C and N were determined with an elemental analyzer (LECO) (Robertson et al., 1999). Soil temperature was measured at 10, 12 and 16 h integrating 10 cm depth. Microbial biomass C was measured using a modified chloroform fumigation–extraction technique (Jenkinson and Powlson, 1976), using a cylindrical piece of mesh in order to improve the diffusion efficiency of the chloroform. The C concentration in the unfumigated sample (0.5 M K_2SO_4), represents the most labile organic matter fraction in the soil, termed extracellular organic C (EOC) (Hofman and Dusek, 2003). The 1–4 β -glucosidase activity was assayed using p-nitrophenyl- β -glucopyranoside as a substrate dissolved in acetate buffer (pH 5). The reaction was stopped with 0.2 ml of NaOH (0.1 N) (Sinsabaugh et al., 1999). Enzyme activity and microbial biomass were determined within 15 days of sampling.

We evaluated differences in abiotic and biotic characteristics of microsites with two-way ANOVA (log transformed in the case of non-normal distributions). We used a non-parametric Kruskal–Wallis test for total soil C and N data that could not be normalised. Statistical analyses were performed using Statistica 6 software (Stat-soft, Inc., Tulsa, OK, USA). Pearson partial and multiple correlation coefficients were used to evaluate the degree of association between β -glucosidase activity and the other variables for each season and across all seasons. We calculated multiple correlation coefficients and the fifth order partial correlation coefficient between β -glucosidase and each variable named in Table 2 (Sokal and Rohlf, 1995) using a numerical Python script (Ascher et al., 2001) that performed the algebraic calculations over the correlation matrix.

Microsites generated by the different life-forms varied significantly and exerted a strong and significant control on abiotic soil characteristics (Table 1, Figs. 1 and 2). Bare soil had 50% or less carbon and nitrogen than the other microsites (Table 1). Microclimatic soil variables, including thermal amplitude and soil water content showed a significant season–life-form interaction with the exception of pH (Fig. 1). Thermal amplitude was most marked in




Fig. 1. pH, water content and thermal amplitude in soils under different life-forms for three sampling dates and the annual average. Thermal amplitude was defined as the difference between the maximum and minimum temperature registered during the day. Results for two-way ANOVA of life-form and season are shown. Bars indicate means (+SE), and an alpha of 0.05 was used in all cases.

spring, and the presence of shrubs appeared to buffer temperature extremes (Fig. 1C). Soil nitrate under shrubs was approximately 3-fold greater than in the rest of the life-forms ($P < 0.001$, Fig. 2A), consistently across seasons. By contrast, soil ammonium showed peaks of high concentration under particular life-forms depending on the season (Fig. 2B).

Soil microbial biomass C demonstrated both seasonal and spatial variation, except for the moss microsite that was constant and elevated throughout the year (Fig. 3A). EOC was greater beneath shrubs in all seasons, but additionally, there were pulses of EOC in the moss microsites in summer and fall (Fig. 3B). β -Glucosidase activity in soils under shrubs was greatest, and was similar to the moss microsite throughout the year ($P < 0.05$, Fig. 3C); fall was the season with lowest β -glucosidase activity across microsites (Fig. 3C). Bare soil showed minimum values of enzymatic activity

Table 1

Means (SE) for vegetation and bare soil cover and total soil C, N, and C/N ratios for all soil microsites in the semi-arid Patagonian steppe. Different letters indicate significant differences ($P < 0.05$) among soils collected under dominant shrub and grasses species, mosses and bare soil (Kruskal–Wallis test).

Life-form	% Cover	n	Soil %C	Soil %N	Soil C/N ratio
Shrubs	11.8	18	1.4 (0.04) ^a	0.11 (0.002) ^a	12.2 (0.11) ^a
Grasses	16	18	0.7 (0.01) ^{bc}	0.06 (0.008) ^{bc}	9.9 (0.06) ^{bc}
Mosses	1.4	6	0.8 (0.04) ^{ac}	0.06 (0.003) ^{ac}	12.4 (0.30) ^{ac}
Bare soil	51.4	6	0.4 (0.01) ^b	0.04 (0.001) ^b	9.6 (0.32) ^b

Download English Version:

<https://daneshyari.com/en/article/2025642>

Download Persian Version:

<https://daneshyari.com/article/2025642>

[Daneshyari.com](https://daneshyari.com)