



Trilobolide–porphyrin conjugates: On synthesis and biological effects evaluation



Pavla Tomanová^a, Silvie Rimpelová^b, Michal Jurášek^a, Miloš Buděšínský^c, Lucie Vejvodová^d, Tomáš Ruml^b, Eva Kmoníčková^{d,*}, Pavel B. Drašar^{a,*}

^a Department of Chemistry of Natural Compounds, Institute of Chemical Technology in Prague, CZ-166 28 Prague, Czech Republic

^b Department of Biochemistry and Microbiology, Institute of Chemical Technology in Prague, CZ-166 28 Prague, Czech Republic

^c Institute of Organic Chemistry and Biochemistry, CZ-166 10 Prague, Czech Republic

^d Institute of Pharmacology and Toxicology and Biomedical Centre, Faculty of Medicine in Pilsen, Charles University in Prague, CZ-301 66 Pilsen, Czech Republic

ARTICLE INFO

Article history:

Received 15 June 2014

Received in revised form 14 August 2014

Accepted 25 August 2014

Available online 6 September 2014

Keywords:

Trilobolide

Porphyrin

Nitric oxide

Fluorescence microscopy

ABSTRACT

Trilobolide (Tb), a potent natural counterpart of thapsigargin, is a sesquiterpene lactone of guaianolide type isolated from horse caraway (*Laser trilobum*, L. Borkh). Tb exerts remarkable pharmacological properties based on irreversible inhibition of sarco/endoplasmic reticulum calcium ATPase (SERCA), thus being of increasing interest for cancer cure. Additionally, another pharmacological activity of Tb, as well as of thapsigargin, was reported in several studies, Tb as being an effective inducer of nitric oxide and cytokine production. These extraordinary biological properties move these molecules in further pre-clinical evaluation.

Because of ubiquitous character of SERCA expression, development of specifically targeted bioactive molecules is inevitable. Since it is well known that porphyrins are preferentially taken up by cancer cells, we have designed and synthesized novel Tb–porphyrin conjugates. Copper-catalyzed azide–alkyne cycloaddition was used to link Tb with porphyrin at once. Two model conjugates of Tb and porphyrin were synthesized and properly characterized. Employing naturally occurring fluorescence properties of porphyrins, we investigated the intracellular localization of the conjugates employing fluorescence microscopy in living cells. Intriguingly, the prepared conjugates localized both in mitochondria and lysosomes of HeLa and LNCaP cells. Furthermore, the cytotoxicity of Tb–porphyrin conjugates was assessed in a number of human cancer cell lines and rat peritoneal cells. Likewise in cancer cell lines, viability of rat peritoneal cells was not affected by the tested conjugates. Interestingly, we observed dose-dependent nitric oxide (iNOS) production induced by the tested conjugates. The effect was related to the type of a linker used and the overall size of the molecule. Another potent immunobiological effects are under evaluation.

In summary, the results presented here indicate notable immunobiological potential of the prepared Tb conjugates. Moreover, they could help to decipher the molecular mechanism of Tb for its possible biomedical applications.

© 2014 Elsevier Inc. All rights reserved.

1. Introduction

Porphyrins are a class of naturally occurring compounds which have become intensively studied in last decades due to their unique photochemical properties. The ability of porphyrins to passively target tumors *in vivo* enables their utilization in cancer therapy and diagnostics [1]. The passive tumor targeting by porphyrins is based on enhanced retention and permeability effect of solid tumors [2]. Photodynamic therapy (PDT) is a non-invasive thera-

* Corresponding authors.

E-mail addresses: eva.kmonickova@lfp.cuni.cz (E. Kmoníčková), pavel.drasar@vscht.cz (P.B. Drašar).

peutic approach based on a use of light-sensitive molecules, photosensitizers. PDT is used worldwide for treatment of a number of diseases, including age-related macular degeneration, psoriasis and cancer [3–8]. Moreover, the photophysical properties of porphyrins allow visualization of their localization as well as of their conjugates both *in vitro* and *in vivo*. Conjugation of the macrocycle with counterpart of choice facilitates its transport via drug- or receptor-mediated endocytosis, affects delivery to a specific location within the cells and generally improves biological effects. Therefore, many drugs based on porphyrins are designed as pro-drug systems to enhance their physico-chemical and pharmacokinetic properties [9]. Many groups have focused on conjugation of porphyrins with other biologically active compounds, such as

saccharides [10,13], peptides [12,14], steroids [15] and polymers [16]. Conjugation is also responsible for specific accumulation in cells [10], which determines biological effects of the conjugate. Mitochondria [11], endoplasmic reticulum [10] and nucleus [12] are believed to be the main organelles for the pharmacotherapeutic intervention. Unfortunately, until now most of the conjugates accumulate in endosomes and lysosomes, as it was described for porphyrins conjugated to saccharides [10,13], peptides [14], dendrimers [15,16], and retinoids [17].

In this study, we present synthesis of two porphyrin conjugates containing a sesquiterpene lactone, trilobolide (Tb). Kmoníčková et al. [18] reported immunostimulatory properties of Tb, which is able to induce interferon gamma (INF- γ) secretion and nitric oxide (NO) release [18]. Another pharmacological feature of Tb is its potency to inhibit sarco/endoplasmic reticulum Ca^{2+} -ATPase (SERCA). The SERCA inhibition leads to accumulation of calcium ions in cytosol, and a sustained increase of Ca^{2+} induces apoptotic pathway and cell death. [19]. We imaged the transfer of the prepared compounds in cells of various tumor origins. Biological activities of the bioconjugates, such as induction of NO release in rat macrophages have also been evaluated. Moreover, we assessed cytotoxic potency of Tb–porphyrin conjugated in cancer cell lines of various histogenic types.

2. Results and discussion

2.1. Chemistry

We chose copper-catalyzed azide-alkyne cycloaddition [20,21] (CuAAC) as the key reaction of the designed synthesis. Thus, both parts of the conjugate, Tb and porphyrin, were modified to provide intermediates suitable for click chemistry. The chemical transformation of Tb is displayed in Scheme 1 and all of the experimental synthetic details are described in Supplementary Section 1. We synthesized the carboxymethyloxime (CMO) derivative **1c** in three steps. Briefly, Tb was transformed to its demethylbutanoyl derivative (**1a**) by mild solvolysis and successively the sole secondary hydroxy group was oxidized using pyridinium chlorochromate (PCC). Obtained 8-oxo-Tb derivative **1b** was transformed to Tb–CMO **1c** by the reaction with *O*-(carboxymethyl)hydroxylamine hemihydrochloride under pyridine catalysis. Finally, we introduced terminal alkyne moiety by the reaction of **1c** with propargyl alcohol in presence of *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide (EDC). Obtained propargylester **1d** was used in subsequent click reaction.

The synthesis of 5-(4-carboxyphenyl)-10,15,20-triphenylporphyrin derivatives is displayed in Scheme 2. First we prepared the

basic porphyrin according to a previously described method [22], see Supplementary information 1.2.5 and Scheme S1. Thereafter, we introduced azido PEG₃-amine and propargyl moiety into the porphyrin molecule using EDC chemistry (see Scheme 3).


Desired Tb–porphyrin conjugates were prepared using standard click chemistry protocol with CuI and Tris[(1-benzyl-1*H*-1,2,3-triazolyl)-methyl]amine (TBTA) [23] as an accelerator of the product formation. The analytical data including NMR, IR, HRMS, and optical rotation characteristics are described in Supplementary Section 1.2. Both conjugates showed absorbance maxima at 424 nm (Soret band) and emission maxima at 605 and 655 nm (excitation wavelength of 429 nm; see Supplementary 1.3 and Fig. S1), which are typical porphyrin spectral properties. Prior to biological testing, all samples were re-purified by column chromatography and their purity was checked by chromatographic methods.

2.2. Intracellular localization of Tb–porphyrin conjugates

After successful synthesis of Tb–porphyrin conjugates, we studied their intracellular localization in several human cancer cell lines: HeLa, LNCaP, U-2 OS, MCF-7, and MiaPaCa-2. We performed live-cell fluorescence microscopy of these red-emitting conjugates (**3** and **4**) for time intervals ranging from 20 min up to 24 h. Detectable fluorescent emission intensity of the tested derivatives **3** and **4** was observed only after 2 h of incubation with the model cell lines. Thus, the kinetics of Tb–porphyrin conjugates was significantly decreased when compared with the rapid intracellular uptake of Tb–Bodipy occurring already after 20 min of incubation with human cancer cells reported by Jurášek et al. [24]. In HeLa cells, Tb–porphyrin **3** (5 μM) localized in network-like patterned organelles, probably of mitochondrial origin, after 2 h of incubation. The localization of **3** did not change over prolonged incubation periods of 5 h (see Fig. 1), 16 h, and 24 h. Compound **4** (5 μM) was distributed in vesicles of endosomal or lysosomal origin in HeLa cells after 2 h and the intracellular distribution was retained at least up to 24 h. Compound **4** occurred partially also in large fluorescent aggregates probably caused by decreased water solubility. We have observed similar behavior of both compounds in living prostatic cancer cells (LNCaP), data shown in Supplementary Section in Fig. S2, and in human osteosarcoma cells (U-2 OS, see Fig. S3), the only difference was more pronounced aggregation of compound **4**.

2.3. NO production in primary macrophages

Within a group of sesquiterpene lactones, Tb possesses strong activity in stimulating nitric oxide (NO) production by immune



Scheme 1. Synthesis of functionalized Tb via carboxymethyloxime derivative. Reagents and conditions: (a) Et_3N , MeOH, 17 h, rt, yield 69%; (b) PCC, CH_2Cl_2 , rt, 2.5 h, yield 74%; (c) *O*-(carboxymethyl)hydroxylamine hemihydrochloride, pyridine, MeOH, rt, 70 min, yield 89%; (d) propargyl alcohol, EDC, HOBT, DMAP, CH_2Cl_2 , 0 °C → rt, 12 h, yield 86%. Compound **1e** was available in our laboratory.

Download English Version:

<https://daneshyari.com/en/article/2027818>

Download Persian Version:

<https://daneshyari.com/article/2027818>

[Daneshyari.com](https://daneshyari.com)