



HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtbOriginal article <http://dx.doi.org/10.1016/j.apjtb.2016.04.011>

Trichomonas vaginalis infection in a low-risk women attended in Obstetrics and Gynaecology Clinic, Universiti Kebangsaan Malaysia Medical Centre



Norhayati Mokhtar¹, Nor Liyana Ismail¹, Phoy Cheng Chun¹, Mohamad Asyrah Sapie¹,
Nor Farahin Abdul Kahar¹, Yusof Suboh¹, Noraina Abdul Rahim¹, Nor Azlin Mohamed Ismail²,
Tengku Shahrl Anuar^{3*}

¹Department of Parasitology and Medical Entomology, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia

²Department of Obstetrics and Gynaecology, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia

³Department of Medical Laboratory Technology, Faculty of Health Sciences, Universiti Teknologi MARA, Puncak Alam Campus, Selangor, Malaysia

ARTICLE INFO

Article history:

Received 22 Dec 2015

Received in revised form 4 Jan,

2nd revised form 30 Mar 2016

Accepted 27 Apr 2016

Available online 11 Jun 2016

Keywords:

Trichomoniasis

Incidence rate

Wet mount

Giemsa staining

Culture

ABSTRACT

Objective: To investigate the presence of trichomoniasis among women attending the Obstetrics and Gynaecology Clinic, Universiti Kebangsaan Malaysia Medical Centre.

Methods: A total of 139 high vaginal swabs were taken from the subjects and sent to the laboratory in Amies gel transport media. The specimens were examined for the presence of *Trichomonas vaginalis* using wet mount, Giemsa staining and cultured in Diamond's medium. Sociodemographic characteristics and gynaecological complaints were obtained in private using structured questionnaire applied by one investigator.

Results: The median age was 32 years, with an interquartile interval of 9.96. Most of the subjects were Malays (76.9%) and the remaining were Chinese (15.1%), Indians (2.2%) and other ethnic groups (5.8%). One hundred and thirty eight (99.3%) of the women were married and 98.6% had less than 6 children. More than half (75.5%) of the women's last child birth was less than 6 years ago. Forty seven percent of them were involved in supporting administrative work and 64.7% of the women gave a history of previous or current vaginal discharge.

Conclusions: The present study reported zero incidence rate of trichomoniasis. The low incidence rate was postulated due to all women who participated in this study were categorized into a low-risk group.

1. Introduction

Trichomonas vaginalis (*T. vaginalis*), which is an obligate single-celled protozoan parasite, is the most common curable

sexually transmitted disease (STD) and despite high prevalence rates ranging from 5% to 75% has received little attention globally [1]. It is predominantly spread via unprotected intercourse with an infected partner but might also be spread through the fingers after masturbation [2]. There are several factors that associated with high incidence of infection such as poor personal hygiene, multiple sex partners, low socioeconomic status and underdevelopment [3]. Globally, *T. vaginalis* infection affects approximately 57–180 million people with the majority living in developing countries [2,4].

This infection is strongly related with the existence of other STD such as chlamydia, gonorrhoea and HIV. Previous history of STD is an important sign of trichomoniasis among adolescents and better STD treatment can decrease the incidence of HIV infection [5,6]. *T. vaginalis* infection usually involves the urogenital tract, affecting both female and male of the reproductive age. The common sites of infection are vagina

*Corresponding author: Dr. Tengku Shahrl Anuar, Department of Medical Laboratory Technology, Faculty of Health Sciences, Universiti Teknologi MARA, Puncak Alam Campus, Selangor, Malaysia.

Tel: +60 3 3258 4425

Fax: +60 3 3258 4599

E-mail: tengku9235@puncakalam.uitm.edu.my

The study protocol was performed according to the Helsinki declaration and approved by Ethical Committee of Universiti Kebangsaan Malaysia Medical Centre. Informed consent was sought and obtained from individuals prior to participation in this study.

Foundation Project: Supported by the UKMMC Fundamental Research Grant (Grant No. FF-165-2011) from Universiti Kebangsaan Malaysia.

Peer review under responsibility of Hainan Medical University. The journal implements double-blind peer review practiced by specially invited international editorial board members.

and urethra in female and male, respectively. However, most of them may appear asymptomatic especially in male and they become the source of infection [7]. Symptoms typically occur after a period of incubation of 4–28 days. In women, the usual presenting complaint is vaginal discharge (yellowish to greenish in color), which often has a foul smell and is frothy. There may be complaints of itching sensation in the vulva, vagina and medial aspects of the thighs. They may also have urethritis, dyspareunia, cystitis, cervicitis or infection of the Skene's ducts and Bartholin's gland [8].

Trichomoniasis can cause a serious discomfort to women such as adverse pregnancy outcomes, manifested by preterm rupture of membranes, low-birth-weight infants, infertility and cervical cancer. It also increases the transmission of HIV [9]. The Centers for Disease Control and Prevention guidelines recognize that typical diagnosis is made by microscopic examination of wet mount preparations and the measurement of vaginal pH, which has a sensitivity of approximately 60% [10]. When there is a high index of suspicion for trichomoniasis unconfirmed by microscopic examination, culture of vaginal specimens should be performed as this method improves the sensitivity to 95% [11].

In Malaysia, there is a minimal research done to survey the prevalence of *T. vaginalis* infection in a low-risk group or in general population. Most of the researches focused on sub-populations which are known to be at high-risk for contracting and spreading STD such as sex workers and drug abusers. Hence, this study was undertaken to investigate the presence of *T. vaginalis* among women, which were categorized in a low-risk group.

2. Materials and methods

2.1. Study design and study subjects

The study was a cross-sectional report performed from April to May 2012 in the Obstetrics and Gynaecology Clinic, Universiti Kebangsaan Malaysia Medical Centre, Bandar Tun Razak, Kuala Lumpur, Malaysia. A total of 139 reproductive-age women (age 19–49 years) were enrolled randomly when they came for follow up at Obstetrics and Gynaecology Clinic, Universiti Kebangsaan Malaysia Medical Centre. Subjects who were virgin, underage and HIV positive were excluded from this study. Subject information was obtained from participants by use of structured questionnaire. They were interviewed in a confidential setting regarding their sociodemographic characteristics, reproductive history, presenting complaints, history of any vaginal discharge and risk behavior. Each subject voluntarily completed the questionnaire before specimen collection and clinical examination.

2.2. Sampling and sample processing

High vaginal swabs were taken by a well-trained medical officer by scraping at the wall of posterior fornix and immediately placing the swab into the Amies gel transport media. Specimens were labeled with reference numbers and tests were conducted anonymously. The specimens were transported in Amies gel agar which has the ability to maintain viability of *T. vaginalis* up to 24 h and transported to culture laboratory on the same day of sample collection for laboratory assessment [12].

Laboratory trainings were given to all staffs prior to the commencement of the study. The specimens were investigated for the presence of *T. vaginalis* using three different methods which were wet mount, Giemsa staining and cultivation in Diamond's medium. The first swab was used to produce a wet mount after mixing with normal saline onto a microscope slide covered with a cover slip for direct microscopic examination. Motile and dead *T. vaginalis* were observed microscopically at a magnification of 400× and 20 fields. In Giemsa staining, the smears were air-dried and fixed immediately with methyl alcohol prior to staining in order to preserve the morphology of the *T. vaginalis*. The smeared glass slides were reviewed by an experienced laboratory technologist for quality control. A third swab specimen was immediately placed in 5 mL of Diamond's medium which was pre-warmed to room temperature and kept in screw-capped tubes prior to use. Cultures were incubated at 37 °C for 7 days and examined using the wet mount method every day in order to detect *T. vaginalis* motile trophozoites.

2.3. Statistical analysis

The data were analyzed using the SPSS software program for Windows version 20 (Chicago, IL, USA). For descriptive analysis, rate (percentage) was used to describe the prevalence of *T. vaginalis*. *P*-values less than 0.05 were considered significant.

2.4. Ethical approvals

The study protocol (Reference No. UKM 1.5.3.5/244/FF-165-2011) was reviewed and approved by the Ethical Committee of Universiti Kebangsaan Malaysia Medical Centre. Informed consent was sought and obtained from individuals prior to participation in this study.

3. Results

Most of the subjects were Malays (76.9%), Chinese (15.1%), other ethnic groups (5.8%) and finally Indians (2.2%). Thirty one percent of them aged between 26 and 31 years, followed by those aged between 32 and 37 years (29.5%). Most subjects involved in this study were young, with a median age of 32 years (interquartile interval of 9.96).

One hundred and thirty eight (99.3%) subjects were married with a stable sexual partner. Almost all of them (98.6%) have less than 6 children in the family. More than half (75.5%) of the women's last child birth was less than 6 years ago. Forty seven percent of them were involved in supporting administrative work, whereas the other half of the subjects were almost equally categorized into professional work (27.3%) and housewife (25.9%). Among the subjects, 90 (64.7%) of them complained of previous or current vaginal discharge. Forty four subjects with vaginal discharge (48.9%) also complained of vaginal itchiness.

Examination of 139 high vaginal swabs using wet mount and Giemsa staining technique were found negative for *T. vaginalis*, whereby culture of 139 high vaginal swabs on Diamond's medium also revealed no growth.

4. Discussion

Trichomoniasis is a well-known sexually transmitted infection in the world and is considerably prevalent in a high-risk

Download English Version:

<https://daneshyari.com/en/article/2032375>

Download Persian Version:

<https://daneshyari.com/article/2032375>

[Daneshyari.com](https://daneshyari.com)