



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb

Document heading

doi:

©2012 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

Antifungal activity of different natural dyes against traditional products affected fungal pathogens

R. Mari selvam¹, A.J.A. Ranjit Singh^{2*} and K. Kalirajan³¹ Department of Zoology, Sri Paramakalyani College, Alwarkurichi, Tamilnadu, India.^{2*} Corresponding Author, Principal, Sri Paramakalyani College, Alwarkurichi, Tamilnadu, India.³ PG & Research Department of Chemistry, Sri Paramakalyani College, Alwarkurichi, Tamilnadu, India.

ARTICLE INFO

Article history:

Received 20 August 2012

Received in revised form 16 September 2012

Accepted 10 December 2012

Available online 28 December 2012

Keywords:

Turmeric

Terminalli

Henna

Guava

Traditional crafts

Humidity

ABSTRACT

Objective: In the present study to evaluate the anti fungal activity of natural dyes against traditional products affected fungal pathogens. **Methods:** Many traditional craft products affected fungal pathogens were isolated using potato dextrose agar medium. The isolated fungus were identified by morphological and microscopically characterization using Alexopolus manual. 50 μ l of Turmeric, Terminalli, Guava and Henna natural dyes were poured into the wells of the culture plates. If antifungal activity was present on the plates, it was indicated by an inhibition zone surrounding the well containing the natural dye. **Result:** At a dose level of 50 μ l of terminalli dye was able to inhibit the growth of all the fungi tested. The absorbance rate of natural dyes analyzed by UV Spectrophotometer. The absorbance rate is high in terminalli (2.266) and turmeric (2.255). **Conclusions:** Natural dyes were bound with traditional products to give good colour and good antimicrobial activity against isolated fungal pathogens

1. Introduction

The various types of traditional craft work being organic origin attack microorganisms due to the carbohydrate content in it. Biological degradation affect the usage, strength utility and value of the products leading to decay and disintegration, splits or cracks, unsightly strains and blotches. So it is essential to treat the raw material and finished products to extend durability and prolong of life. The traditional craft products made from Grass reed leaf and fiber; Bamboo; Areca nut; Palmyra leaf and coconut shell etc. These products are often damaged by fungal biofilm. So they were rejected by exporters and local consumers. So the microbiological intervention is to be applied to prevent the microbial damage in craft products. Many of the fiber products get fungal attack during monsoon seasons with high relative humidity. This fungal attack from a layer over the craft products and consumers reject these products as the fungal infection makes an ugly and dark colour biofilm over the products. This is a major problem the traditional crafts worker endures to address this imperative need, a

microbiological preventive for fungal inhibition is a need of the day.

In the last few decades there has been considerable interest in the active compounds present in natural dyes. In example the major curcuminoid is called curcumin (diferuloyl methane), which makes up approximately 90% of the curcuminoid content in turmeric, followed by demethoxycurcumin and bide methoxycurcumin [20]. Hundreds of in vitro and animal studies have been published describing the antioxidant, anti-inflammatory, antiviral, and antifungal [4, 19] properties of curcuminoids [3, 5, 11, 29, 30]. The curcuminoid give turmeric its bright yellow color. Turmeric oil was tested for its antifungal activity against *Aspergillus flavus*, *Aspergillus parasiticus*, *Fusarium moniliforme* and *Penicillium digitatum* [8]. The natural dye was tested for antimicrobial activity against *E.coli*, *Bacillus subtilis*, *Pseudomonas aeruginosa* and *Styphalococcus aureus* [2]. The extract of henna leaves showed obvious antibacterial activity against *E.coli* [14, 27, 16]. Antibacterial activity of natural dye coated wool fabrics reported [13, 12]. When combined with amphotericin B or Fluconazole, curcumin provided a greater fungicidal effect in the treatment of systemic fungal infections such as candidiasis and candidemia [25, 26]. The natural dyes are using many

*Corresponding author: R.Mari selvam, Department of Zoology, Sri Paramakalyani College, Alwarkurichi, Tamilnadu, India.
Email: honeyselvam01@gmail.com

purposes in example hair dyeing, food preparation and etc [24]. The incorporation of curcumin dye with natural fibers that are used to make handicraft products will add value to the traditional crafts and help to store perishable materials in baskets made of natural fibers. Considerable research work has been undertaken on the application of natural dyes in the coloration of textiles around the globe in the recent past [2, 9, 10, 15, 28].

2. Material and methods

2.1. Preparation of different natural dyes

The four different dyes were prepared using Continuous Stream Distillation Process.

2.1.1. Turmeric dye

1. Turmeric roots were collected from agricultural field at Alwarkurichi region.
2. The roots were dried and powdered. Hundred grams of powdered turmeric roots were boiled in a vessel with 2 liters of water.
3. After 45 min heating the turmeric emitted a strong smell and the water turned a dark orange color.
4. Finally 5ml of lemon juice used as a mordant.

2.1.2. Terminalli dye

1. The Terminalli seeds were dried and powdered and hundred grams of powdered terminalli seeds were boiled in a vessel with 2 liters of water.
2. After 45 min heating the terminalli emitted a strong smell and the water turned a dark brown color.
3. Finally alum used as a mordant.

2.1.3. Guava dye

1. The guava stem outer cuticle collected from the garden and powdered.
2. 100gms of powdered guava cuticle poured in 2 liters of water and boiled with low temperature at 45 minutes.
3. After 45 min, the guava emitted colour. The water turned a gray colour.
4. Finally alum used as a mordant.


2.1.4. Henna dye

1. Henna leaves were collected from agricultural field in our college campus and it to form a paste.
 2. 100gms of henna leaf paste were boiled in a vessel with 2 liters of water.
 3. After 45 min, the paste emitted a strong smell and the water turned a greenish yellow colour.
 4. Finally 5ml of lemon juice used as a mordant.
- After prepared dyes were used for further testing.

2.2. Major groups of antifungal compounds from natural dyes

Plants have an almost limitless ability to synthesize

aromatic substances of different functional groups, most of which are phenols or their oxygen substituted derivatives [6]. Most are secondary metabolites, of which at least 13,000 have been isolated that is less than 10% of the total [21]. In many cases, these substances serve as plant defense mechanisms against predation by microorganisms, insects, and herbivores. Some plants used for their odors (terpenoids), pigment, (quinones and tannins) and flavor (terpenoid capsaicin from chili peppers) were found to be endowed with medicinal properties. Some of the herbs and spices used by humans as season food yield useful medicinal compounds.



The major chemical compound in turmeric is Curcumin, Terflavin B and Chebulinic acid are present in terminallia, Lawsone is active compound of henna and β-Carotene and D & L- Arabinose are present in guava plants. These compounds were good inhibitory effects against fungal pathogens [23, 18].

2.3. Isolation of fungal pathogens

Many of the fiber products (Traditional craft products) are got fungal attacks during monsoon seasons with high relative humidity. The fungal attacked traditional products collected from local traditional craft workers. The collected traditional products (Areca nut plates) were entering to laboratory condition. The fungal pathogens were isolated from collected traditional products using potato dextrose agar medium.

2.4. Fungal identification

The isolated fungus were identified by morphological and microscopically characterization using Alexopolus manual.

2.4.1. Morphological identification

The morphological identification of fungal pathogens using Potato dextrose agar medium and Rose Bengal agar medium. The fungus growth were different from one medium to

Download English Version:

<https://daneshyari.com/en/article/2033645>

Download Persian Version:

<https://daneshyari.com/article/2033645>

[Daneshyari.com](https://daneshyari.com)