



Silencing of miR-124 induces neuroblastoma SK-N-SH cell differentiation, cell cycle arrest and apoptosis through promoting AHR

Tsui-Chin Huang^a, Hsin-Yi Chang^b, Cheng-Yu Chen^b, Pei-Yi Wu^c, Hsinyu Lee^{a,c}, Yung-Feng Liao^d, Wen-Ming Hsu^{e,*}, Hsuan-Cheng Huang^{f,*}, Hsueh-Fen Juan^{a,b,*}

^a Department of Life Science, National Taiwan University, Taipei, Taiwan

^b Institute of Molecular and Cellular Biology, National Taiwan University, Taipei, Taiwan

^c Institute of Zoology, National Taiwan University, Taipei, Taiwan

^d Institute of Cellular and Organismic Biology, Academia Sinica, Taipei, Taiwan

^e Department of Surgery, National Taiwan University Hospital and National Taiwan University College of Medicine, Taipei, Taiwan

^f Institute of Biomedical Informatics and Center for Systems and Synthetic Biology, National Yang-Ming University, Taipei, Taiwan

ARTICLE INFO

Article history:

Received 19 September 2011

Revised 11 October 2011

Accepted 12 October 2011

Available online 20 October 2011

Edited by Tamas Dalmay

Keywords:

Neuroblastoma

MicroRNA

miR-124

Cell differentiation

AHR

ABSTRACT

Neuroblastoma is the most common extracranial solid tumor in children. We investigate whether miR-124, the abundant neuronal miRNA, plays a pivotal role in neuroblastoma. Knockdown of miR-124 promotes neuroblastoma SK-N-SH cell differentiation, cell cycle arrest and apoptosis. Further miR-124 is predicted to target aryl hydrocarbon receptor (AHR) which may promote neuroblastoma cell differentiation. We validate that miR-124 may suppress the expression of AHR by targeting its 3'-UTR. These results suggest that miR-124 could serve as a potential therapeutic target of neuroblastoma.

© 2011 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

1. Introduction

Neuroblastoma, an embryonic tumor derived from the autonomic nervous system neural-crest tissues, is the second most common extracranial malignant tumor of childhood and the most common solid tumor of infancy [1]. Neuroblastoma arises from fetal precursors of the sympathetic nervous system [2]. Molecular

Abbreviations: miRNA, microRNA; AHR, aryl hydrocarbon receptor; 3'-UTR, 3'-untranslated mRNA region; Pre, pre-miR-124 precursor; Inh, anti-miR-124 miRNA inhibitor; Mock-Pre, precursor negative control; Mock-Inh, inhibitor negative control

* Corresponding authors. Addresses: Department of Surgery, National Taiwan University Hospital and College of Medicine, National Taiwan University, No. 7, Chung-Shan South Rd., Taipei 100, Taiwan. (W.-M. Hsu), Institute of Biomedical Informatics and Center for Systems and Synthetic Biology, National Yang-Ming University, No. 155, Sec. 2, Linong Street, Taipei 112, Taiwan. Fax: +886 2 28202508 (H.-C. Huang), Department of Life Science, Institute of Molecular and Cellular Biology, Graduate Institute of Biomedical Electronics and Bioinformatics, National Taiwan University, No. 1, Sec. 4, Roosevelt Rd., Taipei 106, Taiwan. Fax: +886 2 2367 3374 (H.-F. Juan).

E-mail addresses: isurg76@gmail.com (W.-M. Hsu), hsuancheng@ym.edu.tw (H.-C. Huang), yukijuan@ntu.edu.tw (H.-F. Juan).

defects in differentiation are one of the causes in neuroblastoma development.

MicroRNAs (miRNAs) are a class of small non-coding RNAs approximately 21 nucleotides in length that regulate diverse cellular processes including decision of cell lineage, stem cell division, development in evolutionarily divergent organisms, tumorigenesis, and tumor cell metastasis [3,4]. These evidences suggest that miRNAs may be valuable therapeutic agents for their roles in modulating the differentiation of human tumor cells. MiR-124 is a conserved miRNA that has been recorded in 60 organisms for 94 sequences in mirBase and been found to be abundant in nerve system, endocrine gland, and embryo [5]. Most studies showed that this neuronal abundant miRNA could modulate the neurite outgrowth during neurogenesis and regulate the differentiation in neuron. For instance, miR-124 can inhibit SCP1 expression in embryonal carcinoma cell P19 and shows a complementary expression pattern during spinal cord development [6]. Overexpression of miR-124 in P19 cells contributes to the control of neurite outgrowth by regulation of the cytoskeleton remodeling [7]. In the mouse neuroblastoma cell lines, CAD and Neuro2a, miR-124 promotes the differentiation by targeting PTBP1 mRNA encodes a

global repressor of alternative pre-mRNA splicing in nonneuronal cells [8]. During adult neurogenesis in mice, miR-124 is an important regulator of the temporal progression along the subventricular zone stem cell lineage to neurons [9] and during the apical to basal transition of neural precursor cells in the mouse embryonic neocortex [10].

However, the functions of miR-124 in neuronal differentiation remain controversial. Cao et al. showed that experiments to alter expression of miR-124 in neural cells did not appear to affect differentiation in the chick neural tube [11]. In the early developmental stage of optic cup formation in *Xenopus*, miR-124 targeting NeuroD1 plays an anti-neural role in promoting cell proliferation [12]. Furthermore, the role of miR-124 in differentiation shows diversity in a genome wide small RNA sequencing project. Upon differentiation stimulation of retinoic acid (RA), transphorbol ester (TPA), dibutyryl cAMP (dcAMP), or fetal bovine serum (FBS) withdrawn, the expression of miR-124 is upregulated in human SHSY5Y and mouse N1E-115 neuroblastoma cells but downregulated in mouse Neuro2a and NG-108 neuroblastoma cells [13]. Study of dysregulation in miR-124 thus showed decisive in regulation of neuroblastoma.

Aryl hydrocarbon receptor (AHR) is a ligand-activated transcription factor that resides in the cytoplasm as an inactive form by binding of chaperon proteins such as 90 kDa heat shock protein [14,15] and forms heterodimer with AHR nuclear translocator (ARNT) [16] to activate xenobiotic response element located upstream of the target genes such as CYP1A1 due to activation of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD)-related ligands binding and translocation into the nucleus [17,18]. Nevertheless, AHR is postulated to play important roles not only in the regulation of xenobiotic metabolism but also in the regulation of differentiation in pro-inflammatory T cells [19], ketarinocytes [20], myeloblastic leukemia cells [21], and Neuro2a cells [22].

Here in this study, we reveal an undescribed regulation of miR-124 in neuroblastoma that loss of miR-124 resulted in upregulation of AHR leading differentiation in the neuroblastoma cell line SK-N-SH.

2. Materials and methods

2.1. Patients and treatment

The clinical evaluation and use of tumor samples in each patient were approved by the Institutional Review Board of National Taiwan University Hospital, Taipei. During a period of 16 years (from December 1990 to December 2006), tissues were immediately frozen and stored at -80°C prior to real-time RT-PCR analyses.

2.2. Cell culture

Neuroblastoma cell line SK-N-SH and SK-N-BE were maintained in Dulbecco's Modified Eagle's Medium (DMEM)/F12 with 10% (v/v) fetal bovine serum (FBS) (GIBCO, New York, USA). A549 cells were maintained in DMEM. Cells were cultured at 37°C in a humidified atmosphere containing 5% CO_2 .

2.3. miRNA transfection

Cells were seeded into six-well plates and transfected using Turbofect™ in vitro Transfection Reagent (Fermentas, Burlington, USA) according to the manual. Pre-miR-124 precursor (Pre), anti-miR-124 miRNA inhibitor (Inh), precursor negative control (Mock-Pre) and inhibitor negative control (Mock-Inh) were purchased from Ambion Inc. (Austin, TX). Each of them was transfected into cells at a concentration of 200 nM as previous described [23].

2.4. RNA extraction and real-time RT-PCR

Total RNA was extracted from cells or tissues with Trizol reagent according to the instruction (Invitrogen, Carlsbad, USA). The miR-124 level was quantified by quantitative real-time RT-PCR using TaqMan MicroRNA Assay kits (Applied Biosystems, Foster City, USA) with U6 small nuclear RNA (snRNA *RNU6B*) as an internal normalized reference. Their relative levels were measured in triplicate on a Prism 7300 real-time PCR machine (Applied Biosystems). For the quantification of AHR, GAP-43, CRT, and NSE, extracted RNA was reverse transcribed to cDNA using an oligo(dT)12 primer and RevertAid™ Reverse Transcriptase (Fermentas). Their relative levels were measured in triplicate on a MyiQ Single-Color Real-Time PCR Detection System (Bio-Rad, Hercules, USA) with GAPDH as internal control.

2.5. Luciferase reporter assay

The reporter constructs containing AHR 3'-untranslated mRNA region (3'-UTR) were cloned into the pMIR-REPORT vector (Ambion) using PCR-generated fragment. The mutated and deleted AHR 3'-UTR were amplified from the Luc-AHR-WT vector. The reporter assays were carried out as described previously with a transfection control [23].

2.6. Western blot analysis

Total cellular protein was separated by electrophoresis in SDS-polyacrylamide gels before transfer to PVDF membrane (Millipore, Bedford, MA, USA). Anti-GAP43, anti-AHR, and anti-Ki67 were obtained from Abcam (Cambridge, UK), Santa Cruz (CA, USA), and Millipore, respectively. Others were purchased from Cell Signaling Technology (MA, USA). Signals were visualized by chemiluminescence using the ECL reagent (Millipore).

2.7. Statistical analysis

Statistical analysis of miRNA expression in tumor cells was performed on \log_2 transformed fold change data. Student's *t*-test and *P*-values were tested for multiple comparisons by controlling the false discovery rate.

3. Results and discussion

3.1. MiR-124 regulates neuroblastoma cell differentiation

To investigate the role of miR-124 in neuroblastoma, we performed the loss- and gain-of-function of miR-124 in neuroblastoma cells. It was evident that the miR-124 inhibitor induced an increase in the number of cells bearing neurites compared to control cells (Figs. 1A and S1). The changes in cell morphology and neurite length observed in the presence of Inh suggested induction of neuronal differentiation. To determine whether the Inh-induced morphological changes were accompanied by biochemical patterns resembling the neuronal phenotype, we measured the levels of neuronal-specific markers including growth-associated protein 43 (GAP-43), calreticulin (CRT) and neuron-specific enolase (NSE) by qRT-PCR. These three known differentiation markers were all elevated after inhibition of miR-124 (Fig. 1B). In addition, the protein expression level of GAP43 is also strongly increased after inhibition of miR-124 (Fig. 1C). The results suggest that miR-124 may function as a cell fate regulator which modulates cell differentiation in SK-N-SH cells.

Download English Version:

<https://daneshyari.com/en/article/2048715>

Download Persian Version:

<https://daneshyari.com/article/2048715>

[Daneshyari.com](https://daneshyari.com)