

Sialic acid is an essential moiety of mucin as a hydroxyl radical scavenger

Yuki Ogasawara^{a,*}, Tomoyuki Namai^a, Fumihiko Yoshino^b, Masaichi-Chang-il Lee^b,
Kazuyuki Ishii^a

^a Department of Environmental Biology, Meiji Pharmaceutical University, 2-522-1 Noshio, Kiyose, Tokyo 204-8588, Japan

^b Department of Clinical Care Medicine, Division of Pharmacology and Electron Spin Resonance Laboratories, Kanagawa Dental College, Yokosuka, Kanagawa, Japan

Received 10 March 2007; revised 23 April 2007; accepted 23 April 2007

Available online 30 April 2007

Edited by David Lambeth

Abstract In this work, we examined the antioxidant role of mucin, a typical sialic acid containing high-molecular weight glycoprotein. The function of mucin as a hydroxyl radical ($\cdot\text{OH}$) scavenger was characterized using bovine submaxillary gland mucin (BSM). Non-treated BSM effectively protected DNA from the attack of $\cdot\text{OH}$; however, desialylated BSM lost this potential. Moreover, we estimated the scavenging effects of BSM against $\cdot\text{OH}$ generated by UV irradiation of hydrogen peroxide using ESR analysis. Our results indicate that BSM has $\cdot\text{OH}$ scavenging ability and sialic acid in mucin is an essential moiety to scavenge $\cdot\text{OH}$.

© 2007 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

Keywords: Oxidative stress; Mucin; Reactive oxygen species; Sialic acid; Hydroxyl radical

1. Introduction

Hydroxyl radical ($\cdot\text{OH}$) is known as one of the most highly reactive and harmful oxygen derived free radicals in a living organism [1]. This radical reacts with various cellular components including lipid, protein and DNA to oxidatively modify or decompose them [2,3]. It has been proposed that mucus layers lining the respiratory tract and gastrointestinal system may be physiologically important free radical scavenger [4]. The surface epithelium of both the respiratory and gastrointestinal tract is covered by an extracellular and renewable layer of gelatinous mucus. This mucus contains mucin, uric acid, ascorbic acid and reduced glutathione (GSH). Mucins are large, abundant, complicated, filamentous glycoproteins that are present at the interface between many epithelia and their extracellular environments [5,6]. Salivary mucins, which are well

known large glycoproteins act as a functional barrier capable of modulating the untoward effects of the oral environment [7,8]. Halliwell and co-workers showed that when pig gastric and respiratory tract glycoproteins were reacted with systems that generate $\cdot\text{OH}$ there was an increase in thiobarbituric acid reactivity which is indicative of oxidative damage to the carbohydrates [9,10]. Interestingly, recent studies suggest that mucin synthesis is induced by oxidative stress [11,12]. However, there have been no intensive studies to elucidate the relation between mucin and $\cdot\text{OH}$ or other reactive oxygen species (ROS) to date. Therefore, in order to characterize the antioxidant activity of mucin against $\cdot\text{OH}$, we examined the role of sialic acid on the terminal position in the glycoprotein using bovine submaxillary mucin (BSM) under physiological conditions.

2. Materials and methods

2.1. Chemicals

BSM and ascorbic acid were purchased from Sigma Chemical Co. (St. Louis, MO, USA). 1,2-Diamino-4,5-methylenedioxybenzene (DMB), 5,5-dimethyl-1-pyrroline-*N*-oxide (DMPO), 4-hydroxy-2,2,6,6-tetramethyl-piperidine-*N*-oxide (TEMPOL) and hydrogen peroxide (atomic absorption grade) were obtained from Wako Pure Chemical Co. (Osaka, Japan). Sialidase (*Arthrobacter ureafaciens*) was from Nakarai Tesque (Kyoto, Japan). All other reagents were of the highest grade commercially available.

2.2. DNA strand cleavage assay

The DNA-nicking assay was performed according to the method of Kukiela [13,14] with minor modification using pBluescript II SK[−] DNA. Hydroxyl radicals were generated by incubating the following reagent at the indicated final concentrations in 0.5 ml of PBS (pH 7.4) at 37 °C for 20 min: 50 μM H_2O_2 , 5 μM FeCl_3 , 25 μM EDTA, 10 μM ascorbic acid and 0.5 μg of DNA. The iron salt was premixed with EDTA before addition to the reaction mixture and the reaction was started by the addition of ascorbic acid.

2.3. Electron spin resonance measurement

A reaction mixture containing 10 mM DMPO, 0.3% H_2O_2 and test samples in 250 μl was prepared in a 1.5 ml test tube. The mixture was transferred to a cell and illuminated at 365 nm using a PAN UV lamp. After 20 s of UV-irradiation, the DMPO- $\cdot\text{OH}$ spin adduct was measured with a JEOL 200ESR spectrometer (X-Band Microwave Unit, JEOL, Japan). ESR measurement conditions were the following: microwave power 8 mW, microwave frequency 9.45 GHz, modulation amplitude 0.08 mT, time constant 0.03 s, sweep time 1 min, sweep width 5 mT and center field 335.95 mT. The signal intensity of DMPO- $\cdot\text{OH}$ was normalized against Mn^{2+} signal, which was used as an internal standard. The component signals observed in these spectra were identified and quantified as reported [15,16]. The double integrals

*Corresponding author. Fax: +81 424 95 8529.

E-mail address: yo@my-pharm.ac.jp (Y. Ogasawara).

Abbreviations: BSM, bovine submaxillary gland mucin; DMB, 1,2-diamino-4,5-methylenedioxybenzene; ESR, electron spin resonance; DMPO, 5,5-dimethyl-1-pyrroline-*N*-oxide; EDTA, ethylenediamine-tetraacetic acid; HEPES, *N*-(2-hydroxyethyl)-piperazine-*N'*-(2-ethanesulfonic acid); HPLC, high-performance liquid chromatography; LC–MS, liquid chromatography–mass spectrometry; NANA, *N*-acetyl neuraminic acid; NGNA, *N*-glycolyl neuraminic acid; ROS, reactive oxygen species; TEMPOL, 4-hydroxy-2,2,6,6-tetramethyl-piperidine-*N*-oxide

of DMPO-OH experimental spectra were compared with those of a 1 μ M 4-hydroxy-2,2,6,6-tetramethyl-piperidine-*N*-oxide (TEMPOL) sample measured under identical settings to estimate the concentration of spin adduct.

2.4. Preparation of desialylated BSM

Desialylated BSM was prepared by treatment of intact BSM with sialidase (60 mU/ml) in 50 mM acetate buffer (pH 5.4). After 90 min incubation at 37 °C the solution was dialyzed in PBS to remove the hydrolyzed free NANA and NAGA. A control of BSM was treated in 50 mM acetate buffer without sialidase and the solution was dialyzed in PBS. After determination of core protein concentration of BSM in the dialyzed solutions, to confirm the complete desialylation of BSM, the NANA and NGNA contents were determined by the pre-labeling HPLC assay using DMB [17] with acid hydrolysis of all sialic acid after saponification of BSM by treatment with 1.0 M KOH [18].

2.5. Determination of sialic acid in mucin following exposure of \cdot OH

BSM was exposed to hydroxyl radicals generated by incubation of the following reagents at the indicated final concentrations in 0.5 ml of PBS (pH 7.4) at 37 °C for 24 h: (1) 1 mM H_2O_2 , (2) 20 μ M FeCl_3 , 100 μ M EDTA and 50 μ M ascorbic acid. (3) 1 mM H_2O_2 , 20 μ M FeCl_3 , 100 μ M EDTA and 50 μ M ascorbic acid. Residual sialic acid concentrations were determined in all reaction mixtures and filtrates by the HPLC method described above.

2.6. Detection of free sialic acid released from BSM by liquid chromatography–mass spectrometry (LC–MS)

After exposure of BSM to the \cdot OH generation system (1 mM H_2O_2 , 20 μ M FeCl_3 , 100 μ M EDTA and 50 μ M ascorbic acid) for 24 h, the reaction mixture was ultrafiltered. A portion of the filtrate was applied to LC–MS to detect free NANA and NGNA according to the method of Valianpour et al. [19] with minor modification. The HPLC system consisted of a Shimadzu LC/MS (LCMS-2010A, Shimadzu, Kyoto, Japan). The mass spectrometer was used in the negative electrospray ionization (ESI) mode. Sample (10 μ l) was loaded on an Inertsil aminopropyl column (2.1 \times 50 mm; 5 μ m particle size; GL Sciences Inc. Tokyo, Japan).

2.7. Other method

Protein concentrations were determined by the method of Lowry [20] using bovine serum albumin as a standard.

2.8. Statistics

Each experiment was performed at least four times and the results are expressed as means \pm S.D. Data were analyzed by an analysis of variance (ANOVA). Post-hoc comparisons of means between groups were performed using Bonferroni's correction with a significance level of 0.05.

3. Results

3.1. Protection of DNA from \cdot OH damage

As shown in Fig. 1, the effect of BSM on \cdot OH scavenging showed dose dependent behavior. In addition, free NANA and NGNA exerted a slight scavenging effect, however the effect was weak compared to BSM containing the same concentration of sialic acid (Fig. 1A). The DNA protection ability of sialic acids in mucin was equal to or higher than that of GSH and mannitol at same concentration (Fig. 1B).

When 2.67 mg/ml BSM containing 1 mM sialic acid was treated in 50 mM acetate buffer with or without sialidase, the concentration of residual sialic acid in BSM obtained after dialysis was 19.78 μ M (desialylated BSM) or 820 μ M (Control). Thus, we used desialylated BSM in which over 97.5% of sialic acids was removed compared to control BSM. As shown in Fig. 2, BSM treated in acetate buffer has a scaveng-




Fig. 1. Dose–response and efficacy of BSM on DNA damage induced by \cdot OH exposure. Electrophoresis of plasmid using agarose gel (1.0%) performed following exposure to \cdot OH generated by Fenton reaction. Experiments were carried out for 20 min at 37 °C using Fenton reaction mixture (final concentrations in 50 μ l of PBS: 50 μ M H_2O_2 , 5 μ M FeCl_3 , 25 μ M EDTA, 10 μ M ascorbic acid) A: Lane 1, plasmid; lane 2, Fenton reaction mixture plus plasmid; lane 3, Fenton reaction mixture plus plasmid and 0.14 mg/ml BSM containing 50 μ M sialic acid; lane 4, Fenton reaction mixture plus plasmid and 0.27 mg/ml BSM containing 100 μ M sialic acid; lane 5, Fenton reaction mixture plus plasmid and 0.54 μ g/ml BSM containing 200 μ M sialic acid; lane 6, Fenton reaction mixture plus plasmid and 1.08 mg/ml BSM containing 400 μ M sialic acid. lane 7, Fenton reaction mixture plus plasmid and 200 μ M NANA; lane 8, Fenton reaction mixture plus plasmid and 200 μ M NGNA. B: Lane 1, plasmid; lane 2, Fenton reaction mixture plus plasmid; lane 3, Fenton reaction mixture plus plasmid and 1 mM GSH; lane 4, Fenton reaction mixture plus plasmid and 100 μ M GSH; lane 5, Fenton reaction mixture plus plasmid and 1 mM mannitol; lane 6, Fenton reaction mixture plus plasmid and 100 μ M mannitol; lane 7, Fenton reaction mixture plus plasmid and 0.54 mg/ml BSM containing 200 μ M sialic acid; lane 8, Fenton reaction mixture plus plasmid and 0.27 mg/ml BSM containing 100 μ M sialic acid.

ing potential, while these effects were not observed with desialylated BSM.

3.2. Scavenging activity against \cdot OH generated by UV irradiation of hydrogen peroxide

To detail the mechanism of \cdot OH scavenging ability by BSM, we estimated the effect on a transient metal independent \cdot OH generating system. In this experiment, we used completely desialylated BSM and the corresponding control BSM treated in acetate buffer without sialidase. Four characteristic signal lines were observed 20 s after UV irradiation of H_2O_2 with DMPO (Fig. 3A). The effects of BSM and desialylated BSM on DMPO-OH signal intensity were estimated in triplicate at the concentration of 0.16 mg/ml (corresponding to 60 μ M sialic acid in control BSM). As shown in Fig. 3B, BSM significantly inhibited DMPO-OH spin adduct formation, while a significant effect was not observed with desialylated BSM.

3.3. Decrease of sialic acid in mucin by exposure to \cdot OH

Fig. 4 shows the total sialic acid contents (NANA plus NGNA) in the total reaction mixture after incubation with various substrates for \cdot OH generating reactions. Total sialic acid contents in BSM were significantly decreased following exposure to \cdot OH generated by the Fenton reaction, while significant decreases in sialic acid were not observed with iron (III) – ascorbic acid or hydrogen peroxide in the reaction mix-

Download English Version:

<https://daneshyari.com/en/article/2049934>

Download Persian Version:

<https://daneshyari.com/article/2049934>

[Daneshyari.com](https://daneshyari.com)