

# Regulation of Stat3 transcriptional activity by the conserved LPMSP motif for OSM and IL-6 signaling

Wei Sun<sup>a</sup>, Marylynn Snyder<sup>a</sup>, David E. Levy<sup>b</sup>, J. Jillian Zhang<sup>a,\*</sup>

<sup>a</sup> Department of Physiology and Biophysics, Weill Medical College of Cornell University, 1300 York Avenue, New York, NY 10021, USA

<sup>b</sup> Department of Pathology, Kaplan Cancer Center, New York University School of Medicine, New York, New York 10016, USA

Received 19 July 2006; revised 15 September 2006; accepted 25 September 2006

Available online 2 October 2006

Edited by Ivan Sadowski

**Abstract** To achieve maximal transcriptional activity in response to gp130 cytokines, Serine-727 (Ser-727) of Stat3 is phosphorylated. Ser-727 resides in the LPMSP motif, the only conserved sequence among the transcription activation domains of several STATs. We show here that in addition to Ser-727, other residues in this LPMSP motif are also required for Stat3 activity in response to cytokine signaling through regulation of Ser-727 phosphorylation and recruitment of the transcription co-activator CBP/p300 to the promoters of Stat3-target genes for transcription activation. Hence, we have demonstrated a critical role for the whole conserved LPMSP motif in JAK-STAT signaling.

© 2006 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

**Keywords:** Stat3; IL-6; Oncostatin M; Transcription; Serine phosphorylation

## 1. Introduction

Stat3 is one member of the STAT family which is essential for signaling of a wide range of cytokines and growth factors by regulating gene transcription [1]. Stat3 can be activated through tyrosine phosphorylation by a variety of cytokines whose receptors share the gp130 chain, including IL-6, oncostatin M (OSM) and leukemia inhibitory factor (LIF) [2]. Due to its pleiotropic activity, Stat3 plays important roles in many biological processes including oncogenesis, angiogenesis and tumor metastasis [3,4].

The transcriptional activity of Stat3 is mediated by its transcription activation domain (TAD) located in the carboxyl-terminal end of the molecule [5]. The STAT TADs in general can function as independent domains for gene activation by recruiting transcription co-activator CBP/p300 (cAMP-responsive element binding protein [6–10]). The Stat3 TAD contains a conserved sequence motif, LPMSP, also present in the TADs of Stat1 and Stat4 [11]. To achieve maximal transcription activation, the serine residue in the LPMSP motif is also phosphorylated in response to ligand stimulation and required for the maximal transcription activities of Stat1, Stat3 and Stat4 [11,12]. In vivo analyses of S727 of Stat1 and Stat3

using knock-in mouse models showed that S727 plays important physiological roles [13,14]. However, the biochemical function underlying the physiological importance of the phospho-serine and conserved LPMSP motif is not completely understood.

In this report, we demonstrate that other residues in this conserved motif play distinct roles for regulating serine phosphorylation and co-activator recruitment. Our results demonstrate that the whole conserved LPMSP motif, rather than the phospho-S727 alone, is a key regulatory element that controls the maximal transcription activities of STATs.

## 2. Material and methods

### 2.1. Cell culture and antibodies

The STAT3<sup>-/-</sup> mouse embryonic fibroblasts (MEFs) [13] are maintained in DMEM supplemented with 10% bovine calf serum (Hyclone Laboratories Incorporated). Antibodies: anti-phosphoserine-Stat3 and anti-phosphotyrosine-Stat3 (Cell Signaling Technology); anti-Stat3 (Transduction Laboratories); anti-p300 (Santa Cruz Biotechnology). Recombinant human IL-6, IL-6R and mouse OSM were from R&D. Ligand concentrations used were 20 ng/ml IL-6 + 200 ng/ml of soluble IL-6R and 25 ng/ml OSM for lengths of time as indicated in each experiment.

### 2.2. Chromatin immunoprecipitation (ChIP) assay

ChIP assays were performed as previously described [15] with the following antibodies: 2 µg of anti-Stat3 or 2 µg of anti-p300.

### 2.3. Plasmid constructions

Stat3 cDNA was amplified by PCR and subcloned into the R<sub>c</sub>CMV mammalian expression vector (Invitrogen). Specific point mutations in Stat3 were generated with the QuikChange site mutagenesis kit (Stratagene) and confirmed by DNA sequencing.

### 2.4. Transfection experiments

Transient transfections of Stat3<sup>-/-</sup> MEF cells were done with the Lipofectamine 2000 method (Invitrogen). Luciferase assays were performed using the Dual-Luciferase Reporter System (Promega). For generation of stable clones, Stat3<sup>-/-</sup> MEF cells were transfected with the calcium-phosphate transfection kit (Invitrogen) and G418-resistant clones were screened by Western blotting.

### 2.5. Real-time RT-PCR analyses

Total RNAs were prepared using TRIzol (Invitrogen). Real-Time RT-PCR analysis was performed as previously described [16]. Primer pair sequences are C/EBP delta: 5'ATCGACTTCAGCGCTACAT, 5'CTAGCGACAGACCCACAC; c-fos: 5'CCGACCTGCCTGCAAGAT, 5'AATCCAGGGAGGCCACAGA; Tenascin C: 5'TGTGTGCTTCGAAGGCTATG, 5'GCAGACACACTCGTTCTCCA; GAPDH: 5'AGACACCAGTAGACTCCAGG, 5'ACGACCCCTTCATTGACC.

\*Corresponding author. Fax: +1 212 746 6226.

E-mail address: jjz2002@med.cornell.edu (J.J. Zhang).

### 3. Results

To understand the physiological significance of the conserved LPMSP motif for STAT-mediated transcription activation in response to cytokine signaling, we generated a panel of point mutations in the Stat3 LPMSP motif (Fig. 1A). The various forms of Stat3 mutants were transfected into Stat3<sup>-/-</sup> MEF cells and phosphorylation of Y705 and S727 were analyzed by Western blotting. OSM induced strong Y705 phosphorylation in all forms of Stat3 (Fig. 1B, middle panel). For S727 phosphorylation, the level of S727 phosphorylation in wildtype Stat3 increased in response to OSM treatment (Fig. 1B, top panel). Similar increases of S727 phosphorylation were detected in the Stat3L724A mutant and the M726A mutant (Fig. 1B, top panel). Mutation of P725A reduced S727 phosphorylation partially (Fig. 1B, top panel) while Stat3P728A had no detectable level of S727 phosphorylation (Fig. 1B, top panel). These results suggest that only the conserved residue P728 is important for OSM-induced Stat3 S727 phosphorylation.

To further determine the effect of these mutations on the transcriptional activity of Stat3, the Stat3<sup>-/-</sup> MEF cells were transiently transfected with the various forms of Stat3 together with a luciferase reporter containing Stat3 binding sites [13]. Mutations of S727A and P728A reduced Stat3 transcription activity while the Stat3P725A mutant had normal transcription activity compared to the wild type (Fig. 2A). Strikingly, mutations of L724A or M726A significantly reduced the transcription activity of Stat3 in response to OSM (Fig. 2A) even though these mutants had normal S727 phosphorylation (Fig. 1B). The various forms of Stat3 were expressed at similar levels (Fig. 2B). These results demonstrate that in addition to S727, other residues in this LPMSP motif are important for Stat3-mediated gene activation and that S727 phosphorylation alone is not sufficient for Stat3 activity.




Fig. 2. The conserved LPMSP motif is required for Stat3 transcriptional activity for OSM signaling. (A) The various indicated versions of Stat3 plasmids, a Stat3 luciferase reporter [13] and an internal control Renilla luciferase reporter (Promega) were transiently transfected into Stat3<sup>-/-</sup> MEFs. Twenty-four hours after transfection, the cells were either left untreated or treated with OSM for 6 h and harvested for luciferase assays. Results shown are luciferase activities normalized against the internal control and the mean + S.D. of four experiments. (B) The lysates used for luciferase assays in (A) were further analyzed by Western blotting for the expression of the various forms of Stat3 mutants.

To understand the molecular mechanism underlying the defect in transcription activity of these Stat3 mutants, particularly those mutants that have normal S727 phosphorylation but lack transcriptional activity such as the L724A and M726A mutants, we studied the ability of the Stat3 mutants to recruit CBP/p300 to the Stat3 target gene c-fos [15]. In




Fig. 1. Requirement of the conserved Proline-728 for Stat3 S727 phosphorylation in response to OSM. (A) A schematic diagram of Stat3 and the sequence of the transcription activation domain. The conserved LPMSP motif is in bold. TAD, transcription activation domain; Y705, tyrosine-705; S727, serine-727. (B) Stat3<sup>-/-</sup> MEF cells were transiently transfected with Stat3 expression plasmids for 24 h and left untreated or treated with oncostatin M (OSM) for 30 min. Whole cell extracts were analyzed by Western blotting. PS, phospho-S727; PY, phospho-Y705; WT, wild-type; LA, leucine-724 to alanine; SA, serine-727 to alanine; P1A, proline-725 to alanine; MA, methionine-726 to alanine; P2A, proline-728 to alanine.

Download English Version:

<https://daneshyari.com/en/article/2050299>

Download Persian Version:

<https://daneshyari.com/article/2050299>

[Daneshyari.com](https://daneshyari.com)