

# Chloroplast nitrite uptake is enhanced in *Arabidopsis* PII mutants

Sylvie Ferrario-Méry<sup>a,\*</sup>, Christian Meyer<sup>a</sup>, Michael Hodges<sup>b</sup>

<sup>a</sup> Unité de Nutrition Azotée des Plantes, INRA, Route de St. Cyr, 78026 Versailles Cedex, France

<sup>b</sup> Institut de Biotechnologie des Plantes, Unité Mixte de Recherche 8618 Centre National de la Recherche Scientifique, Bât. 630, Université Paris-Sud 11, 91405 Orsay Cedex, France

Received 4 February 2008; revised 22 February 2008; accepted 25 February 2008

Available online 4 March 2008

Edited by Ulf-Ingo Flügge

**Abstract** In higher plants, the PII protein is a nuclear-encoded plastid protein that regulates the activity of a key enzyme of arginine biosynthesis. We have previously observed that *Arabidopsis* PII mutants are more sensitive to nitrite toxicity. Using intact chloroplasts isolated from *Arabidopsis* leaves and <sup>15</sup>N-labelled nitrite we show that a light-dependent nitrite uptake into chloroplasts is increased in PII knock-out mutants when compared to the wild-type. This leads to a higher incorporation of <sup>15</sup>N into ammonium and amino acids in the mutant chloroplasts. However, the uptake differences do not depend on GS/GOGAT activities. Our observations suggest that PII is involved in the regulation of nitrite uptake into higher plant chloroplasts. © 2008 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

**Keywords:** *Arabidopsis*; *AtGLBI* gene; Chloroplasts; NO<sub>2</sub><sup>-</sup> uptake; PII protein; T-DNA insertion mutant

## 1. Introduction

PII is a highly conserved protein that plays a role in the sensing of carbon/nitrogen (C/N) balance and energy status in bacteria, cyanobacteria and plants. It has been shown to be involved in the regulation of inorganic nitrogen uptake and assimilation in both bacteria and cyanobacteria. The PII protein interacts with various target proteins that include signal transduction proteins, key metabolic enzymes and metabolite transporters (see [1–3] for review). In cyanobacteria, NtrC, a subunit of a nitrate/nitrite transporter is believed to be regulated by PII [4]. In all known cases, PII – target protein interactions depend on the Mg-ATP and 2-oxoglutarate (2-OG) levels in the cell [1]. In photosynthetic cyanobacteria and eubacteria, PII activity is also modified by phosphorylation or uridylylation, respectively [5].

In higher plants, a single PII homolog (*GLBI*) has been identified in *Arabidopsis*, rice, tomato, castor bean, alfalfa, and pine [6–8]. PII is a nuclear-encoded chloroplastic protein displaying 50% identity to bacterial PII proteins. As in bacterial systems, plant PII can bind 2-oxoglutarate in the presence of Mg-ATP [9]. However, there is no evidence for PII phosphor-

ylation albeit the conservation of the phosphorylated serine residue found in *Synechococcus* PII [10]. To date, the only PII target protein discovered in plants is conserved between higher plants and cyanobacteria. This PII interacting protein is the chloroplastic *N*-acetyl glutamate kinase (NAGK) that is activated in vitro and shows a reduced retro-inhibition by arginine when the NAGK-PII complex is formed [11,12]. In the leaves of PII knock-out mutants grown under ammonium nutrition conditions, arginine, ornithine and citrulline levels were reduced by 50%, thus giving the first physiological evidence of the activation of NAGK by PII in planta [11].

The identification of new PII targets in plants is essential to understand the role of this protein and the use of PII mutants should be useful to decipher PII function. Indeed, *Arabidopsis* PII mutants appeared to be more sensitive to NO<sub>2</sub><sup>-</sup> toxicity, as judged by the mortality of in vitro grown seedlings [13]. Interestingly, this observation suggests that PII might be involved in the regulation of NO<sub>2</sub><sup>-</sup> metabolism. This function could be reminiscent of the role of PII in cyanobacteria where it has been shown to regulate a nitrate/nitrite transporter [4]. In higher plants, NO<sub>2</sub><sup>-</sup> is synthesised in the cytosol by reduction of NO<sub>3</sub><sup>-</sup> by nitrate reductase (NR). NO<sub>2</sub><sup>-</sup> is then translocated to the chloroplast to be further reduced into ammonium by the nitrite reductase. It is noteworthy that NO<sub>2</sub><sup>-</sup> does not accumulate in plant tissues. This could be partly due to the regulation of NR activity by chloroplastic redox poise and the oxaloacetate/malate shuttle that would deliver reductant for NR activity. In addition, an inhibition of NR activity by phosphorylation and subsequent binding of 14-3-3 proteins occurs in the dark, thus avoiding NO<sub>2</sub><sup>-</sup> accumulation in the dark when NADH could be limiting [14]. However, the occurrence of a highly efficient transport system across the chloroplast envelope to avoid NO<sub>2</sub><sup>-</sup> toxicity is conceivable. However, the molecular mechanism of NO<sub>2</sub><sup>-</sup> uptake into chloroplasts is largely unknown in higher plants. It is possible that NO<sub>2</sub><sup>-</sup> is transported across the plastid membrane by permeation of the nitrous acid [15] versus a NO<sub>2</sub><sup>-</sup> channel or a NO<sub>2</sub><sup>-</sup> transporter [16,17]. However the uptake of NO<sub>2</sub><sup>-</sup> into intact pea chloroplasts shows saturable kinetics thus supporting the presence of a NO<sub>2</sub><sup>-</sup> transport system [16,17]. In *Chlamydomonas*, *Nar1-1* encodes a chloroplast membrane protein involved in high affinity NO<sub>2</sub><sup>-</sup> transport [18,19]. It is a member of the formate/nitrite transporter family [19], however no orthologs have been hitherto identified in the *Arabidopsis* genome. Recently, a NO<sub>2</sub><sup>-</sup> transporter has been identified in higher plants [20]. It is a member of the proton-dependent oligopeptide transporter family but it displays little homology with the NO<sub>2</sub><sup>-</sup> transporters described in cyanobacteria and *Chlamydomonas*.

\*Corresponding author. Fax: +33 130833096.  
E-mail address: ferrario@versailles.inra.fr (S. Ferrario-Méry).

**Abbreviations:** Col, Colombia ecotype; Chl, Chlorophyll; DCMU, 3-(3,4-dichlorophenyl)-1,1-dimethylurea; MSO, methionine sulfoximine; WS, Wassilewskija ecotype

In this study we have tested the hypothesis that PII has a role in the regulation of  $\text{NO}_2^-$  uptake into higher plant chloroplasts, reminiscent of its role in cyanobacterial  $\text{NO}_2^-$  uptake [4]. Using intact chloroplasts isolated from *Arabidopsis* leaves and  $^{15}\text{N}$ -labelled nitrite we show that a light-dependent  $\text{NO}_2^-$  uptake into chloroplasts is increased in PII knock-out mutants when compared to the wild-type.

## 2. Materials and methods

### 2.1. Plants and growth conditions

PIIV1 and PIIS2 mutant lines of *Arabidopsis* Wassilewskija ecotype (WS) and Colombia ecotype (Col) have been described previously and contain undetectable amounts of the PII protein [13]. Wild-type plants and PII mutants were grown on sterilised compost and watered daily with a complete nutrient solution containing 10 mM  $\text{NO}_3^-$  and 2 mM  $\text{NH}_4^+$  [21]. Plants were grown in a growth chamber at 60% relative humidity, a light intensity of  $150 \mu\text{mol m}^{-2} \text{s}^{-1}$  and a day–night regime of 8 h at 22 °C and 16 h at 20 °C, respectively. The total rosette of 6 week-old plants was used for intact chloroplast isolation. Plants were maintained in darkness during 24 h before being harvested in order to reduce chloroplastic starch levels.

### 2.2. Chloroplast purification

Chloroplasts were obtained from 40 g *Arabidopsis* leaves of wild-type and PII mutant plants. Leaves were washed in distilled water at 4 °C for 20 min and then ground in a blender for 2 s in a 20 mM Tricine KOH (pH 8) homogenization buffer maintained at 4 °C and containing 450 mM sorbitol, 10 mM EDTA, 10 mM  $\text{NaHCO}_3$  and 0.1% BSA (100 ml/10 g of leaves). The homogenates were immediately filtered through several layers of gauze and centrifuged at  $700 \times g$  for 5 min at 4 °C. The pellets were resuspended in 3 ml buffer (RB) containing 20 mM Tricine KOH (pH 7.6), 300 mM sorbitol, 2.5 mM EDTA and 5 mM  $\text{MgCl}_2$ . The resulting solution was transferred to the top of a 40% percoll solution in RB (30 ml total volume) and then centrifuged for 3 min at  $2500 \times g$  using a swing out centrifuge bucket at 4 °C. Intact chloroplasts were pelleted while broken chloroplasts and other membrane debris remained at the top of the percoll solution. The chloroplast suspensions were diluted (3-fold) in RB and centrifuged for 4 min at  $5180 \times g$  at 4 °C and the pellets were resuspended in 1–2 ml of RB. The chlorophyll (Chl) content of the chloroplast suspension was calculated from the absorption at 652 nm after extraction of an aliquot (5  $\mu\text{l}$ ) in 80% acetone [22]. The chloroplast suspensions were stored at 4 °C in the dark until their use for  $\text{NO}_2^-$  uptake measurements.

Chloroplast intactness was determined by measuring the light-dependent oxygen production of ‘intact’ and osmotically broken chloroplasts using an  $\text{O}_2$  electrode and ferricyanide (as an electron acceptor) [23]. The intactness was estimated by the ratio  $[(A - B)/A \times 100]$  where  $A$  is the ferricyanide-dependent  $\text{O}_2$  evolution measured after the osmotic shock and  $B$  the  $\text{O}_2$  evolution measured before the osmotic shock.

### 2.3. $\text{NO}_2^-$ uptake by isolated chloroplasts

$\text{NO}_2^-$  uptake by chloroplasts was measured in a ‘plexiglass cell’ (maximum volume of 3 ml), with gentle agitation, at 20 °C and either a light intensity of  $50 \mu\text{mol m}^{-2} \text{s}^{-1}$  or in darkness. The chlorophyll concentration of the chloroplast suspension in the ‘cell’ was adjusted to  $0.1 \text{ mg ml}^{-1}$  in 2 ml of RB. Chloroplasts in the measuring cell were either illuminated 5 min before adding  $^{15}\text{NO}_2\text{Na}$  (light treatment) or maintained in the darkness (dark treatment). The final concentration of  $^{15}\text{NO}_2\text{Na}$  was 0.25 mM (98%  $^{15}\text{N}$  enrichment) for the uptake studies or 0.1–2.5 mM for the determination of kinetic parameters as indicated in the text. Aliquots (300  $\mu\text{l}$ ) of the chloroplast suspension were collected at the times indicated in the text (from 0 to 30 min) and immediately centrifuged. The supernatant was stored at –20 °C until the measurement of nitrite. The pellets were rinsed in 300  $\mu\text{l}$  of RB and centrifuged. The rinsed pellets were stored at –80 °C for  $^{15}\text{N}$  total enrichment analysis. The inhibitor studies were carried out using 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) (50  $\mu\text{M}$ ), methio-

nine sulfoximine (MSO, 1 mM) and azaserine (1 mM). They were added 5 min before  $^{15}\text{NO}_2\text{Na}$  addition. WS and PII mutant chloroplasts were measured simultaneously for each experiment. For the studies of the effect of the addition of  $\text{NH}_4^+$  (1 mM) and  $\alpha\text{KG}$  (1 mM), these metabolites were added 5 min before  $^{15}\text{NO}_2$  addition (as described in [24]).

### 2.4. $\text{NO}_2^-$ measurement

$\text{NO}_2^-$  was measured colorimetrically at 540 nm using *N*-(1-Naphtyl) ethylene diamine dichlorhydrate as in [25].

### 2.5. $^{15}\text{N}$ labelling measurements

For the determination of total  $^{15}\text{N}$  content, the pellets were dried at 70 °C for 48 h, weighed, and analyzed using a continuous-flow isotope ratio mass spectrometer coupled to a carbon/nitrogen elemental analyzer (model ANCA-MS, PDZ Europa, Crewe, UK), as described in [26]. For the  $^{15}\text{N}$  labelling of  $\text{NH}_4^+$  and amino acids in the pelleted chloroplasts, the amino acid extracts from (SSA extracts) were filtered and analysed as described in [27,28].

### 2.6. Transcriptome analyses

The transcriptome analyses were performed with CATMA arrays containing 24576 Gene Specific Tags from *Arabidopsis thaliana* as described in [11]. Two biological repetitions of the same experiment were performed and used for the calculations. The results of the microarray data analysis are shown as the ratio of the intensities for each gene in the two genotypes analysed (after the recalculation from the initial data expressed as a log<sub>2</sub> ratio provided by the CATMA analysis).

## 3. Results

### 3.1. $\text{NO}_2^-$ uptake into isolated chloroplasts is increased in PII mutants

Previous studies suggested PII to have a role in nitrite metabolism or transport due to the increased  $\text{NO}_2^-$  sensitivity observed in PII mutant seedlings grown in vitro on  $\text{NO}_2^-$  as the sole nitrogen source [13]. Since, cyanobacterial PII has been shown to modulate the activity of a  $\text{NO}_3^-/\text{NO}_2^-$  transporter [4], it was decided to investigate  $\text{NO}_2^-$  uptake into chloroplasts isolated from rosette leaves of PII mutants and wild-type plants. The yield of the isolated intact chloroplasts was verified by ferricyanide-dependant  $\text{O}_2$  production measured using an oxygen electrode before (isolated chloroplasts) and after (broken chloroplasts) an osmotic shock. The intactness of the chloroplasts isolated from *Arabidopsis* rosette leaves was found to be between 50% and 80% for the wild-type ecotypes (WS and Col) and the PII mutant lines (PIIV1 and PIIS2) (data not shown). Broken chloroplasts did not exhibit any light-dependent nitrite uptake, and the measured transport rates appeared to be correlated with the degree of chloroplast intactness (data not shown).

$\text{NO}_2^-$  uptake into isolated chloroplasts was estimated by measuring the disappearance of  $\text{NO}_2^-$  from the incubation medium or by  $^{15}\text{N}$  (N total) labelling in the chloroplast pellets after different dark or light treatments. There was a good correlation between the two methods as seen in Fig. 1. Therefore, for further analyses it was decided to measure  $\text{NO}_2^-$  uptake by the disappearance of  $\text{NO}_2^-$  in the chloroplast incubation medium. The effect of  $\text{NO}_2^-$  concentration on the observed light-dependent  $\text{NO}_2^-$  uptake was measured between 0.1 and 2.5 mM  $^{15}\text{NO}_2\text{Na}$ . Uptake was saturated above 0.5 mM  $^{15}\text{NO}_2\text{Na}$  for both the wild-type and PII mutant genotypes (Fig. 2) and a  $K_m$  of approximately 200  $\mu\text{M}$   $\text{NO}_2^-$  was obtained in agreement with previous observations [16,20].

Download English Version:

<https://daneshyari.com/en/article/2050945>

Download Persian Version:

<https://daneshyari.com/article/2050945>

[Daneshyari.com](https://daneshyari.com)