

Bradykinin activates ADP-ribosyl cyclase in neuroblastoma cells: Intracellular concentration decrease in NAD and increase in cyclic ADP-ribose

Haruhiro Higashida^{a,b,c,*}, Alla Salmina^d, Minako Hashii^a, Shigeru Yokoyama^a, Jia-Sheng Zhang^a, Mami Noda^e, Zen-Guo Zhong^f, Duo Jin^a

^a Department of Biophysical Genetics, Kanazawa University Graduate School of Medicine, 13-1 Takara, Kanazawa, Ishikawa 920-8640, Japan

^b Department of Cellular Neurophysiology, Kanazawa University Graduate School of Medicine, Kanazawa 920-8640, Japan

^c The Kanazawa University 21st Century COE Program on Innovative Brain Science for Development, Learning and Memory, Japan

^d Department of Biochemistry and Medical Chemistry, Krasnoyarsk State Medical Academy, Krasnoyarsk 660022, Russia

^e Department of Pathophysiology, Graduate School of Pharmaceutical Sciences, Kyushu University, Fukuoka 812-8582, Japan

^f Senile Brain Disease Institute, Guangzhou University of Traditional Chinese Medicine, China

Received 26 June 2006; revised 27 July 2006; accepted 31 July 2006

Available online 8 August 2006

Edited by Jesus Avila

Abstract ADP-ribosyl cyclase activity in the crude membrane fraction of neuroblastoma × glioma NGPM1-27 hybrid cells was measured by monitoring [³H] cyclic ADP-ribose (cADPR) formation from [³H] NAD⁺. Bradykinin (BK) at 100 nM increased ADP-ribosyl cyclase activity by about 2.5-fold. Application of 300 nM BK to living NGPM1-27 cells decreased NAD⁺ to 78% of the prestimulation level at 30 s. In contrast, intracellular cADPR concentrations were increased by 2–3-fold during the period from 30 to 120 s after the same treatment. Our results suggest that cADPR is one of the second messengers downstream of B₂ BK receptors.

© 2006 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

Keywords: Cyclic ADP-ribose; NAD; ADP-ribosyl cyclase; Bradykinin

1. Introduction

Bradykinin (BK) plays an important role in blood pressure control, inflammation, edema, pain, and neuronal signaling [1]. Recently, BK was shown to be involved in the pathogenesis of particular conditions, such as cardiovascular diseases and glomerular injury in diabetes [2], prostate cancer [3], breast cancer [4], tumor-associated angiogenesis [5], and Alzheimer's disease [6]. These physiological and pathophysiological effects of BK are exerted through BK receptors, B₁ and B₂ [1]. Stimulation of BK receptors leads to activation of phospholipase C

(PLC) [1,7], focal adhesion kinase [8], Pyk2 [9], protein kinase C [10], and Ras or mitogen activated protein kinase [1,9]. BK also activates Ca²⁺ processes due mainly to intracellular Ca²⁺ mobilization by inositol-1,4,5-trisphosphate (InsP₃) from the endoplasmic reticulum [1,7,10].

Ca²⁺ mobilization is triggered not only by InsP₃ but also by Ca²⁺ itself, i.e. Ca²⁺-induced Ca²⁺ release (CICR) via ryanodine receptors [11]. CICR is co-activated by a putative second messenger, cyclic ADP-ribose (cADPR) [11–13]. cADPR is synthesized from β-NAD⁺ by both membrane-bound and cytosolic ADP-ribosyl cyclases in mammalian tissues, including the nervous system [14]. The cADPR synthetic activity is regulated by receptor stimulation with several different mechanisms [12–14]. In one such mechanism, ADP-ribosyl cyclase seems to be coupled directly with neurotransmitter or hormone receptors via different G proteins on the membrane surface [15]. However, the same control of ADP-ribosyl cyclase by BK receptors has not been reported previously. Recently, it was demonstrated that BK increased intracellular calcium, nitric oxide, and cADPR levels, and ADP-ribosyl cyclase activity in coronary arterial endothelial cells subsequently causing smooth muscle dilatation [16]. If this is true, it should be proved that BK decreases intracellular NAD⁺ levels as a consequence of ADP-ribosyl cyclase activation. To address this question, we used a neuronal model cell line of NGPM1-1 neuroblastoma × glioma hybrid cells, expressing endogenous B₂ BK receptors and exogenous M1 muscarinic acetylcholine receptors (mAChRs) [17,18]. We measured ADP-ribosyl cyclase activity in crude membrane fractions of NGPM1-27 cells and showed changes in the content of [³H] NAD⁺ and [³H] cADPR in reaction mixtures. Furthermore, to confirm the signaling from B₂ to ADP-ribosyl cyclase in vivo, the substrate and product concentrations, i.e., intracellular NAD⁺ and cADPR levels, were measured before and after application of BK onto NGPM1-27 cells.

2. Materials and methods

2.1. Membrane preparation

NGPM1-27 cells were cultured as described previously [17]. The cells harvested were suspended in 10 mM Tris–HCl solution, pH 7.3, with 5 mM MgCl₂ at 4 °C for 30 min. The suspension was homogenized

*Corresponding author. Address: Department of Biophysical Genetics, Kanazawa University Graduate School of Medicine, 13-1 Takara, Kanazawa, Ishikawa 920-8640, Japan. Fax: +81 76 234 4236. E-mail address: haruhiro@med.kanazawa-u.ac.jp (H. Higashida).

Abbreviations: BK, bradykinin; PLC, phospholipase C; InsP₃, inositol-1,4,5-trisphosphate; CICR, Ca²⁺-induced Ca²⁺ release; cADPR, cyclic ADP-ribose; mAChR, muscarinic acetylcholine receptor; ADPR, ADP-ribose; ACh, acetylcholine; CCh, carbamoycholine; [β-NAD⁺]_i, intracellular β-NAD⁺ concentrations; [cADPR]_i, intracellular cADPR concentrations

in a glass homogenizer. The resultant homogenate was centrifuged at 4 °C for 5 min at 1000 × *g* to remove unbroken cells and nuclei. Crude membrane fractions were prepared by centrifugation (twice) of homogenates at 105 000 × *g* for 15 min. The supernatant was removed, and the final pellet was dispersed in 10 mM Tris–HCl solution, pH 7.0. In each experiment, membranes were freshly prepared and used immediately for enzymatic reactions.

2.2. ADP-ribosyl cyclase assay

Each 20- μ l reaction mixture contained 50 mM Tris–HCl (pH 7.0); 100 mM KCl; 10 μ M CaCl₂; 2 μ M β -NAD⁺; 0.1 μ M β -[2,8 adenine-³H] NAD⁺ (0.06 μ Bq) and 0.40–7.16 μ g of membrane proteins, according to a formula reported previously [15]. Reaction mixtures were incubated for 0.5–4 min at 37 °C. Reactions were stopped by adding 2 μ l of 10% trichloroacetic acid, and aliquots were centrifuged for 1 min at 2100 × *g*, and 2 μ l of the supernatant were spotted on silica gel plastic thin layer sheets (20 × 10 cm). The layers were developed in the ascending direction for 40–70 min at 23 °C with a mixture of water/ethanol/ammonium bicarbonate (in the ratio 30%: 70%: 0.2 M or 36%: 64%: 0.3 M). The positions of authentic cADPR, ADP-ribose (ADPR) and β -NAD⁺ were detected by UV illumination and of [³H]-labeled products were autoradiographically confirmed in each by Fuji Bas 1000 (Tokyo, Japan). Corresponding areas (about 1 × 0.7 cm) were cut out and the radioactivity was counted in a liquid scintillation counter.

2.3. Intracellular NAD⁺ content

NGPM1-27 cells were cultured on polyornithine-coated dishes (35-mm in diameter) for 4 days. The NAD⁺ content in the supernatant of the heat-inactivated cell homogenate was determined by a slight modification of an enzyme cycling method described as reported previously [15].

2.4. Cycling assay for cADPR

Intracellular cADPR concentrations were measured according to the enzyme cycling assay method described by Graeff and Lee [19]. Briefly, NGPM 1-27 cells were cultured in polyornithine-coated 35-mm dishes. The cells were extracted with 100 μ l of 0.6 M perchloric acid at 4 °C. In order to observe the agonist effect on cADPR levels, agonists were applied on cells in dishes with serum-containing growth medium with no prior change to a fresh experimental medium without serum.

3. Results

3.1. Effects of bradykinin on ADP-ribosyl cyclase activity in NGPM1-27 cells

[³H] cADPR and [³H] ADPR were produced from β -[³H] NAD⁺ by preparation of the crude membrane fraction of NGPM1-27 cells. During an incubation period of 4 min, the majority of radioactivity of β -NAD⁺ was converted to ADPR and/or cADPR, as shown in Fig. 1. The accumulation of radioactivity was greater in the spot of ADPR than cADPR (Fig. 1A). The average specific activity of ADP-ribosyl cyclase, as the rate of [³H] cADPR formation, was 204 ± 43 pmol/min/mg protein (mean ± S.E.M., *n* = 27).

Addition of 100 nM BK to the reaction mixture at zero time increased the rate of [³H] cADPR formation to a greater extent than [³H] ADPR production (Fig. 1C). The average activation by 100 nM BK was 248 ± 47% (*n* = 4) of the control activity (Student's *t* test, *P* < 0.01).

Next, we confirmed the response to muscarinic receptors. A similar level of stimulation (321 ± 41% (*n* = 4)) by 1 μ M carbamylcholine (CCh) of ADP-ribosyl cyclase was obtained (321 ± 41%, *n* = 3; Fig. 1B and C).

3.2. BK-induced decrease in intracellular β -NAD⁺ concentration

To confirm the above effects of BK *in vivo*, we examined agonist-stimulated changes in substrate levels (Fig. 2). Fig. 2A shows the time course of changes in intracellular β -NAD⁺ concentrations ([β -NAD⁺]_i) in NGPM1-27 cells challenged with 300 nM BK. [β -NAD⁺]_i was significantly decreased for 15–60 s after application of BK. The decrease in [β -NAD⁺]_i at 30 s was 77.3 ± 3.0% (*n* = 9, *P* < 0.01) of the pre-stimulation level, and showed partial recovery at 120 s.

3.3. Intracellular cADPR concentrations in NGPM-1 cells

Intracellular cADPR concentrations ([cADPR]_i) were measured by enzyme recycling assay. [cADPR]_i in NGPM 1-27




Fig. 1. Time course of ADP-ribosyl cyclase activity in NGPM1-27 cell membranes. Aliquots (20 μ l) were withdrawn at the indicated times from 140- μ l reaction mixtures containing membrane protein from NGPM1-27 cells in the absence (A) and presence (B) of 1 μ M CCh. Radioactivity in spots migrating with authentic NAD⁺ (open squares), cADPR (closed square), and ADPR (open diamond) on thin-layer chromatogram sheets was measured. Values are the means of two determinations from one representative of three experiments giving similar results. (C) Time course of changes in ADP-ribosyl cyclase activity (rate of formation of [³H] cADPR) in membranes prepared from NGPM1-27 cells. Reaction mixtures were incubated with or without (open circle) 100 nM BK (closed square) or 1 μ M CCh (closed diamond) for the indicated times. Values are the means of two determinations from one representative of three experiments giving similar results.

Download English Version:

<https://daneshyari.com/en/article/2052484>

Download Persian Version:

<https://daneshyari.com/article/2052484>

[Daneshyari.com](https://daneshyari.com)