

Antiviral and Immunostimulant Activities of *Andrographis paniculata*

CHURIYAH^{1*}, OLIVIA BUNGA PONGTULURAN¹, ELRADE ROFAANI¹, TARWADI²

¹Center of Pharmaceutical and Medical Technology, Agency for the Assessment and Application of Technology,
Building 610 LAPTIAB, Puspiptek Area, Serpong–Tangerang 15314, Indonesia

²Biotech Center, Agency for the Assessment and Application of Technology, Building 630, Puspiptek Area,
Serpong–Tangerang 15314, Indonesia

Received March 26, 2014/Accepted November 27, 2014

Andrographis paniculata (Burm. f.) Nees is a medicinal plant which was reported to have anti HIV, anti pathogenic bacteria and immunoregulatory activities. The research purpose was to investigate the activity of *Andrographis paniculata* ethanol extract as antiviral and immunostimulant. *A. paniculata* leaves oven-dried, then grinded and macerated with ethanol 90%, and the extract then analyzed using High Performance Liquid Chromatography (HPLC) to determine the content of active compounds andrographolide. The antiviral activity of the extract was determined by observing its ability on inhibiting virus load in A549 cells transfected with Simian Retro Virus (SRV) by Real Time – Polymerase Chain Reaction (RT-PCR) analysis. The immunostimulant activity of extract was determined by its ability to induce lymphocytes cell proliferation using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. Our result indicated that the *A. paniculata* ethanol extract inhibited the SRV virus titer similar to the positive control Lamivudine, and it was not toxic to the A459 cell line. Furthermore, low concentration (1 µg/mL) of *A. paniculata* extract could stimulated lymphocyte cell proliferation about 38% compared to the control lymphocyte cell without any treatment.

Keywords: *Andrographis paniculata*, anti-viral, immunostimulant

INTRODUCTION

Andrographis paniculata (Burm. f.) Nees (Acanthaceae) is a medicinal plant used in many countries. Major constituents of *A. paniculata* are diterpenoids, flavonoids and polyphenols. Among the single compounds extracted from *A. paniculata*, andrographolide is the major compound in terms of bioactive properties and abundance.

Methanolic extracts of *A. paniculata* as well as chloroform and hexane extracts were reported to inhibit the growth of bacterial and fungal pathogens (Bobbarala *et al.* 2009). The aqueous extract of *A. paniculata* against anti-HIV was ruled out by testing the inhibitory activities against HIV in the H9 cell line (Chao & Lin 2010). A phase I dose-escalating clinical trial of andrographolide in HIV positive patients reported a significant rise in the mean CD4⁺ lymphocyte level of HIV patients. Andrographolide inhibited HIV-induced cell cycle dysregulation, leading to a rise in CD4⁺ lymphocyte levels in HIV-1 infected individuals (Calabrese *et al.* 2000). Andrographolide, neoandrographolide and 14-deoxy-11,12-didehydroandrographolide

demonstrated viricidal activity against herpes simplex virus 1 (HSV-1) (Wiat *et al.* 2005). The *A. paniculata* ethanol extract and andrographolide inhibit the expression of Epstein-Barr virus (EBV) (Lin *et al.* 2008). A recent *in vitro* study investigated the anti-influenza activity of *A. paniculata* in canine kidney cell line infected with H1N1, H9N2, or H5N1 (Chen *et al.* 2009). Andrographolide and their derivatives displayed anti-HIV activity *in vitro* by inhibiting gp120-mediated cell fusion of HL2/3 cells (expressing gp120 on its surface) with TZM-bl cells (expressing CD4 and co-receptors CCR5 & CXCR4) (Uttekar *et al.* 2012).

Many of andrographolide analogues indicating immunostimulatory properties (Chao & Lin 2010). Andrographolide was reported to have immunoregulatory activities with different effects in different immune disease models, playing a role as a modulator of altered immune responses. In tumor-bearing mice it enhanced natural killer cell activity, increased secretion of IL-2 and IFN- γ by T cells and thereby inhibited the tumor growth. However, in autoimmune encephalomyelitis mice, it interfered maturation of dendritic cells, induced antigen-specific tolerance and thus prevented detrimental autoimmune responses (Wang *et al.* 2010). *In vivo* study using

*Corresponding author. Phone/Fax: +62-21-7560536,
E-mail: churiyah_zafrollah@yahoo.com

mice model have shown that *A. paniculata* was a potent stimulator of immune system in two ways, namely the antigen specific response that antibodies was made to counteracted invading microbes, and non specific immune response that macrophage cell scavenged and destroyed invader (Kumar *et al.* 2011).

The aim of our research was to investigate the activity of the ethanol extract of *A. paniculata* as antiviral and immunostimulator. The activity of extract as immunostimulant was determined on inducing lymphocytes cell proliferation using MTT assay, while as antiviral was observing its ability on inhibiting SRVirus titer in A549 cells transfected with SRV by RT-PCR.

MATERIALS AND METHODS

Preparation of Ethanol Extracts of *A. paniculata*. Simplicia *A. paniculata* leaves were oven-dried, then grinded with a grinder until becomes a fine powder. Furthermore, the powders was extracted by maceration with ethanol 90% and stirred for 24 h. The macerate then filtered with a vacuum pump and evaporated with a rotary evaporator at 40 °C to obtain a viscous extract. The ethanol extract then analyzed using High Performance Liquid Chromatography (HPLC) to determine the content of active compounds andrographolide. About ten milligrams of extract was dissolved in 10 mL of methanol and then filtered with a 0.45 µm pore size of 13 mm diameter membrane. The HPLC was setting by the linear gradient system from 70:30 to 100:0 then 70:30 with methanol:distillate water eluent, with a flow rate of 1 mL/min for 20 min and analyzed at the wavelength of 254 nm.

Isolation and Proliferation of Peripheral Blood Lymphocyte. Blood samples were taken aseptically from healthy donors and stored in sterile tubes which already contained 0.1% EDTA as an anticoagulant. Blood samples then aseptically transferred into a sterile centrifuge tube and centrifuged for 5 min at 214 g. Heavier part of the blood (red blood cells) will be at the bottom and the plasma will be on top.

The plasma was removed and placed into the new sterile conical tube then diluted with 50% Roswell Park Memorial Institute (RPMI) 1640 basal medium, mixed gently, then layered onto Ficoll histopaque. After centrifuged at 594 g for 30 min, the white range buffy coat collected into the new conical tube contained the basal medium, and centrifuged at 95 g for 50 min, then washed twice with the basal medium. The lymphocyte cells were resuspended in RPMI 1640 medium with 10% heat inactivated fetal bovine serum (FBS) and 1% Pen-Strep antibiotic (100 Units/mL Penicillin and 100 µg/mL Streptomycin).

Proliferation of lymphocyte cells was measured using MTT method. The yellow color of MTT changed to purple by mitochondrial succinate dehydrogenase of living cells. Briefly, 5×10^5 cells per well in a 96-well plate were stimulated with different concentration of extract (Table 1 for the details of extract concentration) and mitogen as positive control which were 5 µg/mL of Phytohaemagglutinin (PHA) and Concanavaleine A (ConA). Control wells received only medium without extract or mitogen addition. After 72 h of culturing at 37 °C in a humidified incubator with 5% CO₂, 10 µL of 5 mg/mL MTT was added to each well. After incubation at 37 °C with 5% CO₂ for 4 h, 100 µL of 10% SDS was added and mixed to release the purple colour from the cells. Optical density at 570 nm was measured using microplate reader (Arokiyaraj *et al.* 2007)

Cytotoxicity Assay. Continuous A549 cells were grown in the culture medium containing DMEM medium with 10% FBS, and 1% Pen-Strep antibiotics, and incubated at 37 °C in a humidified incubator with 5% CO₂. Confluent A549 cell was trypsinized and seeded at 5×10^4 cells/well in 96-well microplates, incubated for 24 h at 37 °C to allowed cell attached onto the plate. Then the culture medium was discarded and fresh medium culture contained different concentrations of the extract were added and incubated for 24 h. The 50% inhibitory concentration (IC₅₀) was defined by the MTT method. Briefly, the medium of the treated cells was removed, then 100 µL of medium containing 0.5 mg/mL MTT was added to each well. After incubation at 37 °C with 5% CO₂ incubator for 4 h, 100 µL of 10% SDS was added and mixed to release the purple color from the cells. Last incubation was performed again in the dark for 24 h, before optical density was measured at 570 nm using ELISA reader (Sadeghi-aliabadi *et al.* 2008; Heny *et al.* 2012).

Inhibitory Effect of the Viral Titer on Infected Cell. The A549 infected SRV cell were seeded on 12 well plates with density of 10^4 cells in 2 mL medium culture/well, then incubated at 37 °C and 5% CO₂ incubator. After 24 h, medium was discarded and added with the 3 mL fresh medium containing 50 µg/mL of *A. paniculata* extract or 50 µg/mL lamivudin as positive control. After 24 h incubation, 500 µL of medium was replaced with 500 µL fresh medium containing the same concentration of extract or lamivudin. Medium replacement was done every day until 5 days incubation and the plate was kept in -20 °C until use.

RNA Extraction and PCR Analysis. Viral RNA of SRV was extracted from the infected cells using Viral RNA Extraction Kit (Qiagen, Hilden, Germany).

Download English Version:

<https://daneshyari.com/en/article/2085818>

Download Persian Version:

<https://daneshyari.com/article/2085818>

[Daneshyari.com](https://daneshyari.com)