

# Nutrient Intake and Digestibility of Cynomolgus Monkey (*Macaca fascicularis*) Fed with High Soluble Carbohydrate Diet: A Preliminary Study

DEWI APRI ASTUTI<sup>1\*</sup>, IRMA HERAWATI SUPARTO<sup>2,3</sup>, DONDIN SAJUTHI<sup>2,3</sup>, I NENGGAH BUDIARSA<sup>3</sup>

<sup>1</sup>Department of Animal Nutrition and Feed Technology, Faculty of Animal Husbandry, Bogor Agricultural University, Darmaga Campus, Bogor 16680, Indonesia

<sup>2</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Darmaga Campus, Bogor 16680, Indonesia

<sup>3</sup>Primate Research Centre, Bogor Agricultural University, Darmaga Campus, Bogor 16680, Indonesia

Received March 18, 2009/Accepted December 22, 2009

High carbohydrate as obese diet is not yet available commercially for monkeys. Therefore, this preliminary study was to carry out nutrient intake and digestibility of cynomolgus monkeys (*Macaca fascicularis*) fed with high soluble carbohydrate diet compared to monkey chow. Five adult female macaques (average body weight 2.67 kg) were made to consume freshly diet. Commercial monkey chows (contains 3500 cal/g energy and 35% starch) were fed to three adult females (average body weight 3.62 kg). Nutrient intakes and digestibility parameters were measured using modified metabolic cages. Result showed that average of protein, fat, starch, and energy intakes in treatment diet were higher than control diet (T-test). Fat intake in the treatment diet was three times higher, while starch and energy intakes were almost two times higher than monkey chow. Digestibility percentage of all nutrients were the same in both diets except for the protein. The study concludes that the freshly prepared high sugar diet was palatable and digestible for the cynomolgus monkeys. Further studies are in progress to develop obese diet high in energy content based on fat and source of starch treatments.

Key words: *Macaca fascicularis*, obese, metabolic cages, intake, digestibility

## INTRODUCTION

Obesity is a global problem caused by metabolic syndrome and affecting an estimated of 300 million people worldwide (WHO 2002). Its prevalence is increasing in both developed and developing countries throughout the world. The prevalence of the metabolic syndrome is high among obese children and adolescents and it becomes more severe if not intervened. This disorder can lead to diabetes mellitus which usually followed by phases of obesity.

Bennet *et al.* (1995) reported that diet containing 4.2 Calories/kg of energy and 50-70% of soluble carbohydrate (sucrose and dextrin) can cause rhesus monkeys (*Macaca mulatta*) to become obese. In particular, diabetes mellitus in rhesus monkeys (*Macaca mulatta*) is preceded by phases of obesity and hyperinsulinaemia which is similar to Type 2 (non-insulin-dependent) diabetes mellitus in man (de Koning *et al.* 1993).

Riis (1983) reported that a high plasma concentration of acetate, glucose and insulin will stimulate fatty acid synthesis. Increased concentration of fat in the diet results in higher uptake of fatty acid whereas diet that causes the elevation of concentrations of glucose and insulin favor esterification and tend to increase the lipid balance. A number of obesity studies

have been carried out to investigate the effect stress using bonnet monkey (*Macaca radiata*) (Kaufman *et al.* 2007), genetic aspect using baboon (Anthony *et al.* 2003) and to evaluate energy consumption on the nighttime to body weight gain using female rhesus (*Macaca mulatta*) (Sullivan *et al.* 2005). Obesity is caused by two major causes of obesity are genetic factor (50-90%) and environmental factors (e.g. feeding and level of activity) (Chen *et al.* 2000). The aim of this study was to determine nutrient intake and digestibility of high sugar and starch diet as oppose to monkey chow in cynomolgus monkey.

## MATERIALS AND METHODS

**Preparation of Treatment Diet.** Eight adult female cynomolgus monkeys (*Macaca fascicularis*) with age range 6-8 years (estimated by dental examination) and initial body weight 2.67-3.62 kg, obtained from PT. IndoAnilab Bogor, Indonesia. All monkeys were subjected to standard Veterinary Care and Use Committee (ACUC). Animals were held in individual cages positioned to allow them to see and hear each other.

The monkeys were adjusted to the new obese diet for two weeks prior to actual feeding trial of four months. The animals were divided into two groups with unequal replication. The first group consisted of five animals (average body weight 2.67 kg) fed with the high sugar and starch diet whereas the second group consisted three animals (average body weight

\*Corresponding author. Phone: +62-251-8626213,  
Fax: +62-251-8622842, E-mail: dewiapriastuti@yahoo.com

3.62 kg) fed with control diet (monkey chow). Monkeys were fed 120 g (as fed) of diet per head per day. They were fed once a day at 08:00 hours followed by one piece of 100 g *Ambon* banana at 12:00 hours. Water was available *ad libitum*. The treatment diet contained 3570 calories/g, 8.10% fat, and 50.57% starch with ingredient consisting of 15% sugar, 12.5% dextrin, 40% wheat flour, 4% tallow, 6% vegetable oil, 7% fish meal, 5% maizena, 5.5% heated soybean meal, 1.5% minerals, 2% CMC, and 1.5% agar-agar as fiber source. The proximate analysis of treatment diet and monkey chow are as shown in Table 1.

The nutrient intakes and digestibility parameters were measured using modified metabolic cages during the last five days of the study. Nutrient intake was calculated by multiplying dry matter feed intake with nutrient concentration of the diet. Sample of feces (10% of daily total production) were collected from each group and directly frozen until requirement for proximate determination (Table 2). Apparent nutrient digestibility (%) was calculated by the difference in nutrient intake and nutrient in the feces (Mc Donald *et al.* 2002) as described in the following formula:

Table 1. Proximate analysis of high soluble carbohydrate diet and monkey chow

| Nutrients (%)        | Treatment diet | Monkey chow |
|----------------------|----------------|-------------|
| Dry matter           | 71.90          | 89.00       |
| Protein              | 10.15          | 15.00       |
| Crude fat            | 8.10           | 5.00        |
| Mineral              | 2.93           | 8.08        |
| Crude fiber          | 0.15           | 3.49        |
| Starch               | 50.57          | 34.95       |
| Gross energy (cal/g) | 3570           | 3500        |

Laboratory Feed Technology, Department Animal Nutrition, IPB (2006).

Table 2. Proximate analysis of feces

| Nutrients (%)            | Treatment diet | Monkey chow |
|--------------------------|----------------|-------------|
| Dry matter               | 31.24          | 25.80       |
| Protein                  | 24.20          | 28.60       |
| Crude fat                | 1.08           | 1.87        |
| Crude fiber              | 3.47           | 4.51        |
| Starch                   | 58.61          | 50.23       |
| Gross energy (cal/g)     | 2394           | 1269        |
| Av.Feces excretion (g/d) | 26.60          | 31          |

Laboratory Feed Technology, Department Animal Nutrition, IPB (2006).

Table 3. Nutrient consumption of *Macaca fascicularis* fed with different diet

| Treatment    | DM (g/d)       | CP (g/d)      | CL (g/d)     | Starch (g/d)  | Energy (Cal)    |
|--------------|----------------|---------------|--------------|---------------|-----------------|
| High SC diet | 103.31 ± 3.12a | 10.47 ± 1.01b | 4.73 ± 0.61a | 39.25 ± 0.67a | 242.14 ± 11.10a |
| Monkey chow  | 90.04 ± 5.01b  | 13.50 ± 1.40a | 1.78 ± 0.22b | 25.84 ± 0.12b | 129.33 ± 4.55b  |

Different letter in the same column was significantly different (P < 0.05).

Table 4. Nutrient excretion of *Macaca fascicularis* fed with different diet

| Treatment    | DM (g/d)    | CP (g/d)     | CL (g/d)      | Starch (g/d) | Energy (Cal) |
|--------------|-------------|--------------|---------------|--------------|--------------|
| High SC diet | 3.49 ± 0.12 | 3.59 ± 1.01a | 0.04 ± 0.001  | 2.04 ± 0.07  | 8.50 ± 1.10a |
| Monkey chow  | 3.99 ± 0.01 | 1.84 ± 0.40b | 0.007 ± 0.000 | 2.00 ± 0.12  | 5.07 ± 0.55b |

Different letter in the same column was significantly different (P < 0.05).

$$\text{Apparent nutrient digestibility (\%)} = \frac{(\text{Nutrient intake} - \text{nutrient in feces})}{\text{Nutrient intake}} \times 100\%$$

Analysis of dry matter (DM), crude lipid (CL), crude protein (CP), energy, and starch (NFE = nitrogen-free extractives) were carried out according to standard procedure of AOAC (1995).

Data were analyzed using T-test for comparing means of treatment diet and control (monkey chow) diet. Statistical analysis was carried out using Minitab Release 14 Program (Minitab Inc. 2003). Means of treatment diet and control (monkey chow) diet were tested for significance using T-test.

## RESULTS

The average ration eaten by the five female monkeys were 90% of the total portion which was comparable with the monkey chow group. Total dry matter of feed intake for treatment diet was about 4.0% of body mass, while for monkey chow was 3.5% of body mass (Table 3).

The average values of fat, starch and energy intakes in the treatment diet were higher (P < 0.01), while protein intake was lower (P < 0.05) than control (monkey chow) diet. Percentage of the digestibility of nutrient was the same for both diets (around 96%), however protein digestibility for treatment diet (65%) was lower (P < 0.05) than control diet (86%). It seems to be high protein excretion in the treatment diet which almost 2 times compare to control diet (Table 4).

## DISCUSSION

Dierenfeld *et al.* (2006) reported that total dry matter feed intake of proboscis monkey was 3% of body mass. The high sugar and starch in diet caused the high dry matter intake. This resulted in fat intake in treatment diet three times higher than in the monkey chow diet (4.73 vs 1.78 g/d), whereas starch and energy intakes were almost two times higher than monkey chow (39.25 vs 25.84 g/d and 242.14 vs 129.33 Cal/d). Monkey fed treatment diet consumed on average 242 Cal/head/day. Under normal condition, the energy intake of these animals has met its requirement. However to develop obesity, it appears more caloric intake (300 Cal/head/day) is required (Bennett *et al.* 1974). Feeding studies using intragastric feeding with 600-1,050 Calories of energy diet was shown to

Download English Version:

<https://daneshyari.com/en/article/2085971>

Download Persian Version:

<https://daneshyari.com/article/2085971>

[Daneshyari.com](https://daneshyari.com)