



Lab Resource: Stem Cell Line

## Derivation of the human embryonic stem cell line RCe007-A (RC-3)

P.A. De Sousa<sup>a,b,c,\*</sup>, B. Tye<sup>a</sup>, K. Bruce<sup>a</sup>, P. Dand<sup>a</sup>, G. Russell<sup>a</sup>, J. Gardner<sup>a</sup>, J.M. Downie<sup>a</sup>, M. Bateman<sup>a</sup>, A. Courtney<sup>a</sup><sup>a</sup> Roslin Cells Limited, Nine Edinburgh Bio-Quarter, 9 Little France Road, Edinburgh EH16 4UX, UK<sup>b</sup> Centre for Clinical Brain Sciences, University of Edinburgh, UK<sup>c</sup> MRC Centre for Regenerative Medicine, University of Edinburgh, UK

## ARTICLE INFO

## Article history:

Received 25 February 2016

Accepted 29 February 2016

Available online 9 March 2016

## ABSTRACT

The human embryonic stem cell line RCe007-A (RC-3) was derived from a blastocyst voluntarily donated as unsuitable and surplus to fertility requirements following ethics committee approved informed consent under licence from the UK Human Fertilisation and Embryology Authority. The cell line shows normal pluripotency marker expression and differentiation to the three germ layers *in vitro*. It has a normal 46XX female karyotype and HLA and blood group typing data is available.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Resource table

Name of stem cell construct	RCe007-A
Alternative name	RC-3, RC3
Institution	Roslin Cells Ltd.
Person who created resource	B. Tye, K. Bruce, P. Dand, G. Russell, J. Gardner
Contact person and email	Paul.desousa@roslincells.com; Paul.desousa@ed.ac.uk Janet.downie@roslincells.com Aidan.courtney@roslincells.com Malcom.bateman@roslinfoundation.com
Date archived/stock date	17 October 2007 (pre seed bank at passage 5 on feeders) 21 July 2008 (banked at passage 35)
Type of resource	Biological reagent: cell line
Sub-type	hESC, research grade
Origin	Blastocyst with ICM and trophoblast
Key transcription factors	Oct4 (confirmed by flow cytometry and immunocytochemistry), Nanog, Sox 2 (confirmed by immunocytochemistry)
Authentication	See Quality Control test summary, Table 1
Link to related literature (direct URL links and full references)	N/A
Information in public databases	<a href="http://hpscereg.eu/cell-line/RCe007-A">http://hpscereg.eu/cell-line/RCe007-A</a>
Ethics	Informed consent obtained. Scotland A Research Ethics committee approval obtained (07/MRE00/56). Conducted under the UK Human Fertilisation and Embryology Authority licence no R0136 to centre 0202.

## Resource details

RCe007-A (RC-3) was derived from a fresh blastocyst that was surplus to requirement or unsuitable for clinical use. The cell line was derived by whole embryo outgrowth on mitotically inactivated human fibroblast (HDF) feeder cells using HDF conditioned medium and expanded under feeder free conditions.

RCe007-A (RC-3) was shown to be pluripotent by expression of Oct-4, Nanog, Sox-2, Tra-1-60 and Tra-1-81, but not SSEA-1, using immunocytochemistry (Table 1, Fig. 1). By flow cytometric analysis, the expression of pluripotency markers Oct-4 and SSEA-4 was 86.2% and 96.8%, respectively, whereas low expression of the differentiation marker SSEA-1 (5.9%) was observed (Fig. 2). Differentiation to the three germ layers, endoderm, ectoderm and mesoderm, was demonstrated using embryoid body formation and expression of the germ layer markers  $\alpha$ -fetoprotein,  $\beta$ -tubulin and muscle actin (Fig. 3).

HLA Class I and II typing is available and blood group genotyping gave the blood group O<sub>1</sub>O<sub>1</sub> (Table 2).

## Verification and authentication

The cell line was analysed for genome stability by G-banding (Fig. 4) and showed a normal 46XX female genotype. The cell line is free from mycoplasma contamination as determined by RT-qPCR.

## Materials and methods

## Ethics

Derivation of hESC from surplus to requirement and failed to fertilise/develop embryos was approved by The Scotland A Research

\* Corresponding author at: Centres for Clinical Brain Sciences & Regenerative Medicine, University of Edinburgh, Chancellor's Building, 49 Little France Crescent, Edinburgh, EH16 4SB Scotland, UK.

**Table 1**  
Summary of quality control testing and results for RCe007-A (RC-3).

Classification	Test	Purpose	Result
Donor screening	HIV 1 + 2 Hepatitis B Hepatitis C	Donor screening for adventitious agents	Negative
Phenotype	Immunocytochemistry	To assess levels of staining for the pluripotency markers Oct 4 and Nanog	Expression of Oct-4, Nanog, Sox-2, Tra-1-60 and Tra-1-81
	Flow cytometry	Assess antigen levels & cell surface markers commonly associated with hESC	Oct 3/4: 86.2% SSEA-4: 96.8% SSEA-1: 5.9%
Genotype (details provided in Table 2)	Blood group genotyping (DNA Analysis)	To establish blood group of the line	O <sub>1</sub> O <sub>1</sub>
	Karyology (G-banding)	Confirmation of normal ploidy by G-banding	46XX
	HLA tissue typing	To establish full HLA type I and II genotype of the line	HLA typed Class I and Class II
Microbiology and virology	Mycoplasma Endotoxin	Mycoplasma testing by RT-qPCR Screening for endotoxin levels	Negative 3.18 EU/mL
Morphology	Photography	To capture a visual record of the line	Normal
Differentiation potential	Embryoid body formation	To show differentiation to three germ layers	Expression of muscle actin, $\beta$ -tubulin and $\alpha$ -feto protein

Ethics Committee and local ethics board at participating fertility clinics and conducted under licence no R0136 from the UK HFEA with informed donor consent.


#### Cell culture

Fresh embryos were cultured EmbryoAssist (Origio (Medicult), Denmark) until Day 3 or BlastAssist (Origio) after Day 3 of development. Embryos were cultured at 36.5–37.5 °C, 5.0 ± 0.5% CO<sub>2</sub>, 5.0 ± 0.5% O<sub>2</sub> in drops under paraffin oil (Origio) and transferred to fresh medium at least every 2–3 days.

By Day 8 of development, or when spontaneous hatching occurred, embryos were placed in derivation conditions consisting of mitotically inactivated neonatal human dermal fibroblasts (HDFs) (ThermoFisher Scientific (Cascade Biologics), Paisley, UK) on tissue culture plastic precoated with 2 µg/cm<sup>2</sup> human laminin (Sigma Aldrich, Dorset, UK) as per manufacturer's recommendation. If required, assisted hatching was performed by removing the zona pellucidae mechanically using Swemed cutting tools (Vitrolife, Göteborg, Sweden).

HDF cells were cultured in DMEM (Lonza, Slough, UK), 10% FCS (GE Healthcare (PAA), Buckinghamshire, UK) and 2 mM L-glutamine (ThermoFisher Scientific). HDF were mitotically inactivated using gamma irradiation at 50 Gy using a Gammacell Elite 1000 machine. For use as a feeder layer, irradiated HDFs were plated at 2–50,000 cells/cm<sup>2</sup> in HDF conditioned medium (80% Knockout-DMEM, 20% Knockout serum replacement (KOSR), 1 mM glutamine, 0.1 mM  $\beta$ -mercaptoethanol, 1% nonessential amino acids, and 4 ng/ml human bFGF (all ThermoFisher Scientific) over 24 h intervals over 7 days) supplemented with an additional 24 ng/ml human bFGF. Cells were cultured at 36.5–37.5 °C, 5.0 ± 0.5% CO<sub>2</sub>, 5.0 ± 0.5% O<sub>2</sub> and 50% medium exchanged 6 days a week.

The established cell line was expanded and banked using CellStart matrix and Stempro hESC Serum Free Medium (ThermoFisher Scientific). Passaging was performed mechanically using an EZ passage tool (ThermoFisher Scientific). hESC lines were expanded to 25–30 wells of a 6-well plate and cryopreserved in 0.5–1 ml KOSR based cryopreservation solution (75% KO-DMEM, 15% Xeno-free KOSR (ThermoFisher Scientific) and 10% DMSO (Origen Biomedical, Texas, USA)) or Cryostor CS10 (Biolife Solution, Washington, USA).



**Fig. 1.** RCe007-A (RC-3) expresses pluripotency markers Oct-4, Nanog, Sox-2, Tra-1-60 and Tra-1-81, but not SSEA-1. Cell nuclei are counterstained with DAPI (blue).

Download English Version:

<https://daneshyari.com/en/article/2094035>

Download Persian Version:

<https://daneshyari.com/article/2094035>

[Daneshyari.com](https://daneshyari.com)