



AHL communication is a widespread phenomenon in biomining bacteria and seems to be involved in mineral-adhesion efficiency

Lina María Ruiz^a, Susana Valenzuela^a, Matias Castro^a, Alex Gonzalez^a, Marine Frezza^b, Laurent Soullère^b, Thore Rohwerder^c, Yves Queneau^b, Alain Doutheau^b, Wolfgang Sand^c, Carlos A. Jerez^a, Nicolas Guilian^{a,*}

^a Bacterial Communication Unit, Laboratory of Molecular Microbiology and Biotechnology, Department of Biology, Faculty of Sciences and Institute for Cell Dynamics and Biotechnology, University of Chile, Santiago, Chile

^b Laboratoire de Chimie Organique, UMR 5246-ICBMS CNRS, Université Lyon 1, INSA, Lyon, France

^c Biofilm Centre Aquatische Biotechnologie, Universität Duisburg – Essen, Duisburg, Germany

ARTICLE INFO

Available online 24 June 2008

Keywords:

Quorum sensing

Acyl-homoserine lactones

Biofilm

Biomining

A. ferrooxidans

ABSTRACT

Biofilm development plays a pivotal role in the bioleaching process. Quorum sensing (QS) is recognized as one of the main regulatory ways of extracellular polymeric substances (EPS) production and biofilm formation in bacteria. Previous works revealed that *Acidithiobacillus ferrooxidans* ATCC 23270 strain possesses a functional QS type AI-1 system. However, there is neither available data concerning the presence of a QS system in other *A. ferrooxidans* strains nor any other biomining bacterial species. Thus, in this work, different strains of *A. ferrooxidans* (ATCC 19859, ATCC 1927, ATCC 33020, DSMZ 583, DSMZ 9464), *Acidithiobacillus thiooxidans* (DSMZ 504, DSMZ 9463, DSMZ 11478) and *Leptospirillum ferrooxidans* (DSMZ 2391, DSMZ 2705) were screened for their ability to produce QS-signaling molecules [acyl-homoserine lactone (AHLs)]. Thin-layer chromatography (TLC) analysis revealed that all the *A. ferrooxidans* and *A. thiooxidans* strains produced AHL-signaling molecules while both tested *L. ferrooxidans* strains did not. Nevertheless, by using bioinformatic tools to screen the genome sequence of a *Leptospirillum* sp. type III belonging to an AMD biofilm, a QS type AI-1 locus in which the structural organization is different from that of *A. ferrooxidans* has been identified. The effect of synthetic AHLs and AHL-analogues on attachment of *A. ferrooxidans* ATCC 23270 to pyrite was analyzed. Interestingly, preliminary results suggested that some of these molecules change the rate and extent of bacterial attachment to pyrite. The production of AHLs by *A. ferrooxidans* and *A. thiooxidans* strains, the identification of a QS type AI-1 locus in *Leptospirillum* sp. type III and the impact of AHLs and AHL-analogues on the pyrite adhesion of *A. ferrooxidans* cells open new perspectives for the (bio)mining industry to eventually improve bioleaching process and control the acid mine drainage (AMD) contamination.

© 2008 Elsevier B.V. All rights reserved.

1. Introduction

The attachment and adhesion of acidophilic bacteria to the sulfidic energy source and the subsequent biofilm formation are prerequisites for mineral dissolution in natural environments and industrial operations. Therefore, a thorough understanding of the molecular mechanisms involved in the control and regulation of biofilm formation would be of great importance to control the bioleaching process from the biomining industry and the acid mine drainage (AMD) damages.

Biofilm formation is a complex process that can be regulated at different levels through diverse mechanisms. The most studied mechanism that controls biofilm development is quorum sensing (QS) signaling (Waters and Bassler, 2005). QS is a cell-to-cell signaling

system that functions by means of diffusible autoinducer (AI) signal molecules that allow the regulation of cellular processes in a manner that depends on the density of the microbial population.

Recently, a functional QS system type AI-1 involving two divergent genes *afel* and *afeR* has been identified in the acidophilic bacterium *Acidithiobacillus ferrooxidans* (Farah et al., 2005; Rivas et al., 2005). Overexpression of the protein Afel in *Escherichia coli* and the characterization of associated synthesis of AHLs demonstrated that Afel is an AHL synthase (Farah et al., 2005). *A. ferrooxidans* produces AHLs with acyl chains whose length oscillates between 8 and 16 carbons and presents substitutions of the type oxo- and hydroxyl- in C-3 (3-hydroxy-C8-AHL, 3-hydroxy-C10-AHL, C12-AHL, 3-oxo-C12-AHL, 3-hydroxy-C12-AHL, C14-AHL, 3-oxo-C14-AHL, 3-hydroxy-C14-AHL, and 3-hydroxy-C16-AHL). The AHL hydroxy substitution was independent of the nature of the culture medium whereas the keto substitution was displayed only in cells grown on sulfur and thiosulfate. Besides, the transcription of *afel* gene is higher in sulfur- and thiosulfate-grown cells with respect to iron-grown cells and appears to be related to phosphate metabolism (Farah et al., 2005). Based on this experimental background,

* Corresponding author. Unidad de Comunicación Bacteriana, Laboratorio de Microbiología Molecular y Biotecnología, Departamento de Biología, Instituto de Dinámica Celular y Biotecnología, Facultad de Ciencias, Universidad de Chile, Santiago, Chile. Tel.: +56 978 7241.

E-mail address: nguilian@uchile.cl (N. Guilian).

different authors suggested that the QS system type AI-1 could regulate biofilm formation in *A. ferrooxidans* (Farah et al., 2005; Rivas et al., 2005; Valenzuela et al., 2006; Valenzuela et al., 2007).

This work attempts to examine if the quorum sensing type AI-1 system identified in *A. ferrooxidans* ATCC 23270 is a widespread phenomenon in *A. ferrooxidans* species and other biomining bacteria. Knowledge of the molecular mechanisms that regulate biofilm formation in biomining bacteria such as *A. ferrooxidans* is limited. Therefore, the involvement of the AHL-signaling molecules in the attachment of the type strain *A. ferrooxidans* ATCC 23270 to mineral surfaces was examined. Since marker exchange mutagenesis for the construction of QS mutants in *A. ferrooxidans* is still very difficult to carry out, we decided to use different synthetic AHLs and AHL-analogues that have been evaluated as potential inhibitors and/or activators of QS in other microorganisms (Castang et al., 2004; Geske et al., 2005; Frezza et al., 2006).

2. Materials and methods

2.1. Bacteria

The following strains of biomining bacteria obtained from culture collections were used: (i) six strains of *A. ferrooxidans* ATCC 23270 type strain, ATCC 19859, ATCC 1927, ATCC 33020, DSMZ 583, DSMZ 9464; (ii) three strains of *Acidithiobacillus thiooxidans* DSMZ 504, DSMZ 9463, DSMZ 11478; two strains of *Leptospirillum ferrooxidans* DSMZ 2391, DSMZ 2705.

2.2. Growth media and culture conditions

A. ferrooxidans strains were grown in ferrous iron-containing modified 9K medium at pH 1.5 as described previously, and growth on elemental sulfur was done at pH 2.5 with 5% (wt/v) sulfur prills (Amaro et al., 1991). Growth of *A. ferrooxidans* strains in thiosulfate was done at pH 4.6 in DSMZ medium 71 containing 20 mM thiosulfate as described before (Ramírez et al., 2004). *A. thiooxidans* and *L. ferrooxidans* strains were grown on their corresponding DSMZ medium.

2.3. Southern analysis

Genomic DNAs were obtained by using the Wizard Genomic DNA Purification Kit for Gram-negative bacteria, as described by Promega. 5 µg of purified genomic DNA was digested by EcoRI and HindIII restriction enzymes (New England Biolabs) at 37 °C during 4 h. After separation of the restriction enzyme-digested DNA fragments by electrophoresis, they were denatured and transferred to a positively charged nylon membrane (Immobilon-NY+; Millipore) by the semi-dry capillary method (Sambrook and Russel, 1989). ³²P labelled-*afel* probe was obtained by using the NEBlot kit (New England Biolabs) according to the manufacturer's recommendations. Prehybridizations and hybridizations were accomplished as described by Sambrook and Russel (1989). ³²P-signal detection was carried out by using the PhosphorImager (Molecular Imager FX; Biorad).

2.4. AHL-signaling molecules determination

Extracts for analytical thin-layer chromatography (TLC) and LC-MS-MS were prepared from 600-ml cultures. Bacteria were removed by centrifugation and the supernatant was extracted twice with equal volumes of dichloromethane (DCM) as described previously (Farah et al., 2005). Extracts were then dried over anhydrous magnesium sulfate, filtered, and evaporated to dryness.

For TLC plate analysis, dried samples were dissolved in 30–60 µl of HPLC-grade ethyl acetate. Samples (20–40 µl) were spotted onto C₁₈ reversed-phase TLC plates (Merck) along with a set of standards and developed with methanol/water (60:40; vol/vol). After development,

the TLC plates were air-dried and AHL-signaling molecules were detected by using as a bioreporter *Agrobacterium tumefaciens* NTL4 (pZLR4) as described previously (Shaw et al., 1997). Briefly, 250 µl of an overnight culture of *A. tumefaciens* NTL4(pZLR4) AHL reporter strain grown in 25-ml LB medium with gentamicin 25 µg/ml was inoculated in AB minimal glucose medium (ABm) and grown during 8 h at 30 °C with shaking. The culture was then mixed with an equal volume of 1.5% TOP agar containing 80 µg/ml 5-bromo-4chloro-3-indolyl-β-D-galactopyranoside (X-gal), and the preparation was spread over the surface of developed TLC plates. Overlaid TLC plates were incubated overnight at 30 °C in a closed plastic container. Production of AHL is visualized by the development of a blue color.

For LC-MS-MS analysis, residues were dissolved in 0.5 ml of HPLC-grade acetonitrile and analyzed by using reverse-phase liquid chromatography coupled with positive-ion electrospray ionization and ion trap mass spectrometry (LC-MS-MS) (Morin et al., 2003). In the case of 3-hydroxy-AHLs, their identification was done by comparison with synthetic 3-hydroxy-AHLs based on three criteria: MS-MS fragmentation product ions ([M+H-H₂O]⁺ and *m/z* 102), their relative intensities and HPLC retention times.

2.5. Attachment assay

This assay was carried out only with the type strain of *A. ferrooxidans*, ATCC 23270. It is based on the reduction of the number of planktonic cells in the solution due to the attachment of the bacteria to a mineral sulfide substratum (Gehrke et al., 1998; Harneit et al., 2006). Pyrite (FeS₂) (Suior mine in Baia Mare, Romania) was the substratum used for the attachment assay. Ground FeS₂ was wet sieved to a grain size of 50–100 µm (Test sieves, Retsch, Germany). To remove ferric iron, the FeS₂ was boiled in 6 N HCl with stirring for 1 h. Then, the HCl was removed by washing extensively with distilled H₂O. Next, sulfur compounds were extracted from the FeS₂ by washing twice with acetone. Finally, FeS₂ aliquots were put under vacuum and gassed with nitrogen and sterilized at 115 °C for 48 h (Schippers and Sand, 1999). For the attachment assay, 5 × 10⁹ cells were incubated with 20% (w/v) FeS₂ in a final volume of 50-ml mineral salt solution in 100-ml Erlenmeyer flask. *A. ferrooxidans* ATCC 23270 grown in FeSO₄ medium was harvested in the late exponential growth phase. Harvesting was performed by cross flow filtration (Hemoflow F60S, Fresenius, Germany) or centrifugation for 20 min at 6000 ×g, at 10 °C and the cells thus obtained were resuspended in 50 ml of mineral salts solution. Then, cell concentration was determined using a Thoma counting chamber (depth 0.02 mm, smallest square area 0.0025 mm²). Subsequently, cell concentration was adjusted at 1 × 10⁸ cells per ml. Next, this cell suspension was distributed among 100-ml Erlenmeyer flasks. Bacteria harvested by centrifugation were supplemented with 0.2 g of ferric iron l⁻¹ and kept overnight at 17 °C to regenerate their EPS. Afterwards, AHLs or AHL-analogues were added and the mix was incubated for 1 h at 30 °C, at 180 rpm. Subsequently, 10 g of FeS₂ was added to each Erlenmeyer flask containing the assay mixture and agitated on a benchtop shaker (Type TR, Infors HT, Switzerland) at 120 rpm and room temperature for 8 h. Samples (1 ml) of the culture supernatant were taken at specific time intervals and the cell concentration was determined using a Thoma counting chamber. Cell counting was done with a light microscope (Zeiss) in phase contrast mode with 400-fold magnification.

3. Results

3.1. Quorum sensing type AI-1 system and biomining bacteria

In previous experiments, we reported that *A. ferrooxidans* type strain possesses a functional QS type AI-1 system and is able to produce nine different kinds of AHL-signaling molecules (Farah et al., 2005). To examine if the QS type AI-1 system is a widespread phenomenon in biomining bacteria, we analyzed if other strains of *A.*

Download English Version:

<https://daneshyari.com/en/article/213228>

Download Persian Version:

<https://daneshyari.com/article/213228>

[Daneshyari.com](https://daneshyari.com)