

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: <http://www.elsevier.com/locate/pjnns>

## Case report

# Painful tonic spasms and brainstem involvement in a patient with neuromyelitis optica spectrum disorder

Corina Roman-Filip<sup>a</sup>, Aurelian Ungureanu<sup>b,\*</sup>, Mihaela Cernușcă-Mițaru<sup>c</sup><sup>a</sup>Neurology Department, Academic Emergency Hospital, “Lucian Blaga” University Sibiu, Sibiu, Romania<sup>b</sup>Neurology Department, Academic Emergency Hospital, Sibiu, Romania<sup>c</sup>Academic Emergency Hospital, “Lucian Blaga” University Sibiu, Sibiu, Romania

## ARTICLE INFO

## Article history:

Received 22 May 2015

Accepted 30 October 2015

Available online 24 November 2015

## Keywords:

Painful tonic spasms

Neuromyelitis optica

Brainstem

## ABSTRACT

Neuromyelitis optica (NMO) is an inflammatory-demyelinating disease of the central nervous system classically characterized by optic neuritis and severe myelitis. New diagnostic criteria defined neuromyelitis optica spectrum disorder as limited forms of NMO or diverse neurologic presentations in the presence of specific aquaporin-4 antibodies. We report the case of a 57-year-old woman admitted in our department for recurrent attacks of optic neuritis, tetraparesis with severe painful tonic spasms of the left limbs and brainstem involvement. Painful tonic spasms have been described as movement disorders associated with multiple sclerosis, but a growing number of reports describe them in cases of NMO.

© 2015 Polish Neurological Society. Published by Elsevier Sp. z o.o. All rights reserved.

## 1. Introduction

Neuromyelitis optica (NMO) spectrum disorder (NMOSD) is a severe inflammatory and demyelinating disease of the central nervous system. The pathogenic process is attributed to destruction of aquaporin-4 water channels by specific auto-antibodies (NMO-IgG) [1]. NMOSD largely includes limited forms of NMO, such as optic neuritis (ON) and long extensive transverse myelitis, characteristic brain involvement as hypothalamus, corpus callosum, and brainstem lesions or Asian type of optic-spinal multiple sclerosis (MS) [2]. Painful

tonic spasms (PTS) or paroxysmal painful dystonia were first described as movement disorders associated with lesions of the cervical spinal cord, especially in MS [3]. The characteristics of painful tonic spasm are described as paroxysmal episodes lasting seconds or minutes, accompanied by intense pain and tonic postures of the limbs. Ephaptic transmission between abnormal demyelinated tracts could explain the spasms [4]. Recently, the presence of these manifestations seems to be associated to NMO rather than MS or idiopathic acute transverse myelitis [5]. Alongside with myelitis or isolated, brainstem involvement is found in almost 32% of the patients diagnosed with NMOSD [6].

\* Corresponding author at: Neurology Department, Academic Emergency Hospital, 2-4 Pompeiu Onofreiu St., 550166 Sibiu, Romania. Tel.: +40 745088528/269215050.

E-mail address: [aurelianungureanu@hotmail.com](mailto:aurelianungureanu@hotmail.com) (A. Ungureanu).


<http://dx.doi.org/10.1016/j.pjnns.2015.10.010>

0028-3843/© 2015 Polish Neurological Society. Published by Elsevier Sp. z o.o. All rights reserved.

## 2. Case report

A 57-year-old woman admitted in our department for acute onset of painful spasms of the left limbs associated with gait and visual disturbances. The medical history is positive for a left ON diagnosed in 2001 that was partially responsive to intravenous (IV) methylprednisolone with normal MRI at that time. The patient was diagnosed at that moment with isolated ON syndrome, but no other diagnostic workup was followed. In December 2014, she presented bilateral decrease of visual acuity in both eyes and progressive numbness on the left arm followed by minor weakness with negative MRI findings for demyelinating disease, although two small unspecific lesions were described in the frontal white matter. Spastic gait and mild spasticity in the lower limbs were noted. A third episode of left ON occurred in April 2015, without any response to IV methylprednisolone and with subsequent amblyopia. After 2 weeks, the patient presented to the emergency room with sudden painful paroxysmal tonic flexion and adduction of the left upper limb, sparing the face, occurring every 5–10 min with duration of maximum 10–20 s. Inconsistently, they involved the left lower limb, were elicited by passive and voluntary movements of the left limbs, but the majority were observed as appearing spontaneously. There was no other history of cardiovascular pathology (arterial hypertension, cardiac arrhythmia, cardiac ischaemic disease, and dyslipidaemia),

autoimmune diseases or psychiatric diseases. Expanded neurologic examination showed spastic tetraparesis with left side predominance, bilateral plantar clonus, brisk reflexes, left eye blindness with absent pupillary reflex, decreased right eye visual acuity and bilateral nystagmus. The patient accused paraesthesia on the left hemibody without any clear sensory loss and accused imperious micturition. The patient needed bilateral assistance while walking. The general examination found a blood pressure of 110/80 mmHg, rhythmic heart beats with 76 bpm, but was otherwise normal. The acute presentation of tonic spasms raised the suspicion of epileptic seizures. The electroencephalogram was normal during and between the attacks, together with other neurologic findings making the epileptic tonic seizures less plausible. Brain MRI revealed several small frontal ischaemic lesions, T2 and FLAIR hyperintensities at the level of the medulla oblongata with bilateral corticospinal tracts involvement and contrast enhancement at the level of brainstem–spinal junction (Fig. 1). The lesion was not considered characteristic for anterior spinal artery infarction. An additional finding was a C5–C6 intervertebral disc protrusion with compression of the spinal cord and subjacent post-compressive ischaemic lesion. Laboratory studies showed positive serum and cerebrospinal fluid (CSF) NMO-IgG, elevated level of serum anticardiolipin antibodies, and slight CSF albumin elevation with borderline pleocytosis represented by lymphocytes. IgG quotient raised the suspicion of intrathecal IgG synthesis and oligoclonal



**Fig. 1** – (a) T2 axial sequence showing hyperintense lesion of the corticospinal tracts and paramedian structures of the medulla oblongata. (b) FLAIR sequence with lesions at the same level. (c) Sagittal FLAIR with hyperintensities at the level of medulla oblongata and brainstem–spinal junction. (d) Sagittal T2 showing extensive lesion of medulla oblongata. (e) Discrete gadolinium enhancement at the level of brainstem–spinal junction. (f) C5–C6 disc protrusion with subjacent myelopathy.

Download English Version:

<https://daneshyari.com/en/article/2152654>

Download Persian Version:

<https://daneshyari.com/article/2152654>

[Daneshyari.com](https://daneshyari.com)