

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: <http://www.elsevier.com/locate/pjnns>

## Case report

# Melatonin and cortisol profiles in patients with pituitary tumors

Daniel Zielonka<sup>a,\*</sup>, Jerzy Sowiński<sup>b</sup>, Stanisław Nowak<sup>c</sup>,  
Anna Ciesielska<sup>a</sup>, Jakub Moskal<sup>c</sup>, Jerzy T. Marcinkowski<sup>a</sup>

<sup>a</sup>Department of Social Medicine, Poznań University of Medical Sciences, Poznań, Poland

<sup>b</sup>Department of Endocrinology, Poznań University of Medical Sciences, Poznań, Poland

<sup>c</sup>Department of Neurosurgery, Poznań University of Medical Sciences, Poznań, Poland

## ARTICLE INFO

## Article history:

Received 9 January 2014

Received in revised form

9 October 2014

Accepted 8 December 2014

Available online 17 December 2014

## Keywords:

Melatonin

Cortisol

Pituitary tumors

Biological rhythm

## ABSTRACT

The optic tract section at the optic chiasm is expected to disturb the suprachiasmatic nucleus (SCN) rhythm, circadian rhythm and melatonin secretion rhythms in humans, although detailed studies have never been conducted. The aim of this paper was to describe melatonin and cortisol profiles in patients with a pituitary tumor exerting optic chiasm compression. Six patients with pituitary tumors of different size, four of whom had significant optic chiasm compression, were examined. In each brain, MRI, an ophthalmological examination including the vision field and laboratory tests were performed. Melatonin and cortisol concentrations were measured at 22:00 h, 02:00 h, 06:00 h, and 10:00 h in patients lying in a dark, isolated room.

One of the four cases with significant optic chiasm compression presented a flattened melatonin rhythm. The melatonin rhythm was also disturbed in one patient without optic chiasm compression. Larger tumors may play a role in the destruction of neurons connecting the retina with the suprachiasmatic nucleus (SCN) and breaking of basic way for inhibiting effect to the SCN from the retina.

© 2014 Polish Neurological Society. Published by Elsevier Urban & Partner Sp. z o.o. All rights reserved.

## 1. Introduction

The day–night rhythm of Earth's rotation regulates light-dependent rhythms in living organisms [1]. The melatonin circadian rhythm is the most stable light-dependent rhythm in humans. Even after major surgery, it returns to its pattern in

24 h [2,3]. It has been estimated that melatonin secretion increases at night with a peak at 02:00 h and is low during the day [4]. It has also been proven that photic stimulation received through the retino-hypothalamic tract (RHT) stimulates the suprachiasmatic nucleus of the hypothalamus (SCN). The suprachiasmatic nucleus (SCN) plays a major role in generating the circadian rhythm in humans [5,6]. Recently,

\* Corresponding author at: Department of Social Medicine, Poznań University of Medical Sciences, Rokietnicka Str., No. 5 “C”, 60-806 Poznań, Poland. Tel.: +48 504 609 951; fax: +48 61 8547390.

E-mail address: [daniel.zielonka@gmail.com](mailto:daniel.zielonka@gmail.com) (D. Zielonka).

<http://dx.doi.org/10.1016/j.pjnns.2014.12.004>

0028-3843/© 2014 Polish Neurological Society. Published by Elsevier Urban & Partner Sp. z o.o. All rights reserved.

other important connections were described between the SCN and forebrain and midbrain structures, *inter alia* an intergeniculate leaflet regulating the SCN's activity [7,8]. Retino-hypothalamic fibers synapse in the SCN and neuronal connections exist between the SCN and the intermediolateral gray column in the spinal cord. Preganglionic neurons pass from the spinal cord to the superior cervical ganglion, and postganglionic neurons project from this ganglion to the pineal in the nervi conarii [9,10]. The second most stable human rhythm is the cortisol circadian rhythm [11,12]. It is also related to SCN function, however, it is conducted *via* other connections from the SCN.

The aim of this study was to check how an obstruction of the connection between the retina and the pineal gland can disturb the melatonin circadian rhythm in humans. As an obstruction, pituitary gland tumors that were different in size and origin were analyzed. In the four cases considered in this study, optic chiasm compression which was significant but different in intensity was observed. In two cases no significant optic chiasm compression was stated. The cortisol circadian rhythm was used as a reference since it was expected that it would not be disturbed by pituitary tumors.

It was expected that in cases with significant optic chiasm compression, deviations would appear in the melatonin profile, namely, there would be a reduction in melatonin secretion at night (circadian rhythm 'flattening' at night) if the tumor were prompting nerve fibers innervating the pineal gland, or there would be hypersecretion of melatonin and/or the secretion peak shift if the retino-pineal connection were simply cut off by the tumor.

## 2. Materials and methods

Six patients were analyzed with pituitary tumors of different size and origin, two of these patients did not have optic chiasm compression detected in the Magnetic Resonance Imaging (MRI) examination and four of them had significant compression of the optic chiasm detected in the MRI (Table 1).

In each case a detailed interview regarding any other hormonal disturbances and concomitant disorders was

collected. An MRI of the brain, the vision field, color vision field, a general ophthalmological examination and laboratory tests were also performed.

In all patients, medications were discontinued two days prior to hospital admission; all patients were awaiting the same surgical treatment. No history was reported of prolonged sleep-wake cycle disturbances caused by other reasons before admission, of prolonged use of drugs that could influence melatonin secretion, or of ocular comorbidities. After admission the patients were kept in a dark, isolated rooms for 12 h, from 22:00 to 10:00 and in a horizontal position for 13 h from 21:00 to 10:00 the next day. Melatonin and cortisol concentrations were measured at 22:00 h, 02:00 h, 06:00 h and 10:00 h. The melatonin ELISA kit was used to detect the melatonin level in the blood serum. The minimum detectable concentration of melatonin was 3 pg/ml. Intra-assay precision varied from 3.5% to 10.5%. Inter-assay precision varied from 9% to 19%. The plasma cortisol concentration was estimated by electrochemiluminescence immunoassay (ECLIA). The minimum detectable concentration of cortisol was 0.5 nmol/l. Intra-assay precision varied from 1.5% to 1.7%. Inter-assay precision varied from 1.8% to 2.8%. The acrophase, mesor, trough and bathyphase were determined for the rhythm analysis peak.

Because there was no control group, the patients' secretion curves were compared with data from the literature [13,14]. Secretion patterns in healthy individuals as described in the literature were compared with respect to age with the results of the patients in this study. Hormone peak concentrations in different points in time that were different from those in the references were interpreted as a disturbance in secretion.

All patients gave informed written consent according to the International Conference on Harmonization – Good Clinical Practice (ICH-GCP) guidelines (<http://www.ich.org/LOB/media/MEDIA482.pdf>). Ethical approval was obtained from the Bioethics Committee of the Poznan University of Medical Sciences.

## 3. Results

An abnormal melatonin rhythm was observed in two of six cases. The first case was a patient with pineal microadenoma

**Table 1 – Patients characteristics.**

Patient no and gender (man/woman)	1 woman	2 man	3 man	4 man	5 woman	6 woman
Age of patient	26	25	49	58	64	47
Length of tumor	0.9 cm	1.7 cm	2.5 cm	3.4 cm	3 cm	0.6 cm
Width of tumor	0.9 cm	1.6 cm	2.1 cm	3.4 cm	4 cm	0.4 cm
Altitude of tumor	0.9 cm	1.3 cm	3.4 cm	4.5 cm	5 cm	Not given
Volume of tumor	729 mm <sup>3</sup>	3536 mm <sup>3</sup>	17,850 mm <sup>3</sup>	52,020 mm <sup>3</sup>	60,000 mm <sup>3</sup>	n/a
Pressure on the optic chiasm	No	Yes (mild)	Yes	Yes	Yes	No
Histopathological diagnosis	Micro adenoma	Rathke cleft cyst macroadenoma	Eosinophilic adenoma	Eosinophilic adenoma	Eosinophilic adenoma	Adenoma chromophobe microadenoma
Loss of field of vision	No	No	Yes (half of vision field)	Yes (half right eye – green color, full left eye)	Yes (concentric, red color)	No
Cortisol secretion	Correct	Correct	Correct	Disturbed	Disturbed	Disturbed
Melatonin secretion	Correct	Correct	Correct	Correct	Disturbed	Disturbed
Secretion	Prolactin	No	Prolactin	GH, prolactin	No	ACTH

Download English Version:

<https://daneshyari.com/en/article/2152672>

Download Persian Version:

<https://daneshyari.com/article/2152672>

[Daneshyari.com](https://daneshyari.com)