



# A method for reducing the thickness of the outer egg membrane of the Japanese mitten crab *Eriocheir japonica* to improve the normal zoeal larvae hatching rate of *in vitro* artificial fertilized eggs

Tai Hung Lee \*

Laboratory of Aquaculture Genetics and Genomics, Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato-cho, Hakodate, Hokkaido, 041-8611, Japan

## ARTICLE INFO

### Article history:

Received 2 July 2010

Received in revised form 21 April 2011

Accepted 5 May 2011

Available online 13 May 2011

### Keywords:

Egg membrane

Artificial fertilization

Hatching rate

Crab

Decapod

*Eriocheir japonica*

## ABSTRACT

This study was an attempt to prove that the thickness of the outer egg membrane is the key factor that contributes to the normal zoeal larvae hatching rate of the *in vitro* artificially fertilized eggs of the Japanese mitten crab *Eriocheir japonica* by artificially stretching and thinning the outer egg membrane with the help of water surface tension. The results showed that the artificially thinned outer egg membrane became six times thinner. In addition, the normal zoeal larvae hatching rate of the *in vitro* artificially fertilized eggs rose from 10% to over 66% after the thinning treatment. Discussion focuses on the mechanism of this thinning phenomenon, the role the thinned outer egg membrane plays in the increase of the normal zoeal larvae hatching rate of the *in vitro* artificially fertilized eggs and the utilization of this particular outer egg membrane thinning method.

© 2011 Elsevier B.V. All rights reserved.

## 1. Introduction

In aquaculture sciences, *in vitro* artificial fertilization is essential for performing various techniques such as chromosome engineering, gene manipulations (Purdom, 1993). As far as crabs (brachurans) are concerned, the technique has not been well established. The first success in *in vitro* artificial fertilization in crabs was reported in 1989 by Lee and Yamazaki in the Chinese mitten crab *Eriocheir sinensis*. The normal zoeal larvae hatching rate of the *in vitro* artificially fertilized eggs, however, was less than 20%.

In the author's recent study (Lee, 2009b), it was found that the normal zoeal larvae hatching rate of the *in vitro* artificially fertilized eggs in the Japanese mitten crab *Eriocheir japonica* rose from 10% up to over 90% after the outer membrane of the *in vitro* artificially fertilized eggs was removed. The finding strongly suggested that there is a close relationship between the normal zoeal larvae hatching rate and the outer egg membrane in *in vitro* artificially fertilized eggs. In addition, transmission electron microscopy observations indicated that the outer egg membrane of the *in vitro* artificially fertilized egg was 1.5 to 3 times thicker than that of the naturally spawned egg, suggesting that the abnormal thickness of the outer egg membrane of the *in vitro* artificially fertilized egg might have a negative effect on the normal zoeal larvae hatching rate. I hypothesized that this unusually thick

outer egg membrane may contribute to a decrease in not only the exchanging rate of oxygen but also the clearance rate of the metabolic waste. In addition, the thick membrane may physically suppress the enlargement of the developing embryo and impede the ecdysis of the embryos. All of these result in abnormal development of the zoeae (Lee, 2009b). Furthermore, I pointed out that the removal of the outer egg membrane was not only time-consuming but also extremely exhausting.

The purpose of this study was two-fold: 1) to prove the hypothesis proposed by Lee (2009b) that the thickness of the outer egg membrane is a key factor that contributes to the normal zoeal larvae hatching rate of *in vitro* artificially fertilized eggs, and 2) to introduce a simple and easy method for improving the normal zoeal larvae hatching rate of *in vitro* artificial fertilized eggs by reducing the thickness of the outer egg membrane.

## 2. Materials and methods

The experimental species used for this study was the Japanese mitten crab *E. japonica*, a freshwater crab of commercial value. This crab inhabits the rivers of Japan and is closely related to the Chinese mitten crab *E. sinensis* in terms of taxonomy and morphology (Sakai, 1976; Peng, 1986; Dai, 1988; Li et al., 1993; Gao and Watanabe, 1998; Li and Li, 1999; Li and Zou, 1999; Xie et al., 1999; Zhao and Li, 1999; Zhao et al., 2002; Lee et al., 2004). Copulated adult females of the Japanese mitten crab *E. japonica* were collected from the estuary of the Shiodomari River in Hakodate, Hokkaido. They were maintained

\* Tel./fax: +81 138 40 5590.

E-mail address: [thlee@fish.hokudai.ac.jp](mailto:thlee@fish.hokudai.ac.jp).

in 80% seawater (salinity: 27.6 ppt) in a well-aerated, close circulation system at 20 °C at the Faculty of Fisheries Sciences, Hokkaido University.


### 2.1. To reduce the thickness of the outer egg membrane of the *in vitro* artificially fertilized egg by water surface tension

Female crabs were monitored 24 h using a self-made oviposition alarm system (Lee, 2009a). When the alarm system signaled that a female crab was beginning to oviposit, unfertilized ripe eggs were obtained directly from its ovary as soon as possible. *In vitro* artificial fertilization was then carried out using the methods described in Lee and Yamazaki (1989; 1990). Following *in vitro* artificial fertilization, the eggs were rinsed three times with filtered (Millipore filter: 0.20 µm) sea water (salinity: 27.6 ppt).

For the thinning of the outer egg membrane, part of the rinsed fertilized eggs were immediately placed at the bottom of the sterile plastic Petri dishes filled with filtered (Millipore filter: 0.20 µm) seawater (salinity: 27.6‰) (Fig. 1A). The seawater was then removed by pipette until part of the outer egg membrane was exposed to air (Fig. 1B). And then filtered (Millipore filter: 0.20 µm) seawater (salinity: 27.6‰) was added once again by pipette. With this treatment, the eggs floated on the water surface and part of the outer egg membrane was exposed to air and stretched by the water surface tension of the seawater (salinity: 27.6 ppt). As a result, part of the outer egg membrane became thin (Fig. 1C and D).

### 2.2. Egg incubation trial for the examination of the effect of the stretched outer egg membrane

After the *in vitro* artificially fertilized eggs had floated overnight, they were brought down to the bottom of the tissue culture flask by pipette. For the experimental groups, three replicates of 50 floating treated eggs were counted and artificially incubated in 50 ml sterile tissue culture flask filled with 30 ml filtered (Millipore filter: 0.20 µm) sea water (salinity: 27.6‰). The tissue culture flasks were placed in an orbital shaker (shaking speed: 30 rpm; orbit: horizontally reciprocating in a motion that resembles the number 8; amplitude: 2 cm) in order to keep the culture solution flowing continuously. For the control groups, the untreated *in vitro* artificially fertilized eggs were counted and artificially incubated in the same way as the experimental groups.



**Fig. 1.** Schematic illustration of the treatment for thinning the outer egg membrane. A: Rinsed fertilized eggs were brought down to the bottom of the dish. B: Seawater (salinity: 27.6‰) was removed until part of the outer egg membrane was exposed to air. C: Seawater (salinity: 27.6‰) was added until the eggs floated on the water surface. D: The outer egg membrane exposed to air was stretched and became thin as a result of the water surface tension of the seawater (salinity: 27.6‰). b = bottom of the dish; j = joint region of the water surface and the outer egg membrane; oma = outer egg membrane exposed to air; oms = outer egg membrane submerged in seawater; ws = water surface.

**Table 1**

The results of the incubation trial in the experimental groups and control groups.

	Control groups	Experimental groups
Normal zoea hatching rate	10.67 ± 1.16	66.67 ± 2.31***
Abnormal zoea hatching rate	54.67 ± 2.31	29.33 ± 3.06***
Rate of the eggs unable to hatch	33.33 ± 2.31	1.33 ± 2.31**
Rate of the eggs dead before hatch	1.33 ± 1.16	2.67 ± 3.06
Total	100.00	100.00

Date represent mean (%) ± S.D. Asterisks indicate significant differences in comparison with values of control groups (\*\* $P < 0.01$ ; \*\*\* $P < 0.001$ ; *t*-test).

The eggs in both the experimental and control groups were artificially incubated at 20 °C until 10 days elapsed after the beginning of hatch. The eggs and hatched zoeae were checked and counted every day under the stereoscopic microscope. In addition, the external features and movement of the hatched zoeae were observed.

*T*-test was conducted in the statistical analysis of the results of artificial incubation in the experimental and control groups, i.e., the average normal zoeal hatching rate, average abnormal zoeal hatching rate, average rate of the eggs that were unable to hatch, and average rate of the eggs that died before hatch.

### 2.3. To examine the diameter of the outer egg membrane exposed to air

After the *in vitro* artificially fertilized eggs were floated on the surface of the filtered (Millipore filter: 0.20 µm) sea water (salinity: 27.6 ppt), twenty of them were collected at 20 min, 30 min, 40 min, 1 hr, 1.5 h, and 24 h. The outer egg membranes exposed to air were photomicrographed with light microscopy for measurement of their diameters.

### 2.4. To examine the thickness of the outer egg membrane of the floating egg

In order to examine the thickness of the outer egg membranes, transmission electron microscopy observations were carried out using the standard methods described in Hayat (1986). *In vitro* artificially fertilized eggs that had floated on filtered (Millipore filter: 0.20 µm) sea water (salinity: 27.6‰) overnight were collected and fixed. Following the fixation of the eggs, several cracks (holes or gaps) were made by pricking the eggs with a surgery blade for the penetration of chemicals to be used in the preparation of the transmission electron microscopy specimens. For comparison of the thickness of the outer egg membranes, all of the transverse sections were cut at the middle of the eggs.

## 3. Results

### 3.1. Eggs incubation trial for the examination of the effect of the thinned outer egg membrane

After 19 days of incubation, hatching was observed in both the experimental and control groups. Within a week, almost all of the zoeae were hatched. Among them some zoeae were normal and some were abnormal. Abnormal zoeae refer to those that lacked normal dorsal and rostrum spines, the first and second maxillipeds, tail forks, and were unable to swim or predate. At the end of the incubation experiment, the average normal zoea hatching rates of the experimental and the control

**Table 2**

Changes of the diameters of the outer egg membranes exposed to air after floating.

Time elapsed after floating	20 min	30 min	1 h	1.5 h	24 h
Diameter(µm)	400.25 ± 22.38	456.30 ± 22.39	669.45 ± 36.92	689.70 ± 33.34	677.85 ± 56.69

Data represent mean (%) ± S.D.

Download English Version:

<https://daneshyari.com/en/article/2422917>

Download Persian Version:

<https://daneshyari.com/article/2422917>

[Daneshyari.com](https://daneshyari.com)