

## Short Communication: Antimicrobial Drug Susceptibility of *Staphylococcus aureus* from Subclinical Bovine Mastitis in Italy

P. Moroni,\*<sup>1</sup> G. Pisoni,\* M. Antonini,† R. Villa,‡ P. Boettcher,†<sup>2</sup> and S. Carli‡

\*Department of Veterinary Pathology, Hygiene and Public Health, University of Milan, via Celoria 10, 20133 Milan, Italy

†Institute of Agricultural Biology and Biotechnology (IBBA), Consiglio Nazionale delle Ricerche (CNR), Milan, Italy

‡Department of Veterinary Sciences and Technologies for Food Safety, University of Milan, via Celoria 10, 20133 Milan, Italy

### ABSTRACT

The antimicrobial susceptibility of 68 *Staphylococcus aureus* isolates collected during 2004 from milk of cows affected by subclinical mastitis was examined. The antimicrobial agents tested were the  $\beta$ -lactams, penicillin G, amoxicillin, ampicillin, cloxacillin, amoxicillin + clavulanate, cephalonium, and cefoperazone; and other drugs including lincomycin, oxytetracycline, doxycycline, and kanamycin. Minimum inhibitory concentrations recorded show that only certain  $\beta$ -lactamase-resistant penicillins (specifically cloxacillin) or penicillin combinations (amoxicillin + clavulanate) were consistently effective against *Staph. aureus*, whereas the other  $\beta$ -lactam derivatives and drugs from other pharmacological groups were either moderately effective or ineffective. Thus,  $\beta$ -lactamase-resistant penicillins are to be considered the antimicrobial agents of choice for treatment of bovine mastitis resulting from infection by *Staph. aureus*.

**Key words:** *Staphylococcus aureus*, bovine, minimum inhibitory concentration

Bovine mastitis is the most costly disease to the dairy industry worldwide, with losses estimated at 2 billion dollars per year in the United States alone. These relevant economic losses are attributable to rejected milk, reduced milk quality, early culling, drug costs, veterinary expenses, and increased labor costs (Hoblet et al., 1991; Gruet et al., 2001). The disease is the most frequent reason for the use of antimicrobial agents on dairy farms (Erskine, 2000). Intramammary infection sustained by *Staphylococcus aureus* may result in clinical or subclinical mastitis and is usually associated with increased SCC. Appropriate treatment of mastitis during the lactation or dry period is an important compo-

nent of any mastitis control program, but the outcome for treatment of mastitis caused by *Staph. aureus* is variable and the probability of curing the disease is not high, primarily because of poor distribution of the drug in the inflamed udder and the occurrence of staphylococci resistant to antimicrobial agents (Pyörälä and Pyörälä, 1994). During lactation, the cure rate of subclinical mastitis ranges widely, and this variability can be due to the choice of antimicrobial agent as well as to factors associated with the infected cow and the quarter. Therefore, cure is likely not a random event (Schukken et al., 1994). Antimicrobial therapy is a primary tool for controlling staphylococcal mastitis, and antimicrobial susceptibility tests can guide the veterinarian in selecting the most appropriate antimicrobial agent for treatment of IMI by *Staph. aureus*. However, despite a variety of available antimicrobial agents, success in the treatment of *Staph. aureus* mastitis, particularly during lactation, is still very low. In fact, *Staph. aureus* pathogens have many characteristics that make them difficult targets for antimicrobial therapy (Sol et al., 2000). For example, they can penetrate the phagocytic cells and survive inside them. This process shields the pathogens from some of the activity of antimicrobial agents, even with drugs that can penetrate the cells. The objective of the present study was to evaluate the antimicrobial resistance patterns of *Staph. aureus* isolates collected from IMI in cows from 44 commercial herds in Italy.

Sixty-eight isolates of *Staph. aureus* taken from individual quarters of 68 cows with subclinical mastitis were used. The isolates were obtained during 2004, from milk samples collected from animals belonging to 42 dairy herds from different regions of Italy. Animals were selected on the basis of having quarter milk SCC greater than 400,000 cells/mL but no sign of clinical mastitis. Herds were selected to represent different prevalences of *Staph. aureus* IMI, ranging from 1 to 100%. The number of isolates for each herd was determined according to the number of lactating animals in each herd: One single isolate was collected from 26 small herds ( $\leq 50$  lactating cows), 2 isolates were collected from 11 medium-sized herds ( $\leq 100$  lactating

Received October 17, 2005.

Accepted February 28, 2006.

<sup>1</sup>Corresponding author: paolo.moroni@unimi.it

<sup>2</sup>Current address: Animal Health and Production Section, Joint IAEA/FAO Division, International Atomic Energy Agency, A-1400 Vienna, Austria.

cows), and more than 2 isolates were collected from 5 large herds (>100 lactating cows). All isolates were identified on the basis of morphology, hemolysis pattern, and gram staining. The gram-positive cocci were tested for catalase and coagulase production. The species were identified by biochemical tests and by the API Staph System (BioMérieux, Rome, Italy) and were then stored at  $-70^{\circ}\text{C}$  in a nutrient broth enriched with 15% glycerol.

In the present study, the antibiotics were selected by considering the approved and most frequently used drugs for the treatment of bovine IMI in Italy, even if these drugs were not representative of a particular antibiotic class. The antimicrobial agents selected and provided by manufacturers as powders were penicillin G (PEN), ampicillin (AMP), amoxicillin (AMX), AMX + clavulanate, cloxacillin (CLX), cephalonium (CFL) and cefoperazone (CFP), kanamycin, oxytetracycline, doxycycline, and lincomycin (LIN). The antimicrobial agents were dissolved in suitable solvents to make stock solutions and then diluted in sterile distilled water according to the methods recommended by the Clinical Laboratory Standards Institute (2002). Minimum inhibitory concentration tests were performed according to the microdilution broth method, as recommended by the Clinical Laboratory Standards Institute (2002a), using U-bottomed 96-well microtiter plates. Serial 2-fold dilutions of the antimicrobial agents were prepared starting from the stock solution of each drug. The dilution schemes differed according to the antimicrobial agent. Inocula were prepared by diluting an overnight (16 to 18 h) Mueller–Hinton broth culture in buffered saline solution to a density of 0.5 on the McFarland turbidity scale and finally diluting it again 40-fold before testing. The MIC was defined as the lowest concentration of the antimicrobial agent at which the bacterial growth was completely inhibited. A reference strain (*Staph. aureus*, ATCC 29213) was inoculated as a control in each plate. The MIC data were summarized, calculating the MIC values for which the isolates were equal to or below 50 and 90% ( $\text{MIC}_{50}$  and  $\text{MIC}_{90}$ , respectively), as well as the minimum and maximum MIC values (range). Resistance and susceptibility, for most of the antimicrobial agents tested, were determined according to Clinical Laboratory Standards Institute (2002a) MIC breakpoints for veterinary pathogens. The interpretive criteria, however, were based on MIC data and drug pharmacokinetic data obtained in humans (taken from Clinical Laboratory Standards Institute, 2002b). *Staphylococcus aureus* was also tested for  $\beta$ -lactamase production by the nitrocefin test (Cefinase, Becton, Dickinson and Co., Sparks, MD).

Numerous data are available in the literature on the susceptibility to antimicrobial agents of *Staph. aureus*

isolates collected from IMI of dairy cattle (Craven et al., 1986; Watts and Yancey, 1994; Owens et al., 1997; Makovec and Ruegg, 2003; Pengov and Ceru, 2003; Tikofsky et al., 2003). Nevertheless, obtaining continually updated MIC values is important to prevent the use of ineffective antimicrobial drugs. Table 1 reports  $\text{MIC}_{50}$  and  $\text{MIC}_{90}$  values of the selected antibiotics against the *Staph. aureus* isolates examined in the present study. All values obtained with the control strain were within the expected ranges for all antimicrobial agents tested. Of the 68 isolates evaluated, none were susceptible to all antibiotics and 64 (94%) were resistant to 3 or more antimicrobial agents. No isolates were resistant to CLX. The  $\beta$ -lactams (penicillins and cephalosporins) are widely used for intramammary treatment of bovine mastitis, but in the present experiment we observed a very poor activity of PEN ( $\text{MIC}_{50} = 0.5$  and  $\text{MIC}_{90} = 2,000 \mu\text{g/mL}$ ). The  $\text{MIC}_{90}$  for PEN reported in previous studies ranged from  $<0.06$  to  $>100 \mu\text{g/mL}$  (De Oliveira et al., 2000; Erskine et al., 2004). In our study, 47 isolates (69%) of *Staph. aureus* were PEN resistant. This proportion was greater than those reported for comparable studies in Argentina (40%; Gentilini et al., 2000), the United States (38.4 to 60.9%; Erskine et al., 2002), and Finland (50%; Myllys et al., 1998), but was lower than that reported for strains isolated from mammary parenchymas of slaughtered dairy cows in Brazil (75%; Costa et al., 2000). Similarly, in the penicillin group, AMP and AMX had very poor in vitro activity ( $\text{MIC}_{50} = 2$  and  $4 \mu\text{g/mL}$ , and  $\text{MIC}_{90} = 500$  and  $1,000 \mu\text{g/mL}$ , respectively), and 100 and 98.5% of the isolates were resistant to these respective antimicrobial agents. Results from other studies for  $\text{MIC}_{90}$  for AMP differed remarkably from our findings; they ranged from only 0.5 to  $4 \mu\text{g/mL}$  (Watts and Salmon, 1997; De Oliveira et al., 2000). This high level of resistance was probably related to the presence of strong  $\beta$ -lactamase producers among the tested staphylococcal isolates. The in vitro data confirmed the influence of  $\beta$ -lactamase production on the microbial susceptibility to  $\beta$ -lactams in general and to PEN in particular. In fact, the difference between  $\text{MIC}_{50}$  and  $\text{MIC}_{90}$  values, with reference to PEN, AMP, and AMX, correlates very well with the identification of 28  $\beta$ -lactamase-producing isolates (58%). Recalculation of the  $\text{MIC}_{90}$  without these strains yielded values of  $0.5 \mu\text{g/mL}$  for PEN and  $4 \mu\text{g/mL}$  for both AMP and AMX. On the other hand, the  $\beta$ -lactamase-resistant CLX and amoxicillin + clavulanate (a widely used  $\beta$ -lactamase inhibitor) were both highly effective, with  $\text{MIC}_{50}$  of 0.25 and 1 to  $0.5 \mu\text{g/mL}$  and  $\text{MIC}_{90}$  of 0.5 and 8 to  $4 \mu\text{g/mL}$ , respectively.

The cephalosporins are usually classified into 3 different generations on the basis of their respective antimicrobial spectra. In the present study, CFL and CFP

Download English Version:

<https://daneshyari.com/en/article/2440997>

Download Persian Version:

<https://daneshyari.com/article/2440997>

[Daneshyari.com](https://daneshyari.com)