

## Short Communication: Canola Meal as a Substitute for Cottonseed Meal in Diet of Midlactation Holsteins

S. M. Maesoomi,\* G. R. Ghorbani,\* M. Alikhani,\* and A. Nikkhah†<sup>1</sup>

\*Department of Animal Science, Isfahan University of Technology, Isfahan 84156, Iran

†Department of Animal Science, University of Manitoba, Winnipeg, MB, Canada R3T 2N2

### ABSTRACT

The growing demand by humans for monounsaturated vegetable oils has provided canola meal (CM) for use in dairy diets because it possesses an excellent nitrogen profile for rumen microbes. Six midlactation cows were used in a replicated 3 × 3 Latin square design with 3 periods of 20 d each. Treatments included diets with 1) CM, 2) 50% CM + 50% cottonseed meal (CSM), and 3) CSM. Total crude protein (CP), nonprotein nitrogen, and rapidly degradable true protein (% of CP) were greater in CM than in CSM. The neutral and acid detergent fibers, slowly degradable true protein, and unavailable CP were lower in CM than in CSM. Daily feeding of 3.4 kg of CM instead of 5.6 kg of CSM enhanced milk percentage of protein and SNF, and improved total tract digestibility of dry matter and CP. Therefore, CM offers an economical substitute for CSM in midlactation diets when commercial access, cost, and quality of CSM are variable.

**Key words:** canola meal, cottonseed meal, protein fraction, midlactation cow

Cottonseed meal (CSM) is used in dairy diets because of its lower price, limited use in nonruminant diets, and greater escape protein when compared with soybean meal (Coppock et al., 1987). However, limited access to CSM superimposed on its low Lys and Met concentrations and variable gossypol content appear to limit its dietary use for lactating cows (Coppock et al., 1987).

In view of the progressive human demand for vegetable oils high in monounsaturated fat and low in ratio of n-6 to n-3 polyunsaturated fatty acids, canola cultivars with largely reduced levels of glucosinolates (GT) have been developed. This progress in plant breeding has concomitantly served the dairy diets with canola meal (CM; Sanchez and Claypool, 1983). Meanwhile, the high ruminal degradability of CP has led researchers

to process CM (Khorasani et al., 1993). The processing of CM aims to minimize the ruminal surplus of soluble N and optimize the small intestinal supply of the most limiting AA for milk synthesis (i.e., Met, Lys, His, and Phe; Schingoethe, 1996). The benefits of increased RUP due to processing CM seem, however, to be offset by conducting the costly, standardized processing techniques. Besides, CM already contains a RUP residue whose strata of AA are remarkably close to that of milk protein (Piepenbrink and Schingoethe, 1998). More importantly, CM possesses an excellent RDP profile that may stimulate microbial growth in the rumen (Piepenbrink and Schingoethe, 1998). Such efficient provision of AA and peptides, readily incorporated into microbial mass (Bach et al., 2005), becomes more prominent in the light of recent timely emphases (Santos et al., 1998; Bach et al., 2005) on significance of RDP for efficient microbial metabolism and milk synthesis.

In the current animal feed market in Iran, CM is similar in price to CSM (185 vs. 195 toman/kg; US \$1 = 916 toman), but is much higher in CP (44 vs. 29%). In other words, on average, CM is cheaper than CSM by about 5 and 60% per kg of DM and CP, respectively. The antinutritive factors in CSM and the adverse environmental impacts of feeding high amounts of unavailable N are additional concerns that justify introducing a commercial alternative for CSM. Therefore, we designed this study to 1) address the significance of using more economical protein sources (e.g., CM) by the dairy industry in Iran, and 2) underline the importance of dietary RDP in the form of CM for lactating cows. We hypothesized that full dietary substitution of highly degradable CP of CM for the less degradable CP of CSM maintains productivity without affecting DMI, nutrient digestibility, or circulating thyronines and albumin. We fractionated CP of CM and CSM, and assessed the effect of dietary substitution of CM for CSM on performance of midlactation cows.

Six (3 primiparous and 3 multiparous) Holstein cows (600 ± 50 kg of BW), 120 ± 4 DIM with 34 ± 2 kg/d (mean ± SE) initial milk yield were used in a replicated 3 × 3 Latin square design. Cows were housed in individual stalls (4 × 4 m) in a roofed area at the Dairy Facilities of Lavark Research Station (Isfahan University of Tech-

Received July 5, 2005.

Accepted November 15, 2005.

<sup>1</sup>Corresponding author: umnikkha@cc.umanitoba.ca

**Table 1.** Ingredients and nutrient composition of the diets (DM basis)

Item	Dietary treatment <sup>1</sup>		
	CM	CC	CSM
Alfalfa hay	49.6	46.2	39.5
Corn grain	13.7	13.7	13.8
Barley grain	16.1	16.2	16.3
Canola meal	14.3	9.0	—
Cottonseed meal	—	9.0	23.6
Whole linted cottonseed	2.4	2.4	2.4
Wheat bran	2.4	2.0	2.9
Sodium bicarbonate	0.9	0.9	0.9
Vitamin and mineral supplement <sup>2</sup>	0.6	0.6	0.6
NE <sub>L</sub> (Mcal/kg) <sup>3</sup>	1.50	1.51	1.52
CP, %	17.2	17.3	17.2
RDP, <sup>3</sup> %	12.4	12	11.4
RUP, <sup>3</sup> %	4.8	5.3	5.8
NDF, %	36.4	37.5	37.8
ADF, %	21.6	22.7	23.2
NFC, <sup>4</sup> %	37.8	37.0	36.1
EE, %	3.0	3.1	3.3
Ca, %	1.0	0.9	0.8
P, %	0.4	0.5	0.5
DCAD, <sup>5</sup> mEq/kg	282	290	303

<sup>1</sup>CM = Canola meal; CC = 50% canola meal + 50% cottonseed meal; CSM = cottonseed meal.

<sup>2</sup>Contained (per kilogram): 19.6% Ca, 9.6% P, 7.1% Na, 1.9% Mg, 0.3% Fe, 0.03% Cu, 0.2% Mn, 0.3% Zn, 100 ppm Co, 100 ppm I, 0.1 ppm Se,  $50 \times 10^5$  IU of vitamin A,  $10 \times 10^5$  IU of vitamin D, and 0.1 g of vitamin E.

<sup>3</sup>Calculated from NRC (2001).

<sup>4</sup>NFC = 100 - (NDF% + CP% + ether extract% + ash%).

<sup>5</sup>DCAD = (Na<sup>+</sup> + K<sup>+</sup>) - (Cl<sup>-</sup> + S<sup>-</sup>).

nology, Iran) from July to September 2004. The experiment consisted of three 20-d periods. The first 15 d of each period were for acclimation to the experimental conditions, and the last 5 d for data collection. Cows were allowed 2 h of outdoor exercise every morning for the entire experiment.

Treatments included the diets containing: 1) CM; 2) 50% CM + 50% CSM; or 3) CSM (Table 1). Diets were formulated (NRC, 2001) to contain equal levels of NE<sub>L</sub>, CP, and NDF (Table 1). Cows were offered TMR twice daily at 0800 and 1400 h at rates to allow for 5 to 10% orts. Fecal samples were obtained from the rectum, and urine was sampled by manual stimulation of the vulva at 1100 h every day of the collection period. Acid insoluble ash was used as an internal marker to determine the apparent total tract digestibility of nutrients. The CP of CM and CSM was fractionated (Table 2) according to the Cornell Net Carbohydrate and Protein System (Sniffen et al., 1992), as described by Licitra et al. (1996). On the last day of each period, 10 mL of blood was collected via jugular vein. The plasma thyroid hormones (triiodothyronine and thyroxine) were measured by standard radioimmunoassay using commercial kits (Kavoshyar Kit, clinical diagnostic, Isfahan, Iran). The serum was assayed for albumin using cellulose acetate

electrophoresis. Cows were milked thrice daily in the milking parlor at 0500, 1300, and 2100 h. Milk samples were analyzed (AOAC, 1990) for fat, protein, SNF, lactose, and TS using a Milk-O-Scan (134 BN Foss Electric, Hillerød, Denmark).

Data were analyzed as a linear mixed model with PROC MIXED of SAS (SAS Institute, 1999). The initial model included the fixed effects of treatment, period, parity, treatment  $\times$  parity, and random effect of cow nested within parity. Because of insignificance ( $P > 0.30$ ), parity and its interaction with treatment were excluded from the model. Period was modeled as a repeated factor with first-order autoregressive [AR (1)] covariance structure to account for the correlation of repeated measures on the same subject (Tempelman, 2004).

The CM contained significantly less ( $P < 0.01$ ) NDF and ADF than did CSM (Table 2). The CM was, however, higher in CP but lower in ether extract than was CSM ( $P < 0.01$ ). The fractionation of CP revealed that NPN (fraction A) and rapidly degradable true protein (fraction B<sub>1</sub>) were greater ( $P < 0.01$ ) in CM than in CSM. However, the slowly degradable (escape) true protein (fraction B<sub>2</sub>), total true protein, and unavailable N (fraction C) were lower ( $P < 0.01$ ) in CM than in CSM (Table 2). Daily DMI, feed efficiency, actual milk yield, milk fat content and yield, and the yield of 4% FCM were not altered by treatments (Table 3). Milk percentage of protein and SNF were increased by dietary substitution of CM for CSM (Table 3).

Complete substitution of dietary CP of CM for that of CSM in midlactation diet improved the apparent total tract digestibility of dietary DM ( $P = 0.01$ ), CP ( $P < 0.01$ ), and OM ( $P = 0.07$ ; Table 4). The concentrations of albumin and thyronines [triiodothyronine (T<sub>3</sub>) and thyroxine (T<sub>4</sub>)] in circulating blood were not impacted by treatments (Table 4).

The significantly different CP content of CM and CSM (Table 2) was in disagreement with Sniffen et al. (1992) who reported comparable CP contents for CM (42.3%) and CSM (44.8%). The discrepancy between the 2 studies could be explained by the lower NDF and ADF in CM than in CSM in our study. The CSM used in the current trial was higher in fiber, but lower in NPN and CP compared with that used in other studies (Sniffen et al., 1992; Licitra et al., 1996). Such interstudy differences in nutrient composition of CSM could be attributed to the variety of the cottonseeds produced under specific agronomic conditions. The processing conditions (e.g., temperature, pressure, and chemicals used) during the extraction of oil from oilseeds could also affect nutrient composition of protein meals. Moreover, the high levels of N associated with ADF in CSM of the present study should have, at least partly, contributed

Download English Version:

<https://daneshyari.com/en/article/2441584>

Download Persian Version:

<https://daneshyari.com/article/2441584>

[Daneshyari.com](https://daneshyari.com)