



CASE STUDY: Feed Intake and Performance of Heifers Sired by High- or Low-Residual Feed Intake Angus Bulls¹

J. Minick Bormann,² PAS, D. W. Moser, and T. T. Marston,³ PAS

Department of Animal Sciences and Industry, Kansas State University, Manhattan 66506

ABSTRACT

The objective of this project was to investigate the effects of selecting sires for residual feed intake (RFI) on the performance of their daughters. Bulls with low or high estimated breeding values (EBV) for RFI were selected from the Angus Society of Australia sire summary and mated to Angus cross commercial cows at the Kansas State University Cow-Calf Unit in 2005 and 2006. The average EBV of low- and high-RFI bulls were -0.55 and 0.27 kg DM, respectively. Heifers born in 2006 were tested for feed intake in 2 groups ($n = 24$, $n = 26$), and heifers born in 2007 ($n = 42$) were sent to a commercial bull test facility for feed intake and BW gain tests. Body weights were collected every 14 d and used to calculate midtest BW and ADG. Actual feed intake was regressed on midtest metabolic BW and ADG to calculate an expected feed intake for each heifer. Residual feed intake was calculated by subtracting the expected intake from the actual intake. There were no significant

differences between heifers sired by low- or high-RFI EBV bulls in RFI, feed intake, G:F, or BW gain ($P > 0.05$). Heifers in this study were being developed on a less energy-dense diet than the diet used to rank their sires. Genetic differences in RFI calculated in growing bulls may not have been expressed on the lower plane of nutrition of these developing heifers.

Key words: beef heifer, breeding value, residual feed intake

INTRODUCTION

Feed costs continue to be a large portion of the total expenses of a beef cattle operation. Much of the focus of genetic improvement in beef cattle has focused on improved quantity and quality of outputs of the production system, such as increased growth rate and carcass quality. Feed continues to be a major cost to the cow-calf, stocker, and feedlot sections of the industry, and there has been much recent interest in improving the feed efficiency of beef production.

Koch et al. (1963) first proposed using residual feed intake (RFI) as a measure of efficiency in beef cattle. Residual feed intake is defined as the difference between actual feed intake

and predicted feed intake based on BW and growth. Animals with negative RFI eat less than is expected for their size and level of production, and are more efficient. Residual feed intake appears to be moderately heritable, with literature estimates for Angus cattle ranging from 0.39 to 0.51 (Arthur et al., 2001a; Moore et al., 2003, 2005). After one generation of selection in an Australian study, bulls, heifers, and steers from the high-RFI ($n = 27$) and low-RFI ($n = 30$) lines had differences in RFI ($P < 0.05$) and in actual intake ($P < 0.05$), showing response to selection in one generation (Herd et al., 1997; Richardson et al., 1998). The performance of animals selected for RFI has not been evaluated outside Australia, so the objective of this study was to investigate the effects of selecting for RFI on the feed efficiency and performance of beef heifers in a typical Midwestern US beef production system.

MATERIALS AND METHODS

This study was conducted under guidelines established by the Kansas State University Institutional Animal Care and Use Committee. In 2005 and 2006, Angus-based commercial cows at the Kansas State University

¹Kansas Agricultural Experiment Station (Manhattan) Publication 09-285-J

²Corresponding author: jbormann@ksu.edu

³Current address: Northeast Research and Extension Center, University of Nebraska, Norfolk, NE 68701.

Cow-Calf Unit were bred to Angus sires that had RFI estimated breeding values (EBV) published by the Angus Society of Australia (Armidale, New South Wales, Australia). The distribution of progeny by sire is shown in Table 1. The sires used had a combination of progeny with feed intake phenotypes and IGF-1 phenotypes. Accuracy of sire EBV ranged from 0.34 to 0.78 at the time they were used. One sire was used only in yr 1, whereas 4 sires were used only in yr 2, and 4 sires were used in both years. Heifers born in 2006 were tested in winter and spring of 2007 using Calan gates (American Calan, Northwood, NH) in 2 groups ($n = 24$, test 1; $n = 26$, test 2). In 2008, heifers that were born in 2007 were tested in the spring in a GrowSafe system (Airdrie, Alberta, Canada) in one group ($n = 42$, test 3). Heifers were removed from the study for failure to train to the equipment, morbidity, pregnancy, and death. After a 14-d adjustment period, feed intake was measured for 42 d, BW gain was measured for 58 d in 2007, and both intake and BW gain were measured for 57 d in 2008, following the recommendations of Archer et al. (1997) and Wang et al. (2006). These studies showed that 35 d was an adequate time to obtain accurate measurements of feed intake, whereas BW gain should be measured for approximately 60 d. In both years, heifers were allowed ad libitum intake of a high-roughage, complete diet [approximately 2.63 Mcal ME/kg DM (11.00 MJ ME/kg DM) in 2007 and

1.9 Mcal ME/kg DM (7.95 MJ ME/kg DM) in 2008]. Test 1 and 2 heifers were scanned by ultrasound for body composition at the end of the BW gain test. Residual feed intake was calculated by the method of Arthur et al. (2001b) within each test group. Body weights were collected every 14 d and used to calculate midtest BW and ADG. Actual DMI was regressed on midtest metabolic BW and ADG to calculate an expected DMI for each heifer. The model for expected feed intake was

$$y_i = b_0 + b_1 \text{ADG}_i + b_2 \text{WT}_i + e_i,$$

where ADG_i is the ADG of animal i , WT_i is the midtest metabolic BW of animal i , and e_i is the error. Expected DMI was calculated within each contemporary (test) group separately. Residual feed intake was calculated by subtracting the expected intake from the actual intake. Data were analyzed using SAS software (SAS Institute Inc., Cary, NC). The model for differences between high- and low-RFI sire groups included test group as a fixed effect and sire within RFI group as a random effect. For analysis of ultrasound traits, test group was a fixed effect, sire within RFI group was a random effect, and sire EBV for the ultrasound trait was included as a covariate. The model for regression of heifer RFI phenotype on sire RFI EBV included the fixed effect of test group. For both sire RFI EBV group and test group, Levene's test for homogeneity of variance was per-

formed on RFI, ADG, DMI, and G:F. To further investigate the relationship between diet and RFI, the differences between sire groups were analyzed within test group. For analysis within group, sire within RFI group was included as a random effect.

RESULTS AND DISCUSSION

Least squares means for daily RFI, ADG, daily DMI, and G:F for the sire groups within test groups are shown in Table 2. There was no difference between the 2 lines for feed intake or RFI either within or across tests. The phenotypic variances between sire groups and test groups were equal ($P > 0.1471$) for RFI, ADG, DMI, and G:F (Table 3). The weighted averages of sire RFI EBV were 0.29 kg for inefficient sires and -0.40 for efficient sires. Therefore, we would expect the progeny groups from these sires to differ in RFI by 0.35 kg (difference in EBV divided by 2). The actual difference in phenotypic RFI between heifers sired by low- or high-RFI EBV bulls was 0.12 kg. The regression of heifer RFI on sire RFI EBV was 0.17 ± 0.32 kg. Theoretically, this value is expected to be 0.5 because half the breeding value of the sire is passed to progeny. These results are in contrast with previous results from the Australian selection experiment. After one generation of selection, they found significant differences ($P < 0.05$) in feed intake and RFI between high and low selection lines in bulls, heifers, and steers (Herd et al., 1997; Richardson et al., 1998). Herd et al. (1997) reported significant differences in intake, of $1,262 \pm 25$ kg for the low-RFI line and $1,354 \pm 24$ kg for the high-RFI line, and total RFI of -19 ± 10 kg for the low-RFI line and $+49 \pm 9$ kg for the high-RFI line. Richardson et al. (1998) found significant differences in daily DMI, of 9.22 ± 0.18 for the low-RFI line and 9.78 ± 0.16 for the high-RFI line, and daily RFI of -0.20 ± 0.11 for the low-RFI line and $+0.17 \pm 0.10$ for the high-RFI line. There may be an effect of plane of nutrition and growth rate of the cattle on RFI. In the Australian stud-

Table 1. Sire residual feed intake (RFI) group (I = inefficient, E = efficient), estimated breeding value (EBV), and number of daughters

Sire	Sire RFI group	EBV, kg DM	Daughters, no.
1	I	0.29	18
2	I	0.26	10
3	I	0.30	3
4	I	0.31	21
5	I	0.19	4
6	E	-0.54	8
7	E	-0.72	7
8	E	-0.41	7
9	E	-0.51	14

Download English Version:

<https://daneshyari.com/en/article/2454354>

Download Persian Version:

<https://daneshyari.com/article/2454354>

[Daneshyari.com](https://daneshyari.com)