



# Effects of *Morinda citrifolia* (Noni) on CD4<sup>+</sup> and CD8<sup>+</sup> T-Cell Activation in Neonatal Calves

V. J. Brooks,\* M. Schäfer,\* P. Sharp,† J. Xu,\* J. Cai,\* N. S. Keuler,‡ R. G. Godbee,§ S. F. Peek,\* R. D. Schultz,† M. Suresh,† and B. J. Darien\*<sup>1</sup>

\*Department of Medical Sciences, †Department of Pathobiological Sciences, and ‡Department of Computing and Biometry, University of Wisconsin, Madison 53706; and §Department of Animal Biotechnology/Veterinary Medicine, Fleischmann Agriculture, University of Nevada, Reno 89557

## ABSTRACT

Developmental immaturity of the immune system renders neonatal calves vulnerable to infectious causes of morbidity and mortality. Juice made from the *Morinda citrifolia* fruit (noni) reportedly has immune enhancing, antibacterial, and antiinflammatory effects. The objective of this study was to evaluate the potential immune-modulating effects of feeding noni puree to neonatal calves on T-cell activation on CD4<sup>+</sup>, CD8<sup>+</sup>, and  $\gamma\delta$  T cells. Sixteen newborn Holstein bull calves were acquired in pairs from local dairies and confirmed to have adequate passive transfer at 24 h of age. The calves were divided into 2 groups: group 1 ( $n = 8$ ) was composed of control calves, and group 2 ( $n = 8$ ) received 30 mL of noni puree twice daily in milk replacer. Blood samples were collected from each calf on d 0, 3, 7, and 14. Mitogen-induced activation of CD4<sup>+</sup>, CD8<sup>+</sup>, and  $\gamma\delta$  T-cell receptor-positive T cells was evaluated by measurement of the up-regulation of the interleukin 2 receptor, CD25, using 2-color flow cytometry. Concanavalin A and phytohemagglutinin were used as global mitogens. Noni

puree-fed calves had an increase in CD25 expression on concanavalin A-stimulated CD4<sup>+</sup> ( $P = 0.03$  for parametric and nonparametric analysis) and CD8<sup>+</sup> T cells ( $P = 0.04$  for parametric analysis) on d 3 of the study, or approximately 4 to 5 d postpartum. There was also an effect over time for CD8<sup>+</sup> T cells ( $P = 0.03$  for nonparametric analysis). Further studies are warranted to determine the cellular mechanisms responsible for these findings and whether noni supplementation in dairy calves translates to improved health and well-being.

**Key words:** *Morinda citrifolia*, neonatal calf, noni, immunomodulation, T cell

## INTRODUCTION

The developmental immaturity of the immune system of the neonatal calf presents a predisposing factor toward increased morbidity and mortality. The naïve immune system of a calf is constantly challenged by a variety of viral and bacterial pathogens found in its environment, and as a result, approximately 8.7% of dairy heifers born alive die before weaning (USDA, 2002).

The *Morinda citrifolia* fruit (noni) is a natural product that has a broad

range of immune-enhancing effects, including antibacterial, antiinflammatory, analgesic, antioxidant, and antitumor effects (Wang et al., 2002; Furusawa et al., 2003). Noni has been shown to induce the release of several immune mediators, many of which have beneficial stimulatory effects and aid in the maturation of the neonatal immune system (Hirazumi and Furusawa, 1999).

Previously, we examined the effects of feeding calves noni puree for the first 2 wk of life on bacterial killing via an ex vivo whole-blood bactericidal assay (Schäfer et al., 2008). That study demonstrated that noni-supplemented calves had more *Escherichia coli* killing power at d 14 compared with control calves, and that enhanced bactericidal activity in noni-supplemented calves increased over time. We postulated that these results may have been due to noni positively affecting pathogen-associated molecule pattern recognition via Toll-like receptor 4, resulting in an increase in bacterial engulfment, phagocytosis, or both, or a cytokine-mediated effect via enhanced T-helper cell recognition of antigens expressed on the MHC II. Because both of these mechanisms result in the activation of T cells (whether directly or as a downstream

<sup>1</sup>Corresponding author: darienb@svm.vetmed.wisc.edu

effect), we decided to examine whether noni supplementation would also have an effect on T-cell activation in neonatal calves.

In the current study, we evaluated the effects of noni supplementation in newborn calves during the first 2 wk of life on T-cell activation in response to the potent mitogens concanavalin A (**ConA**) and phytohemagglutinin (**PHA**) by 2-color flow cytometry. The CD4<sup>+</sup>, CD8<sup>+</sup>, and  $\gamma\delta$  T-cell receptor-positive (**TCR**<sup>+</sup>) T cells were deemed activated if they expressed CD25, the interleukin (**IL**)-2 receptor, which plays a crucial role in lymphocyte proliferation and differentiation (Waters et al., 2003).

## MATERIALS AND METHODS

### Reagents

Before assay, acid citrate dextrose-A was prepared with 2.2 g sodium citrate (dehydrate), 0.8 g citric acid (monohydrate), 2.5 g dextrose, and 100 mL H<sub>2</sub>O. A cell-lysing solution of pH 7.2 was prepared by dissolving 1.5 g (10.6 mM) Na<sub>2</sub>HPO<sub>4</sub> and 0.32 g (2.7 mM) NaH<sub>2</sub>PO<sub>4</sub> in 1 L H<sub>2</sub>O. A restoring solution of pH 7.2 was similarly prepared by dissolving 1.5 g (10.6 mM) Na<sub>2</sub>HPO<sub>4</sub>, 0.32 g (2.7 mM) NaH<sub>2</sub>PO<sub>4</sub>, and 27 g (462.0 mM) NaCl in 1 L H<sub>2</sub>O.

### Animals

Animals for this project were obtained from 6 local dairies. The study and procedures were reviewed and approved by the Institutional Animal Care and Use Committee of the University of Wisconsin-Madison. Within 12 h of birth, 16 newborn Holstein bull calves that had already received 4.0 L of pooled colostrum arrived in pairs at the veterinary medical teaching hospital, where they were housed in individual pens without nose-to-nose contact. The animals were assigned to control or treatment groups in the order in which they were removed from the calf trailer by hospital personnel, who had no further involvement with the study.

Upon arrival, every animal received a physical examination, followed by daily examinations for the duration of the study. At each examination, the following parameters were measured: temperature, ease of cough induction, fecal consistency, and presence and severity of ocular or otic abnormalities. A calf health score was assigned to each animal using the scoring system described previously in Schäfer et al. (2008). Veterinarians blinded to treatment groups oversaw calf health evaluations. Any calf receiving a total health score of  $\geq 5$  for 3 consecutive days was removed from the study and treated appropriately. Adequate passive transfer (IgG >1,000 mg/dL) was confirmed for all calves in the study with the IgG Midland Quick Test Kit (Midland Bioproducts Corp., Boone, IA) at approximately 24 h of age.

Calf pairs consisted of 1 noni puree-fed calf and 1 control calf. Calves were bottle-fed 2 L of nonmedicated milk replacer (Calf Glo, Vita Plus Corp., Madison, WI) reconstituted according to the label of the manufacturer twice daily for the first 7 d and 2.5 L twice daily from d 8 to 14. Noni puree-fed calves received 30 mL noni puree twice a day in milk replacer. Calves had access to 125 g calf starter and 4 L fresh water per day. Of the 8 calf pairs, only 3 had differing health scores between calves on d 0: in 2 pairs the score was 1 unit greater for the control calf, and in 1 pair it was 1 unit greater for the noni puree-fed calf. No calves were removed from the study because of health reasons.

### Peripheral Blood Mononuclear Cell Isolation

The acid citrate dextrose-A anticoagulated blood (25 mL) was collected in Vacutainer tubes (Becton Dickinson, Franklin Lakes, NJ) from the jugular vein from each calf on d 0, 3, 7, and 14. Day-0 samples were obtained from each calf between 36 and 48 h of age and before the first feeding of puree. All other samples were obtained before the respective feeding.

The blood was diluted 1:1 with PBS, and the buffy coat was harvested after centrifugation at  $920 \times g$  for 30 min at 25°C, diluted with plasma, layered over 1.083 Ficoll-Histopaque (Sigma, St. Louis, MO), and centrifuged at  $1,380 \times g$  for 30 min at 25°C. The peripheral blood mononuclear cell (**PBMC**) interphase layer was harvested and the remaining red blood cells were exposed to a lysing solution for 1 min, which was neutralized with a restoring solution (final ratio 1:1:1). The PBMC were washed twice, once with PBS and once with RPMI-1640 (Mediatech Inc., Herndon, VA) at  $280 \times g$  for 10 min, and were subsequently resuspended in RPMI-1640 with 20% fetal calf serum (**FCS**). Cells were stained with Trypan blue to confirm cell viability (>95%) and adjusted with RPMI-1640 with 20% FCS to  $5 \times 10^6$  cells/mL.

### Flow Cytometry

For flow cytometry analysis,  $50 \mu\text{L}$  of PBMC ( $5 \times 10^6$  cells/mL) were cultured on a Costar flat-bottomed, tissue culture-treated, 96-well plate (Corning Inc., Corning, NY). Unstimulated control wells received 100  $\mu\text{L}$  RPMI-1640 + 20% FCS. Stimulated sample wells received 100  $\mu\text{L}$  ConA (final concentration of 6.7  $\mu\text{g/mL}$ ; Sigma) or 100  $\mu\text{L}$  of the M form of PHA (Gibco, Invitrogen Corp., Carlsbad, CA) diluted 1:200 in RPMI-1640 + 20% FCS. Plates were incubated for 72 h at 37°C at 5% CO<sub>2</sub>.

After incubation, the cells were transferred into 12  $\times$  75 mm polystyrene round-bottomed tubes (Becton Dickinson) and washed with PBS at  $520 \times g$  for 7 min at 25°C. The cells were resuspended and stained for 2-color flow cytometry with the primary mouse anti-bovine antibodies: CD4 (IgM), CD8 $\alpha$  (IgM), TCR1-N6 ( $\gamma\delta$  TCR<sup>+</sup>; IgM), and CD25 (IgG2a; 15  $\mu\text{g/mL}$ ; VMRD, Pullman, WA).

Stained cells were incubated for 20 min at 4°C and washed 3 times with PBS at  $520 \times g$  for 7 min at 25°C. The cells were resuspended again and stained with the secondary antibodies: fluorescein isothiocyanate-conjugated

Download English Version:

<https://daneshyari.com/en/article/2454380>

Download Persian Version:

<https://daneshyari.com/article/2454380>

[Daneshyari.com](https://daneshyari.com)