



Short communication

Assessment of luteal function in goats by ultrasonographic image attribute analysis

E.K. Arashiro^{a,*}, J.F. Fonseca^b, L.G.B. Siqueira^c, C.A. Fernandes^d, F.Z. Brandao^a, E. Oba^e, J.H. Viana^c^a Fluminense Federal University, Niteroi, RJ 24230-340, Brazil^b Embrapa Goat Research Center, Sobral, CE 62011-970, Brazil^c Embrapa Dairy Cattle Research Center, Juiz de Fora, MG 36038-330, Brazil^d University of Alfenas, Alfenas, MG 37130-000, Brazil^e São Paulo State University, Botucatu, SP 18618-000, Brazil

ARTICLE INFO

Article history:

Received 18 December 2009

Received in revised form 9 July 2010

Accepted 21 July 2010

Available online 17 August 2010

Keywords:

Corpus luteum

Echotexture

Goat

Ultrasound

ABSTRACT

The aim of this study was to evaluate the potential of luteal echotexture (mean pixel value and heterogeneity), as a tool for assessing luteal function during different phases of the estrous cycle in Toggenburg goats. Sonographic evaluations of the ovaries were performed daily in nulliparous goats ($n = 21$), using a 5 MHz linear rectal probe, commencing at estrus (day 0). Blood samples were collected daily for plasma progesterone RIA and images recorded on VHS tape and then digitized in TIFF format at a resolution of 1500×1125 pixels. A representative elementary area (REA) of 5625 pixels (0.31 cm^2) of these images was analyzed using custom-developed software, for mean pixel value and heterogeneity. Mean plasma progesterone, luteal area and pixels all reached maximum values at approximately days 13 and 14, during luteogenesis. Luteolysis was characterized by an abrupt decrease in blood progesterone concentration following ovulation, and a gradual decline in luteal area and pixel values. The luteal tissue area was positively correlated with plasma progesterone concentration during both luteogenesis ($r = 0.63$; $P < 0.05$) and luteolysis ($r = 0.50$; $P < 0.05$). Weak correlations were recorded between the mean pixel value and luteal tissue area during luteogenesis ($r = 0.34$; $P < 0.05$) and luteolysis ($r = 0.26$; $P < 0.05$). Similarly, weak correlations between the mean pixel value and plasma progesterone concentration were recorded during luteogenesis ($r = 0.24$; $P < 0.05$) and luteolysis ($r = 0.37$; $P < 0.05$). The pixel heterogeneity was not correlated with luteal tissue area or the plasma progesterone concentration at any stage of the estrous cycle. The results show the association between the corpus luteum echotexture and steroidogenic function to be weak and the present ultrasound technology, to have limited potential in evaluating luteal function in goats.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

Real-time transrectal ultrasonography (RTU) has shed light on important reproductive events, and is considered to be one of the most significant technological advances in animal reproduction research, since the introduction of radioimmunoassay (RIA) (Ginther, 1986). In small ruminants, corpora lutea (CL's) evaluation by RTU is equally as efficient as invasive techniques, such as laparotomy and

* Corresponding author at: Embrapa Dairy Cattle Research Center, Laboratory of Animal Reproduction, Rua Eugenio do Nascimento 610, Bairro Dom Bosco, Juiz de Fora, MG 36038-330, Brazil. Tel.: +55 32 3311 7438; fax: +55 32 3311 7401.

E-mail address: eduardoarashiro@hotmail.com (E.K. Arashiro).

laparoscopy (Viñoles et al., 2004; Simões et al., 2005). Thus morphometric characteristics of the CL (diameter, area or volume), as measured by RTU in association with plasma progesterone concentration, have been extensively used to evaluate CL function in large domestic animals (Veronesi et al., 2002), and only more recently in small ruminants (Gonzalez-Bulnes et al., 2000).

An ultrasonographic image is known to be composed of thousands of picture elements (pixels), that numerically represent 256 shades of gray (0 = black; 255 = white), depending on the brightness (Tom et al., 1998). The density and gross features of the tissue can be assessed by RTU. However these cannot be quantified by the human eye, which can only distinguish 18–20 different shades of gray (Pierson and Adams, 1995). The use of computer algorithms (computer-assisted analysis) permits a quantitative assessment of the intensity of each pixel present on an image, allowing an objective evaluation of the CL and thus overcoming the subjectivity of the visual analyses (Singh et al., 2003).

In the past, real-time ultrasonography has been used for the functional evaluation of different tissues, including testes, ovarian follicles and CL's. Regarding luteal function, previous studies on cows and ewes have demonstrated a relationship between the CL echotexture and the physiological changes during different phases of the estrous cycle (Davies et al., 2006; Siqueira et al., 2009). However, only a few studies have been performed using the technique of computer-assisted analysis to determine luteal function in goats (Simões et al., 2007).

The aim of the present study was to establish and validate a methodology for evaluating goat CL's by computer-assisted ultrasonographic image analysis, and to assess its ability in determining luteal function during different phases of the estrous cycle.

2. Materials and methods

2.1. Experimental design, animals and facilities

The experiment was conducted on a dairy goat farm located near the city of Coronel Pacheco, in the Minas Gerais State of Brazil (21°35'S, 43°15'W, 435 m altitude), during the onset of the natural breeding season (March and April).

Twenty-three nulliparous Toggenburg goats (*Capra hircus*), with an average age of 8 months, a mean live weight of 33.5 ± 1.2 kg and mean body condition score (BCS) of 3.5 ± 0.1 (scale of 1–5) (Suiter, 1994) were used in the trial. Two animals were removed from the experiment due to follicular cysts. All does used in the trial exhibited natural estrus during a 48-h period and were free of any reproductive abnormalities.

The does were kept as a group in a single pen and fed Napier grass (*Pennisetum purpureum* v. *Taiwan*) and a concentrate. Water and minerals were available ad lib.

2.2. Ovarian ultrasonographic evaluation

The ovarian ultrasonographic evaluations were performed daily, by the same technician, from overt estrous identification (day 0), throughout an estrous cycle (21 days) using a portable ultrasound device (Aloka SSD 500, Aloka Co, Tokyo, Japan), equipped with an adapted linear rectal 5 MHz transducer. The focus field (F1 and F2) and proximal, distal and total gains were standardized and uniform throughout the experimental observation period.

The number of ovulations was determined by the monitoring of follicles larger than 5 mm after the onset of estrus, and this was later confirmed by the number of CL's observed by ultrasound. The number and tissue area

of each CL, as well as the presence or absence of a central cavity, was also recorded. In those does with more than one ovulation, luteal tissue area was considered as the sum of the luteal area of all CL's present.

2.3. Image capture and computer-assisted analysis

The capture and image analysis procedures used were as described for cattle (Siqueira et al., 2009). Image capture was performed by coupling a video cassette recorder using a co-axial cable and recording a representative good quality frozen image on the VHS tape at the SP velocity, for approximately 10 s. Recorded images were transferred to a computer equipped with a video capture board and saved in the non-compressed Tagged Image File (TIFF) format, with a resolution of 1500×1125 pixels.

Computer analysis was performed using a custom-developed software (Quantoro®), as previously described (Siqueira et al., 2009). Briefly, the procedure consisted of the selection of a specific region on the CL, followed by software calculation of the brightness intensity of each pixel composing that area – which received a numerical value of between 0 (black) and 255 (white). Images with artifacts, or encoding fluid-filled cavities or ovarian stroma were excluded from the analyses. Luteal echotexture was represented by the mean pixel value and the standard deviation of this value (heterogeneity).

A minimum representative area (REA) was determined beforehand, based on the criteria of Van den Bygaart and Protz (1999) and modified for use in ultrasound images. In the present study, an REA of 75×75 pixels was used – which represents 5625 pixels.

2.4. Blood samples and plasma progesterone concentrations

Blood samples were collected daily via the jugular vein, using vacuum tubes containing sodium EDTA. Samples were centrifuged at $894 \times g$ for 20 min at 5 °C and the plasma samples then aspirated and stored at –20 °C in 1.5 ml tubes, until RIA plasma progesterone analyses.

Plasma progesterone concentration was determined by radioimmunoassay (RIA), using commercial RIA kits (Coat-a-Count®, Medlab) at the Endocrine Laboratory of the Veterinary Medicine and Animal Science of the São Paulo State University (UNESP), Brazil. The assay sensitivity was 0.02 ng/mL, and inter- and intra-assay variation coefficients 2.75% and 1.87%, respectively.

2.5. Statistical analyses

Outcome variables (mean pixel value and heterogeneity, luteal tissue area and plasma progesterone concentration) were evaluated for normality using the Lilliefors test, and analyzed for the main effect of time (day of the estrous cycle or hours from luteolysis). Due to the expected differences between does at the time of natural luteolysis, data were analyzed according to estrous cycle period: luteogenesis (days 0–16), and luteolysis, which was normalized to the time that plasma progesterone concentrations decreased to <1 ng/mL in each goat (–48 to +72 h from progesterone <1 ng/mL). Daily differences between means were compared using the Tukey's test. Correlations between the luteal tissue area and the mean pixel value, as well as plasma progesterone concentrations and the mean pixel value were compared using Pearson's correlation method. The SAS MIXED procedure (version 9.1.3, SAS Institute, Inc. Cary, NC) was used for the statistical analysis. A probability of $P < 0.05$ indicated that a difference was significant, and all results are presented as the mean \pm SEM.

3. Results

The luteal tissue area progressively increased ($P < 0.05$) until day 9 of the estrous cycle (1.26 ± 0.08 cm²), while no significant increase was recorded in the subsequent days. Luteal tissue area reached its maximum value on day 15 (1.65 ± 0.02 cm²). A time effect was observed regarding the plasma progesterone concentration and the levels progressively increased until day 9 ($P < 0.05$), reaching a maximum value on day 14 (8.32 ± 0.40 ng/mL) (Fig. 1). Luteolysis was characterized by an abrupt decrease in the plasma progesterone concentration, which reflected values of below 1 ng/mL, within 24 h. Luteal tissue area was positively cor-

Download English Version:

<https://daneshyari.com/en/article/2457366>

Download Persian Version:

<https://daneshyari.com/article/2457366>

[Daneshyari.com](https://daneshyari.com)