



# Efficacy of a novel oral formulation of sarolaner (Simparica™) against four common tick species infesting dogs in Europe



Thomas Geurden<sup>a,\*</sup>, Csilla Becskei<sup>a</sup>, Sarah Grace<sup>b</sup>, Christina Strube<sup>c</sup>, Pdraig Doherty<sup>b</sup>, Julian Liebenberg<sup>d</sup>, Sean P. Mahabir<sup>e</sup>, Nathalie Sloomans<sup>a</sup>, Anne Lloyd<sup>a</sup>, Robert H. Six<sup>e</sup>

<sup>a</sup> Zoetis, Veterinary Medicine Research and Development, Mercuriusstraat 20, B-1930 Zaventem, Belgium

<sup>b</sup> Charles River Laboratories, Pre-Clinical Services, Glenamoy Co. Mayo, Ireland

<sup>c</sup> Institute for Parasitology, Centre for Infection Medicine, University of Veterinary Medicine Hannover, Buenteweg 17, 30559 Hannover, Germany

<sup>d</sup> ClinVet International (pty) Ltd., Uitsigweg, Bainsvlei, 9338 Bloemfontein, South Africa

<sup>e</sup> Zoetis, Veterinary Medicine Research and Development, 333 Portage St., Kalamazoo, MI 49007 USA

## ARTICLE INFO

### Article history:

Received 30 October 2015

Received in revised form 12 February 2016

Accepted 29 March 2016

### Keywords:

Sarolaner

Isoxazoline

*Dermacentor reticulatus*

Oral

Tick

Dog

Dose confirmation

## ABSTRACT

The efficacy of single oral treatment of sarolaner (Simparica™, Zoetis), a novel isoxazoline compound, was evaluated against four tick species known to commonly infest dogs in Europe. Eight laboratory studies were conducted using adult purpose-bred Beagle dogs. In each study, 16 animals were randomly allocated to one of two treatment groups based on pre-treatment host-suitability tick counts. Dogs were infested with 50 unfed adult *Dermacentor reticulatus* (two studies), *Ixodes hexagonus* (three studies), *Ixodes ricinus* (two studies) or *Rhipicephalus sanguineus* (one study) ticks on Days –2, 5, 12, 19, 26 and 33. On Day 0, dogs were treated orally with placebo or sarolaner tablets providing the minimum dose of 2.0 mg/kg bodyweight and tick counts were conducted 48 h after treatment and after each subsequent weekly re-infestation. There were no treatment-related adverse reactions in any of the studies. Dogs in the placebo-treated group maintained tick infestations throughout the studies. Geometric mean live tick counts were significantly ( $P \leq 0.0001$ ) lower in the sarolaner-treated group compared to the tick counts in the placebo group at all time-points. A single oral administration of sarolaner resulted in 100% efficacy against existing infestations of all tick species except *R. sanguineus*, for which the efficacy was 99.7%, within 48 h. Efficacy against weekly re-infestations was  $\geq 97.5\%$  for all tick species for 35 days.

Thus, a single dose of sarolaner administered orally at the minimum dosage of 2 mg/kg, resulted in  $\geq 99.7\%$  efficacy within 48 h against existing tick infestations, and in  $\geq 97.5\%$  efficacy against weekly re-infestations, for at least 35 days after treatment. These studies confirmed that administration of the minimum dose of sarolaner will provide treatment of existing infestations and give at least one month of control against re-infestation by the common tick species affecting dogs in Europe.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Ticks are one of the most common ectoparasites infesting dogs. Tick infestation can lead to nuisance, alopecia and skin irritation. Heavy infestations can even lead to anemia (Dryden and Payne, 2004). Canine tick infestations are thus of direct veterinary importance to the animal, but are also important in the distribution of tick-borne diseases (Needham and Teel, 1991; Beugnet and Marié, 2009). Ticks are responsible for the transmission of a number of disease agents, of which some are zoonotic and some

cause serious, even life-threatening conditions (Dryden and Payne, 2004). Zoonotic infections include Lyme disease caused by *Borrelia burgdorferi*, which is transmitted by *Ixodes* species (Beugnet and Marié, 2009). Other tick-borne pathogens cause predominantly dog-specific infections, such as *Babesia canis*, primarily transmitted by *Dermacentor* spp., and *Ehrlichia canis*, primarily transmitted by *Rhipicephalus sanguineus* (Chomel, 2011; Dantas-Torres et al., 2012).

Tick prevention has historically been based on the monthly use of acaricidal compounds applied as topical formulations (Dryden and Payne, 2004; Rust, 2005) but recently two isoxazoline compounds for use in dogs have been introduced in Europe that provide treatment and prevention of flea and tick infestations after oral treatment (Robertson-Plouch et al., 2008; Rohdich et al., 2014;

\* Corresponding author.

E-mail address: [thomas.geurden@zoetis.com](mailto:thomas.geurden@zoetis.com) (T. Geurden).

Shoop et al., 2014). Sarolaner (Simparica™, Zoetis) is a novel isoxazoline with potent activity against ectoparasites (McTier et al., 2016). The objective of this series of studies was to evaluate the efficacy of sarolaner against four tick species of major importance in Europe, each of which has potential to transmit disease organisms (Beugnet and Marié, 2009). Eight laboratory studies were conducted to evaluate the efficacy of sarolaner against existing tick infestations and against re-infestations for a period of five weeks after treatment.

## 2. Materials and methods

The eight laboratory studies were conducted to evaluate against the following tick species commonly infesting dogs in Europe: *Dermacentor reticulatus* (ornate dog tick; two studies), *Ixodes hexagonus* (hedgehog tick; three studies), *Ixodes ricinus* (castor bean tick; two studies), and *R. sanguineus* (brown dog tick; one study). All studies were conducted in accordance with the World Association for the Advancement of Veterinary Parasitology (WAAVP) guidelines for evaluating the efficacy of parasiticides for the treatment, prevention and control of flea and tick infestation on dogs and cats (Marchiondo et al. 2013) and complied with Good Clinical Practices (VICH guideline GL9, 2000). All studies were approved by the Zoetis Ethical Review Committee and by the study site's Ethical Review Committee.

### 2.1. Animals

All dogs used in these studies had not been treated with an ectoparasiticide for at least 60 days, had demonstrated good tick retention prior to treatment, and were in good health at enrollment. Sixteen different purpose-bred Beagles, including both sexes, were enrolled in each study. Dogs ranged in age from 12 months to 7 years, and weighed between 8.3 kg and 32.0 kg. Female dogs were confirmed not to be pregnant or lactating. Each dog was individually identified by a unique and permanent code (microchip or tattoo). Dogs were housed in individual indoor pens such that no physical contact was possible between them, and the possibility of tick transfer among animals was minimal. Dogs were fed an appropriate maintenance ration of a commercial canine diet for the duration of the study. Water was available *ad libitum*.

### 2.2. Study methods

Day 0 for each study was the day dogs were administered the study treatment. Dogs were acclimated to the study conditions for at least 7 days prior to treatment. For tick infestations, a pre-counted aliquot of approximately 50 adult unfed ticks were placed onto the hair coat and allowed to disperse on the dog. Ticks were applied in an approximate 1:1 sex ratio, with the exception of one *I. ricinus* study in which a 3:2 female to male ratio was used as this was indicated per previous use for that tick strain.

Tick counts were performed by personnel trained in the standard procedures in use at the test facility. Personnel changed protective clothing between dogs to avoid any cross-contamination, and personnel conducting parasite or other observations were unaware of treatment assignments. Initially, the entire dog's entire body was examined, pushing the hair against its natural nap, exposing, counting and removing the ticks. After the manual inspection, an extra-fine tooth comb was used to comb the animal to remove any missed ticks. Each dog was examined for at least 10 min. If ticks were encountered in the last minute, combing was continued in one minute increments until no ticks were encountered. The ticks were examined to assess viability (movement and reaction to CO<sub>2</sub> stimulation) and the numbers of live ticks was quantified.

At least 16 animals arrived into the housing facilities on or before Day –7. General health observations were performed at least once a day from the start of the acclimation period. All dogs were given a physical examination to evaluate general health and suitability for inclusion into the study. The dogs were examined to ensure they were free of ticks and were then infested to determine the host suitability between Day –9 and Day –7. The live attached ticks present on each dog were counted and removed at 48 (±2) hours after infestation. The 16 dogs with the highest counts were selected for inclusion, ranked by decreasing tick count into blocks of two and randomly allocated within block to treatment with placebo or sarolaner tablets. Blocks of dogs were randomly assigned to adjacent pens within the test facility. Dogs were moved into their allocated pens on or before Day –2.

Dogs were weighed and infested with ticks on Day –2. On Day 0, the dogs were dosed orally with placebo (Simparica™ formulation without active ingredient sarolaner) or sarolaner strengths of 5, 10, 20, or 40 mg such that the sarolaner dose was as close as possible to 2 mg/kg without under-dosing.

Each dog was offered its regular food ration ~20 min before dosing. Dogs were hand-pilled to ensure accurate dose delivery. Each dog was observed for a minimum of 1 min after dosing for evidence that the dose was swallowed, and for potential adverse events associated with treatment and then for up to 2 h for any signs of emesis. Dogs were observed for general health and any reaction to treatment approximately 1, 3 and 6 h after treatment. On Day 2, each dog was examined to remove and count ticks. In all studies except two of the three conducted against *I. hexagonus*, animals were subsequently re-infested with ticks on Days 5, 12, 19, 26 and 33. In one *I. hexagonus* study dogs were re-infested only on Days 19 and 26, and in one study only on Day 33. Dogs were examined, combed and parasites counted 48 (±2) hours after each infestation.

Ticks were sourced from multiple laboratory maintained colonies with isolates all originating from Europe. These ticks were originally isolated from the field, and new ticks had been introduced into each colony within the previous ten years.

### 2.3. Data analysis

The individual dog was the experimental unit and the primary endpoint was live tick counts. Tick counts were transformed by the  $\log_e(\text{count} + 1)$  transformation prior to analysis in order to stabilize the variance and normalize the data. Using the PROC MIXED procedure (SAS 8.2, Cary NC), transformed counts were analyzed using a mixed linear model for repeated measures. The model included the fixed effect of treatment, day of study and the interaction between treatment and day of study. The random effects included room, block within room, the interaction between block and treatment within room (animal term) and error. Testing was two-sided at the significance level  $\alpha = 0.05$ . Percent efficacy was calculated using Abbott's formula:

$$\% \text{reduction} = 100 \times \frac{\text{meancount}(\text{placebo}) - \text{meancount}(\text{treated})}{\text{meancount}(\text{placebo})}$$

## 3. Results

### 3.1. Efficacy

Dogs in the placebo-treated group maintained tick infestations throughout the studies (Tables 1–4).

For *D. reticulatus*, efficacy against existing infestations was 100% at 48 h after treatment in both studies. Against subsequent weekly re-infestations, efficacy 48 h after infestation was  $\geq 99.0\%$  in one study and 100% in the second study through 35 days post treatment (Table 1). Efficacy against existing infestations of *I. hexagonus*

Download English Version:

<https://daneshyari.com/en/article/2469837>

Download Persian Version:

<https://daneshyari.com/article/2469837>

[Daneshyari.com](https://daneshyari.com)