

Short communication

First isolation of *Trichinella britovi* from a wild boar (*Sus scrofa*) in Belgium

F. Schynts^a, J. van der Giessen^b, S. Tixhon^c, E. Pozio^d,
P. Dorny^e, J. de Borchgrave^{e,*}

^a Centre d'Economie Rurale (CER), Division de Virologie Animale, Rue du Carmel 1, 6900 Marloie, Belgium

^b Microbiological Laboratory for Health Protection, National Institute of Public Health and the Environment (RIVM),
Antonie van Leeuwenhoeklaan 9, 3720 BA Bilthoven, The Netherlands

^c Agence Fédérale pour la Sécurité de la Chaîne Alimentaire (AFSCA),
Unité Provinciale de Contrôle de la Province de Luxembourg,
Rue du Vicinal 1, 6800 Libramont, Belgium

^d Department of Infectious, Parasitic and Immunomediated Diseases,
Istituto Superiore di Sanita (ISS), Viale Regina Elena 299, 00161 Rome, Italy

^e Institute of Tropical Medicine Antwerp (ITMA), National Trichinella Reference Centre, Nationalestraat 155,
2000 Antwerp, Belgium

Received 24 May 2005; received in revised form 9 August 2005; accepted 1 September 2005

Abstract

Since 1992, when the European Union Council Directive requires that wild boars (*Sus scrofa*) hunted in EU for commercial purpose should be examined for *Trichinella*, the infection has not been detected in wild boars from Belgium, despite serological evidence of the presence of anti-*Trichinella* antibodies in wildlife and previous reports of *Trichinella* larvae in this host species. In November 2004, *Trichinella* larvae were detected in a wild boar hunted near Mettet, Namur province (Southern Belgium). Larvae were identified as *Trichinella britovi* by polymerase chain reaction methods. This is the first report of the identification of *Trichinella* larvae from Belgium at the species level. The detection of *T. britovi* in wildlife in Belgium is consistent with findings of this parasite in other European countries and confirms the need to test game meat for *Trichinella* to prevent its transmission to humans.

© 2005 Elsevier B.V. All rights reserved.

Keywords: *Trichinella britovi*; Wild boar; Belgium; Wildlife; Trichinellosis; Epidemiology

1. Introduction

Four species of *Trichinella* (*Trichinella spiralis*, *Trichinella nativa*, *Trichinella britovi* and *Trichinella pseudospiralis*) are represented in the European

* Corresponding author. Tel.: +32 3 2476271; fax: +32 3 2476268.
E-mail address: jdeburchgrave@itg.be (J. de Borchgrave).

Union (EU) (Pozio, 2001). In these countries, human infections are related to the consumption of meat from game, domestic pigs raised in organic farms in endemic areas or fed with offal from game, and horses imported from countries of Eastern Europe and America (Pozio, 1998; Pozio, 2001; Boireau et al., 2000). The European Union (Directive 92/45/EEC, 1992) obliges the examination of meat of wild boar, pig and horse for the presence of *Trichinella* spp.

In Belgium, *Trichinella* infection has not been detected in domestic pigs or horses and only one outbreak was documented in humans following the consumption of pork from a wild boar (Famerée et al., 1979). In wildlife Famerée et al. (1981) detected *Trichinella* larvae in 6.7% of wild boars (*Sus scrofa*), 2.2% of muskrats (*Ondatra zibethica*), 6.5% of brown rats (*Rattus norvegicus*) and 11.1% of black rats (*Rattus rattus*). However, at that time the parasites were not confirmed, and not identified at the species level. Since 1992, annually 8000 sport hunted wild boars are tested for *Trichinella* in Belgium, as imposed by Directive 92/45/EEC, and infection had until now not been detected. This paper presents the isolation of *Trichinella* larvae by artificial digestion and the subsequent confirmation and characterization of the isolate, from a wild boar in southern Belgium.

2. Materials and methods

Routine inspection by artificial digestion of pooled samples of 5 g of tongue and diaphragm muscle from 20 animals (EU Directive 92/45/EEC) was carried out in the laboratory of the Centre d'Economie Rurale of Marloie in Belgium. To trace back the infected animal, muscle samples of tongue, diaphragm and forearm from the 20 wild boars were digested separately. In order to identify the parasite at the species level, larvae were sent to the *Trichinella* Reference Laboratory of The Netherlands (RIVM, Bilthoven) and tested by a 5S rDNA based PCR followed by DNA sequencing (Rombout et al., 2001; Van der Giessen et al., 2005). For case registration, the larvae were also subjected to a multiplex PCR analysis (Pozio and La Rosa, 2003) in the International *Trichinella* Reference Centre in Italy (ISS, Rome).

3. Results

Five larvae were recovered after pooled sample digestion. Individual digestions revealed that the infected muscles originated from a wild boar shot near Mettet (50.19N, 4.40E). The average parasite load in mixed muscles from the tongue and diaphragm was 0.7 larva/g (LPG). No larvae were detected in 55 g of forearm muscles. The shape and movement of larvae were suggestive for *Trichinella*. The examination of larvae at higher magnification showed the presence of the stichosome and of a row of collateral dots, which are morphological characters of the *Trichinella* genus. In the reference laboratories, larvae were identified as belonging to *T. britovi*. Phylogenetic analysis of the 5S rDNA sequences (RIVM, The Netherlands) showed 99.5% similarity with *T. britovi* AY009943.1 from Genbank.

4. Discussion

Sylvatic carnivores (e.g. red fox, wolf and mustelids) represent the main hosts of *T. britovi*. The infection can be transmitted to wild boars and consequently it can easily reach the human being (Pozio, 1998). In the last decades, the wild boar populations of Europe have increased exponentially favoring *Trichinella* transmission and consequently increasing the biomass of this parasite (Hars et al., 2000). Even if the experimental infection of wild boars shows that swine is not the optimal host for *T. britovi* (Kapel, 2001), epidemiological data including the present work stress the role played by this animal species for spreading the infection in Europe. *Trichinella* infection can be maintained by a sylvatic cycle for decades as has been shown in Ireland, where *T. spiralis* was maintained among the fox population for >30 years, without any documented infection in domestic animals and humans (Rafter et al., 2005).

The only documented case of trichinellosis in Belgium was caused by the consumption of wild boar meat, originating from two home-fed animals in the northern part of the country, where wild boars are not present in natural conditions (Famerée et al., 1979).

In Belgium studies on wildlife species not intended for human consumption suggest low prevalence of *Trichinella* spp. In the season 2003–2004, 199 red

Download English Version:

<https://daneshyari.com/en/article/2472702>

Download Persian Version:

<https://daneshyari.com/article/2472702>

[Daneshyari.com](https://daneshyari.com)