



A novel spotted fever group *Rickettsia* infecting *Amblyomma parvitarsum* (Acari: Ixodidae) in highlands of Argentina and Chile

Maria Ogrzewalska^{a,b,*}, Fernanda A. Nieri-Bastos^a, Arlei Marcili^{c,d}, Santiago Nava^e, Daniel González-Acuña^f, Sebastián Muñoz-Leal^{a,f}, Ignacio Ruiz-Arondo^g, José M. Venzal^h, Atilio Mangold^e, Marcelo B. Labruna^a

^a Departamento de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, São Paulo, SP, Brazil

^b Laboratório de Hantavírus e Rickettsioses, Instituto Oswaldo Cruz, FIOCRUZ, Rio de Janeiro, RJ, Brazil

^c Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, São Bernardo do Campo, SP, SP, Brazil

^d Medicina Veterinária e Bem estar animal, Universidade de Santo Amaro, SP, Brazil

^e Instituto Nacional de Tecnología Agropecuaria, Rafaela, Santa Fe, Argentina

^f Facultad de Ciencias Veterinarias, Universidad de Concepción, Chillán, Chile

^g Departamento de Patología Animal, Facultad de Veterinaria, Universidad de Zaragoza, Spain

^h Laboratorio de Vectores y enfermedades transmitidas, Universidad de la República, CENUR Litoral Norte, Salto, Uruguay

ARTICLE INFO

Article history:

Received 29 April 2015

Received in revised form

21 December 2015

Accepted 7 January 2016

Available online 8 January 2016

Keywords:

Spotted fever group

Rickettsia

Amblyomma parvitarsum

Argentina

Chile

ABSTRACT

The tick *Amblyomma parvitarsum* (Acari: Ixodidae) has established populations in Andean and Patagonian environments of South America. For the present study, adults of *A. parvitarsum* were collected in highland areas (elevation >3500 m) of Argentina and Chile during 2009–2013, and tested by PCR for rickettsial infection in the laboratory, and isolation of rickettsiae in Vero cell culture by the shell vial technique. Overall, 51 (62.2%) out of 82 *A. parvitarsum* adult ticks were infected by spotted fever group (SFG) rickettsiae, which generated DNA sequences 100% identical to each other, and when submitted to BLAST analysis, they were 99.3% identical to corresponding sequence of the *ompA* gene of *Rickettsia* sp. strain Atlantic rainforest. Rickettsiae were successfully isolated in Vero cell culture from two ticks, one from Argentina and one from Chile. DNA extracted from the third passage of the isolates of Argentina and Chile were processed by PCR, resulting in partial sequences for three rickettsial genes (*gltA*, *ompB*, *ompA*). These sequences were concatenated and aligned with rickettsial corresponding sequences available in GenBank. Phylogenetic analysis revealed that the *A. parvitarsum* rickettsial agent grouped under high bootstrap support in a clade composed by the SFG pathogens *R. sibirica*, *R. africae*, *R. parkeri*, *Rickettsia* sp. strain Atlantic rainforest, and two unnamed SFG agents of unknown pathogenicity, *Rickettsia* sp. strain NOD, and *Rickettsia* sp. strain ApPR. The pathogenic role of this *A. parvitarsum* rickettsia cannot be discarded, since several species of tick-borne rickettsiae that were considered nonpathogenic for decades are now associated with human infections.

© 2016 Published by Elsevier GmbH.

1. Introduction

Bacteria of the genus *Rickettsia* (Rickettsiales: Rickettsiaceae) are obligate intracellular organisms that multiply freely in the cytosol of eukaryotic cells of a variety of invertebrate hosts,

including insects and ticks. Many *Rickettsia* species undergo transovarial transmission in their invertebrate hosts, which can sustain rickettsial infection for a number of successive generations. Some *Rickettsia* species infect hematophagous arthropods (e.g., fleas, ticks), which can act as vectors of pathogenic rickettsiae to vertebrate hosts, including humans (Parola et al., 2013). The pathogenic *Rickettsia* species have been divided into two main groups, based on genotypic and phenotypic criteria: the typhus group (TG), typically transmitted by fleas and lice; and the spotted fever group (SFG), mostly transmitted by ticks (Parola et al., 2013). A recent review listed 11 *Rickettsia* species occurring in South America, mostly SFG agents associated with ticks (Labruna et al., 2011). Additionally, a

* Corresponding author at: Laboratório de Hantavírus e Rickettsioses, Instituto Oswaldo Cruz, FIOCRUZ, Av. Brasil 4365, Manguinhos, Rio de Janeiro, RJ 21045-900, Brazil.

E-mail address: maria.ogrzewalska@fiocruz.ioc.br (M. Ogrzewalska).

number of SFG unnamed agents have been reported infecting South American ticks (Parola et al., 2013).

The tick *Amblyomma parvitarsum* (Acari: Ixodidae) has established populations in Andean and Patagonic environments of Argentina, Bolivia, Chile, and Peru, where the adult tick stage feeds chiefly on native camelids, while immature ticks seem to feed primarily on lizards (Muñoz-Leal et al., 2014). Until the present study, no organism had been reported infecting *A. parvitarsum* ticks. Herein, we report a novel rickettsial agent infecting *A. parvitarsum* ticks in Argentina and Chile.

2. Materials and methods

Adults of *A. parvitarsum* were collected in two localities of Argentina during 2009 and 2013, and one locality of Chile during 2012 (Table 1). Ticks were determined following Estrada-Peña et al. (2005). While the 2009 ticks were preserved in absolute ethanol before being sent to the laboratory, the 2012 and 2013 ticks were sent alive to the laboratory, where they were frozen at -80°C upon arrival.

All ticks (except for eight ticks from the 2012 and 2013 collections, as stated below) were individually subjected to DNA extraction by the guanidine isothiocyanate phenol technique (Sangioni et al., 2005) and tested for *Rickettsia* by two different PCR protocols. Firstly, all DNA samples were tested with primers CS-78 and CS-323 targeting a relatively conserved fragment of the citrate synthase gene (*gltA*) that occurs in all *Rickettsia* species (Labruna et al., 2004). Samples yielding a visible amplicon of the expected size (compatible with 398-bp) were then tested by a second PCR protocol with primers Rr190.70F and Rr190.602R, which amplify a 532 bp fragment of the 190-kDa outer membrane protein gene (*ompA*) from only some *Rickettsia* species belonging to the SFG (Regnery et al., 1991).

Attempts to isolate rickettsiae in cell culture were performed with eight ticks (four from the 2012 collection and four from the 2013 collection). For this purpose, each tick was thawed and immediately subjected to the shell vial technique for isolation of rickettsiae in Vero cell culture, as previously described (Labruna et al., 2004). Briefly, cultures of Vero cells were inoculated with tick-body homogenates and incubated at 28°C . The percentage of Vero cells infected with rickettsiae was monitored by the use of Giménez staining of cells scraped from each inoculated monolayer. After the establishment of each isolate in the laboratory (i.e., at least 3 cell passages with >90% infected cells), rickettsial DNA was extracted from the infected cells (Labruna et al., 2004). The extracted DNA was tested in a battery of different PCR protocols, using primer pairs targeting a 632-bp fragment of the *ompA* gene (Roux et al., 1996), a

856-bp fragment of the rickettsial 135-kDa outer membrane protein gene (*ompB*) (Roux and Raoult, 2000), and two overlapping fragments (398-bp and 834-bp) of the *gltA* gene (Labruna et al., 2004). From each of the 8 ticks processed by shell vial, part of the tick homogenate was also tested by PCR targeting fragments of the *gltA* and *ompA* genes, as stated above for ticks. PCR products were purified and sequenced in an automatic sequencer (model ABI 3500 Genetic Analyzer; Applied Biosystems/Thermo Fisher Scientific, Foster City, CA) according to the manufacturer's protocol. The generated sequences were submitted to BLAST analyses (www.ncbi.nlm.nih.gov/blast) to infer the closest similarities available in GenBank.

Partial DNA sequences obtained from the amplified PCR products (*gltA*, *ompA*, and *ompB*) of the *A. parvitarsum* rickettsial isolate were concatenated and aligned with corresponding sequences of different SFG *Rickettsia* species and yet unnamed rickettsial agents available in GenBank using Clustal X (Thompson et al., 1997) and adjusted manually using GeneDoc (Nicholas et al., 1997). Phylogenetic analyses were performed using PAUP version 4.0b10 (Swofford, 2002) to maximum parsimony (MP); the confidence values for individual branches of the resulting tree were determined by bootstrap analysis with 1000 replicates. Bayesian analysis (BA) was performed with MrBayes version 3.1.2 (Huelsenbeck and Ronquist, 2001) software with 1,000,000 generations using the GTR+I+G substitution model. Corresponding sequences of *Rickettsia australis* were used as outgroup.

3. Results

Overall, 51 (62.2%) out of 82 *A. parvitarsum* adult ticks were infected by rickettsiae, including ticks from Argentina and Chile (Table 1). All infected ticks yielded PCR products by both *gltA*- and *ompA*-PCR assays. The overall infection rates per tick gender were 62.1% (18/29) for male ticks and 61.9% (26/42) for female ticks from Argentina (grouping 2009 and 2013 data). In Chile, where no male ticks were collected, 63.6% (7/11) female ticks were infected by rickettsiae. DNA sequences generated from the *ompA*-PCR products were all 100% identical to each other, and when submitted to BLAST analysis, they were 99.3% (444/447-bp) identical to the corresponding sequence of *Rickettsia* sp. strain Atlantic rainforest (JQ906784), 98.9% (471/476-bp) to *Rickettsia africae* (CP001612), 98.7% (469/475-bp) to *Rickettsia endosymbiont of Amblyomma tuberculatum* (JF934878), and 98.5% (469/476-bp) to *Rickettsia sibirica* (CP001612).

Rickettsiae were successfully isolated in Vero cell culture from two ticks, one from Argentina and one from Chile. These two isolates were successfully established in the laboratory, with

Table 1
Rickettsial infection in *Amblyomma parvitarsum* adult ticks from Argentina and Chile.

Locality						Ticks		
Name	Province or Region	Country	Coordinates	Elevation	Date	Source	No. collected	No. infected by <i>Rickettsia</i> (% infection)
Parque Nacional San Guillermo	San Juan	Argentina	29°28' S, 69°19' W	3600	February 2009	Environment ^a	45	29 (64.4)
45 km southwest of provincial route 129 from San Antonio de los Cobres	Salta	Argentina	24°22' S, 66°42' W	4500	February 2013	Environment ^a	26	15 (57.7)
Visviri	Arica and Parinacota	Chile	17°35' S, 69°28' W	4069	August 2012	Vicuñas ^b	11	7 (63.6)
Total							82	51 (62.2)

^a Ticks collected through visual inspection of the soil within vicuñas' resting places.

^b *Vicugna vicugna* (Artiodactyla: Camelidae).

Download English Version:

<https://daneshyari.com/en/article/2473900>

Download Persian Version:

<https://daneshyari.com/article/2473900>

[Daneshyari.com](https://daneshyari.com)