

Multiple Linear Regression Analysis Indicates Association of P-Glycoprotein Substrate or Inhibitor Character with Bitterness Intensity, Measured with a Sensor

KENTARO YANO,¹ SUZUNE MITA,¹ KAORI MORIMOTO,² TAMAMI HARAGUCHI,³ HIROSHI ARAKAWA,¹ MIYAKO YOSHIDA,³ FUMIYOSHI YAMASHITA,⁴ TAKAHIRO UCHIDA,³ TAKUO OGIHARA¹

¹Faculty of Pharmacy, Takasaki University of Health and Welfare, Takasaki, Gunma 370-0033, Japan

²Department of Drug Absorption and Pharmacokinetics, Tohoku Pharmaceutical University, Sendai, Miyagi 981-8558, Japan

³Department of Clinical Pharmaceutics, Faculty of Pharmaceutical Sciences, Mukogawa Women's University, Nishinomiya, Hyogo 663-8179, Japan

⁴Department of Drug Delivery Research, Graduate School of Pharmaceutical Sciences, Kyoto University, Sakyo, Kyoto 606-8501, Japan

Received 17 July 2014; revised 2 October 2014; accepted 7 October 2014

Published online 24 December 2014 in Wiley Online Library (wileyonlinelibrary.com). DOI 10.1002/jps.24232

ABSTRACT: P-glycoprotein (P-gp) regulates absorption of many drugs in the gastrointestinal tract and their accumulation in tumor tissues, but the basis of substrate recognition by P-gp remains unclear. Bitter-tasting phenylthiocarbamide, which stimulates taste receptor 2 member 38 (T2R38), increases P-gp activity and is a substrate of P-gp. This led us to hypothesize that bitterness intensity might be a predictor of P-gp-inhibitor/substrate status. Here, we measured the bitterness intensity of a panel of P-gp substrates and nonsubstrates with various taste sensors, and used multiple linear regression analysis to examine the relationship between P-gp-inhibitor/substrate status and various physical properties, including intensity of bitter taste measured with the taste sensor. We calculated the first principal component analysis score (PC1) as the representative value of bitterness, as all taste sensor's outputs shared significant correlation. The P-gp substrates showed remarkably greater mean bitterness intensity than non-P-gp substrates. We found that K_m value of P-gp substrates were correlated with molecular weight, log P , and PC1 value, and the coefficient of determination (R^2) of the linear regression equation was 0.63. This relationship might be useful as an aid to predict P-gp substrate status at an early stage of drug discovery. © 2014 Wiley Periodicals, Inc. and the American Pharmacists Association *J Pharm Sci* 104:2789–2794, 2015

Keywords: P-glycoprotein; substrate specificity; inhibition; bitterness intensity; Log P ; Transporters; absorption; solubility; Multivariate analysis; Physicochemical properties; multiple linear regression analysis; correlation efficient

INTRODUCTION

P-glycoprotein (P-gp) is an efflux transporter in small intestine,¹ liver,² kidney,³ and brain,⁴ serving to transport xenobiotics and/or toxic compounds out of cells, including tumor cells.^{5,6} Therefore, this transporter is involved in poor gastrointestinal absorption of various drugs and multidrug resistance in tumor cells. P-gp substrates tend to have higher lipophilicity and/or larger molecular weight than nonsubstrates, but the basis of the substrate specificity of P-gp remains unclear. Nevertheless, it is very important to determine whether a compound may be a substrate of P-gp at an early stage of drug discovery, in order to predict its gastrointestinal absorption and distribution to tumor tissues.

Most compounds that are poisonous to humans have a bitter taste.⁷ Bitterness is sensed by bitter taste-sensing type 2 receptors (T2Rs), which are expressed in taste buds on the tongue, palate epithelium, and mucosa of the gastrointestinal tract.^{8–10} Among them, T2R member 38 (T2R38) is expressed in the

gastrointestinal tract and this receptor is stimulated by phenylthiocarbamide (PTC), which is a bitter substance. Interestingly, it was reported that P-gp mRNA and transport activity were upregulated by PTC in Caco-2 cells and rat intestine. Further, this compound is a P-gp substrate.¹¹

This led us to hypothesize that P-gp-inhibitor/substrate character might be associated with bitter taste. To investigate this hypothesis, we measured the bitterness intensity of P-gp substrates and non-P-gp substrates using six taste sensors, which yields values that are well correlated with those obtained in human gustatory sensation tests.^{12,13} We then used multiple regression analysis to examine the contributions of various physical properties, including the first principal component analysis score (PC1) as value indicating a representative character of bitterness intensity obtained by using taste sensors, to P-gp inhibition and substrate recognition. On the basis of the results, we propose an equation to predict whether compounds are P-gp substrates or non-P-gp substrates.

METHODS

Materials

Hanks' balanced salts, rhodamine123 (Rho123), estrone 3-sulfate sodium salt, and theophylline were purchased from Sigma-Aldrich Inc. (St. Louis, Missouri). Oseltamivir was purchased from Sequoia Research Products (Pangbourne,

Abbreviations used: P-gp, P-glycoprotein; T2R, taste-sensing type 2 receptor; C/M ratio, ratio of intracellular concentration to medium concentration; PCA, principal component analysis; PC1, the first PCA score; M.W., molecular weight; PTC, phenylthiocarbamide; Rho123, rhodamine123; CPA, change of membrane potential by adsorption.

Correspondence to: Takuo Ogihara (Telephone: +81-27-352-1180; Fax: +81-27-352-1118; E-mail: togihara@takasaki-u.ac.jp)

Journal of Pharmaceutical Sciences, Vol. 104, 2789–2794 (2015)

© 2014 Wiley Periodicals, Inc. and the American Pharmacists Association

UK). Colchicine, cyclosporin A, etoposide, hydrocortisone, quinidine, verapamil hydrochloride, vinblastine sulfate, 5-FU, antipyrine, caffeine, methotrexate, sodium taurocholate, pravastatin sodium salt, and quinine hydrochloride were purchased from Wako Pure Chemical Industries, Ltd. (Osaka, Japan). Fexofenadine hydrochloride and uric acid were purchased from Tokyo Chemical Industry Company, Ltd. (Tokyo, Japan) and ChromaDex, Inc. (Irvine, California), respectively. Other reagents were of analytical grade commercial products. P-gp-overexpressing porcine kidney epithelial cell line LLC-GA5-COL150 (MDR1 gene-transfected cells) was obtained from RIKEN Cell Bank. Medium 199 (M199), penicillin, streptomycin, and trypsin ethylenediaminetetraacetic acid were purchased from Invitrogen (Carlsbad, California).

Measurement of Bitterness Intensity Using Taste Sensors

The bitterness intensity of P-gp substrates (10 samples) and non-P-gp substrates (nine samples) was measured with several taste sensors (SA402B, Intelligent Sensor Technology), using quinine as a standard, as described previously.¹⁴ The non-P-gp substrates included passive diffusers (antipyrine¹⁵ and theophylline¹⁶) and substrates of multidrug-resistance-associated protein (estrone 3-sulfate¹⁷), breast cancer-resistance protein (methotrexate, pravastatin,¹⁸ and uric acid¹⁹), bile salt export pump (taurocholate²⁰), and an unknown transporter (not P-gp) (caffeine²¹). The detection part of the taste-sensing system consists of AN0, AC0, and BT0 sensors, which act as working electrodes and are composed of lipid/polymer membranes, and a reference electrode. Drugs with bitterness are adsorbed on the hydrophobic part of the membrane and change the membrane potential by altering the charge density.²² Each sensor's electrodes were dipped into the reference solution (V_r) to determine the baseline, and then into the sample solution (V_s). The relative sensor output was represented as the potential difference ($V_s - V_r$) between the sample and the reference solution. When the electrode was immersed into the reference solution a second time, the new potential of the reference solution was defined as V_r' . The difference between the potential of the reference solution before and after sample measurement ($V_r' - V_r$) was defined as CPA (change of membrane potential by adsorption), and is a measure of the aftertaste of bitterness.

Principal Component Analysis of Bitterness Intensity

Because six measures of bitterness intensity obtained from the taste sensor measurement were highly intercorrelated, principal component analysis (PCA) was carried out to condense information from the measurements. Data were standardized to a mean of zero with a standard deviation of unity, and eigenvalues and eigenvectors were then calculated for the correlation matrix. Taking the eigenvectors as loadings, the standardized data matrix was orthogonally decomposed to calculate the corresponding PCA scores. The first PCA score (PC1) was that that gave the largest eigenvalue, followed by PC2, PC3, PC4, PC5, and PC6. A series of calculations were made with the function "prcomp(PCA scores)" from the freely available statistical package R (<http://cran.r-project.org/>).

Cell Culture

LLC-GA5-COL150 cells were cultured in M199 containing 10% fetal bovine serum, 100 U/mL penicillin, 100 µg/mL

streptomycin, and 150 ng/mL colchicine at 37°C in an atmosphere of 5% CO₂ at 95% relative humidity. LLC-GA5-COL150 cells were seeded on 12-well collagen-coated plates (Becton Dickinson Bioscience, Bedford, Massachusetts) at a cell density of 3.0×10^5 cells/mL. Cells were grown for 7 days, and then used to evaluate Rho123 accumulation. The medium was changed to M199 without colchicine the day prior to the accumulation study.

Accumulation Study

Uptake studies with LLC-GA5-COL150 cells were performed at 37°C in Hank's balanced salt solution (HBSS) with 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (pH 7.4) buffer containing 10 µM Rho123 as a substrate of P-gp. Cells were preincubated with or without 100 µM of the 18 test drugs for 15 min. After preincubation, Rho123 uptake experiments were performed in HBSS-HEPES buffer containing 10 µM Rho123 in the presence of 100 µM test drugs for 10 min. Intracellular concentrations of Rho123 were measured using a WALLAC Multilabel/Luminescence Counter (PerkinElmer, Waltham, Massachusetts) at wavelengths of 485 nm (excitation) and 538 nm (emission). The ratio of intracellular concentration to medium concentration (C/M ratio) represented the coefficient of intracellular and extracellular ratio of Rho123 and was determined using the following formula: C/M ratio = intracellular concentration (µM)/extracellular concentration (10 µM)/protein concentration (mg/mL). The accumulation rates of Rho123 by cells treated with test drugs were compared with the control (Rho123 alone).

Statistical Analysis

All data are presented as mean \pm standard error of the mean (SEM). Correlation analysis of six kinds of bitter taste parameter was conducted first. On the basis of the results, PC1 was selected as a representative value of bitterness, and the two-tailed *t*-test was used to compare bitterness intensity between P-gp substrates and non-P-gp substrates. For $p < 0.05$ or $p < 0.01$, differences between means were considered significant. Multiple linear regression analysis for the inhibition rates of P-gp or K_m values (Michaelis constants) of P-gp for its substrates was conducted against various physical properties, including bitterness intensity. Values of physical properties and K_m values of P-gp for its substrates were taken from Sci Finder, ChemDraw, or previous reports.^{23–28} The K_m values of non-P-gp substrates were standardized at 10,000 µM, which indicated that the concentrations of drugs were clearly nonsubstrate.²⁹ The relationship between dependent variable (y) and explanatory variable (x) was represented by the following equation:

$$y_i = \alpha_0 + \alpha_1 \cdot x_1 + \alpha_2 \cdot x_2 + \dots$$

R^2 value is the coefficient of determination.

RESULTS

Measurement of Bitterness Intensity of P-gp and Non-P-gp Substrates Using Taste Sensors

To examine the idea that P-gp-inhibitor/substrate character might be associated with bitter taste, we measured the values of bitterness intensity for Rho123 (a good P-gp substrate), verapamil hydrochloride (a weak substrate), and antipyrine

Download English Version:

<https://daneshyari.com/en/article/2484381>

Download Persian Version:

<https://daneshyari.com/article/2484381>

[Daneshyari.com](https://daneshyari.com)