

# Influence of salt stress on phosphorus metabolism in the roots and leaves of one month old *Prosopis juliflora* (Sw.) DC seedlings

\*Patil, Amol V. and Chavan, P. D.

Department of Botany, Shivaji University, Kollhapur-416 004 (MS) INDIA.

## ABSTRACT

A sand culture experiment was designed to study the effect of sodium chloride salinity on phosphorus metabolism in the roots and leaves of one month old *Prosopis juliflora* (Sw.) DC seedlings. It was found that the P level in the roots as well as leaves was decreased with increasing level of salinity in rooting medium. However, the activities of enzymes acid phosphatase and ATPase were increased in both the parts of seedlings grown in saline conditions. The activities of alkaline phosphatase and inorganic pyrophosphatase were found to be decreased in the root and leaves of seedlings grown under saline conditions.

**Key words:** salinity, phosphorus, enzymes, *Prosopis juliflora*.

## INTRODUCTION

The selection and breeding of salt tolerant crops is regarded as one of the main approaches to deal with a serious problem of salt affected soils throughout the world. In order to achieve this strategy it is necessary to identify the mechanisms of salt tolerance in the plant species well adapted to such problem soils. *Prosopis juliflora* is one such plant species which can successfully grow and complete its life cycle in a variety of problem soils. It is noticed that the plant has successfully established in farmlands of Digraj (Dist. Sangli) which are heavily affected by secondary salinization. *Prosopis juliflora* is a multipurpose plant of great economic potential. The ability of this species to grow on the poorest soil, under arid conditions and on saline soil is well known Pasiecznik et al.<sup>[1]</sup>

According to Dagar and Tomar<sup>[2]</sup> in India about 8.53 million ha land is waterlogged, 5.50 million ha land is saline and 3.88 million ha land is alkaline and more and more land is becoming water logged due to several factors. According to CSSRI these soils can be judiciously utilized for raising forestry, agriculture and horticulture crops. Afforestation programme for saline soil requires the proper selection of

tree species, as the major problems of such soils are high water table, high salinity impeded drainage and less soil aeration for tree growth, Singh<sup>[3]</sup>

Phosphorus metabolism occupies a key position in cellular-biochemistry as it is related with energy relation in respiration and photosynthesis. Hence, an attempt has been made to study the phosphorus metabolism in the roots and leaves of *Prosopis juliflora* seedlings grown under salinity stress in laboratory conditions.

## MATERIALS AND METHODS

For the experiment, seeds were obtained from the pod of *Prosopis juliflora* plants growing in the salt affected agriculture field in Sangli district in the month of April-May. Mechanically scarified seeds were used to raise the seedlings. After the establishment of seedlings for 5 days, they were treated with increasing concentration of salt (100, 200, and 300 mM NaCl) mix with half strength Hoagland solution. The seedling were grown for one month and then analysed for phosphorus metabolism. The method of Sekine *et al.*<sup>[4]</sup> was employed for estimation of Phosphorus from the root and leaves. Fresh leaves and roots were used for the assay of enzymes of Phosphorus metabolism. For enzyme acid phosphatase crude enzyme was prepared in 0.1 M acetate buffer (pH 5) and assayed according to the method of Mclachlan<sup>[5]</sup> The activity of enzyme ATPase was determined following the method

\*Address for correspondence:  
E-mail: prosopis.1912@gmail.com


DOI: 10.5530/pj.2011.25.9

described by Todd and Yoo<sup>[6]</sup> and liberated phosphorus was estimated by the standard method.<sup>[7]</sup> The method described by Weimberg<sup>[8]</sup> was employed for the study of activity of enzyme alkaline phosphatase. A method by Kar and Mishra<sup>[9]</sup> was employed for the determination of the activity of enzyme alkaline inorganic pyrophosphatase. The soluble proteins in the enzyme preparations were determined according to the method of Lowry *et al.*<sup>[10]</sup>

## RESULTS

Phosphorus is an important macronutrient essential for all living organisms. It plays a major role in energy transfer during plant metabolism like respiration, photosynthesis in the form of ATP, NADP and also in cell division and cell expansion. Phosphorus is involved in the formation of cell membrane lipids, which play a vital role in ionic regulation.<sup>[11]</sup> There are many reports indicating suppression of P uptake due to salt stress.<sup>[12,13]</sup> Nieman and Clark<sup>[14]</sup> also found depression of total P in the corn leaves due to salinity at low level of inorganic phosphorus in the nutrient solution. In case of *Prosopis cineraria* seedlings Ramoliya *et al.*<sup>[15]</sup> noticed that phosphorus content was significantly decreased in the leaves with increase in soil salinity while that was gradually decreased in the stem and root tissues. A decrease in P content of root tissue and that increase in the leaf tissue of salt grown *Poncirus trifoliata* was evident in the experiments by Tozly *et al.*<sup>[16]</sup>

*Prosopis juliflora* seedlings have shown a pattern similar to that in *Prosopis cineraria* since in both root and leaves a decline in P content was evident in the seedlings exposed to salt stress (Figure 1) According to Gibson<sup>[17]</sup> phosphorus deficiency induced by salinity could reduce the cellular ability to accumulate optimum concentration of ion without reduced growth. Thus in contrast to Calcium and Potassium nutrition



**Figure 1:** Effect of Sodium chloride salinity on phosphorus content in the roots and leaves of *Prosopis juliflora* (Sw.).

which appears to be quite stable during salt stress in this species, the phosphorus nutrition in *Prosopis juliflora* seems to be sensitive to salt stress. The disturbance in P nutrition can have significant effects on overall plant metabolism in view of a key role of this element in cellular biochemistry.

Effect of sodium chloride salinity on the activity of enzyme acid phosphatase in the leaves and roots of *Prosopis juliflora* is recorded in figure 2(a). It is evident that the activity of this enzyme in both root and leaves is stimulated at all salinity levels except 300 mM NaCl, at which it has decreased in the roots. Enhancement in the activity of acid phosphatase in the leaves of spinach grown under saline condition has been reported by Pan.<sup>[18]</sup> Similar observations have been made by Karadge and Chavan<sup>[19]</sup> in *Sesbania*. Lila Arab and Ehsanpour<sup>[20]</sup> measured acid phosphatase activity in the leaf and stem of *in vitro* grown *Medicago sativa* under saline conditions and found that the activity was increased due to increasing salt concentration. Chakrabarti and Mukharji<sup>[21]</sup> have also found that the salt stress caused to increase the activity of acid phosphatase in the leaf and roots of mung bean. Parida and Das<sup>[22]</sup> studied effect of various levels of salinity (0, 100, 200, 400mM NaCl) on the activity of acid phosphatase in *Bruguiera parviflora* growing under hydroponic culture. Their experiments also revealed that the salinity causes stimulation of activity of this enzyme.

Effect of NaCl salinity on the activity of enzyme alkaline phosphatase in the leaves and roots of *Prosopis juliflora* is depicted in the figure 2 (b). It is evident that the activity of this enzyme is decreased in the root and leaves with increasing level of salt in the medium. Weimberg<sup>[23]</sup> noticed a decrease in the level of alkaline phosphatase in pea seedlings due to NaCl salinity. A contrasting behavior of acid and alkaline phosphatases under saline conditions was noticed by Ahmad and Huq<sup>[24]</sup> in halophytic spinach. In the case of horsegram only lower concentration of salt (25 mM of NaCl) caused the real increase in alkaline phosphatase activity.<sup>[25]</sup> Parida and Das<sup>[22]</sup> noticed that the activity of this enzyme in a mangrove, *Bruguiera parviflora* was increased under varying levels of salinity (0, 100, 200, 300 mM NaCl). The effect of salt stress on alkaline phosphatase was studied by Pan<sup>[26]</sup> in Spinach. He found that the enzyme alkaline phosphatase was inhibited by salinity (> 150 mM NaCl). In case of *Prosopis juliflora* a trend more or less similar to that in Spinach and pea is evident in both root and leaf tissues. Acid phosphatase and alkaline phosphatase in the root and leaves of this plant, however have shown an opposite trend. A difference in ionic balance resulting in a shift in cellular pH might be a reason for such alterations.

Effect of NaCl salinity on enzyme ATPase in the leaves and roots of *Prosopis juliflora* is shown in figure 2(c). It is evident that the activity of enzyme ATPase in the root was

Download English Version:

<https://daneshyari.com/en/article/2495893>

Download Persian Version:

<https://daneshyari.com/article/2495893>

[Daneshyari.com](https://daneshyari.com)