

SHORT COMMUNICATION

Nuphar lutea: *In vitro* anti-leishmanial activity against *Leishmania major* promastigotes and amastigotes

J. El-On^{a,b,*}, L. Ozer^{a,b}, J. Gopas^{a,c}, R. Sneir^{a,b}, A. Golan-Goldhirsh^d

^aThe Shraga Segal Department of Microbiology and Immunology, Faculty of Health Sciences, Ben-Gurion University of the Negev, Israel

^bLaboratory of Parasitology, Soroka University Medical Center, Beer Sheva, Israel

^cDepartment of Oncology, Soroka University Medical Center, Israel

^dAlbert Katz Department of Dryland Biotechnologies, The Jacob Blaustein Institutes for Desert Research, Sede Boqer Campus, Ben-Gurion University of the Negev, Beer Sheva, Israel

Abstract

Several anti-leishmanial drugs of choice are of plant origin. Many of the available drugs against the disease are toxic and in certain cases parasite drug resistance is developed. The development of new compounds is urgently required.

Aims of the study: To determine the leishmanicidal activity of the *Nuphar lutea* plant extract against *Leishmania major* *in vitro*.

Materials and methods: The leishmanicidal activity of methanolic plant extract against *L. major* free living promastigotes and intracellular amastigotes was evaluated, using microscopic examinations and the enzymatic XTT assay.

Results: Methanolic extract of *N. lutea* was highly effective against both *Leishmania* promastigotes and *L. amastigotes* ($IC_{50} = 2 \pm 0.12 \mu\text{g/ml}$; $ID_{50} = 0.65 \pm 0.023 \mu\text{g/ml}$; $LD_{50} = 2.1 \pm 0.096 \mu\text{g/ml}$, $STI = 3.23$). The extract at $1.25 \mu\text{g/ml}$ totally eliminated the intracellular parasites within 3 days of treatment. Also, a synergistic anti-leishmanial activity was demonstrated with *N. lutea* extract combined with the anti-leishmanial drug – paromomycin. The partially purified *N. lutea* active component was found to be a thermo-stable alkaloid(s) with no electrical charge and is resistant to boiling and to methanol, dichloromethane and xylene treatment.

Conclusions: The present study suggests that *N. lutea* might be a potential source of anti-leishmanial compounds.

© 2009 Published by Elsevier GmbH.

Keywords: *Nuphar lutea*; *Leishmania major*; Amastigotes; Promastigotes

Introduction

Leishmaniasis, a zoonotic protozoan disease caused by *Leishmania*, is still considered a major health problem. In the vertebrate host, *Leishmania* are intracellular parasites

(amastigotes) of the macrophage of the skin and lymphatic organs, and in the intestine of the invertebrate host (sand fly) they are transformed into free living flagellates (promastigotes). The clinical manifestations range from simple cutaneous lesion (CL) to progressive disseminated visceral (VL) fatal disease. The number of effective drugs available against the disease (sodium stibogluconate, pentamidine, amphotericin B, miltefosine, paromomycin) is extremely limited, and the development of resistant parasites to most drugs has been

*Corresponding author at: The Shraga Segal Department of Microbiology and Immunology, Faculty of Health Sciences, Ben-Gurion University of the Negev, Israel. Tel.: +972 8 6400 899; fax: +972 8 6477 626.

E-mail address: jelon@bgumail.ac.il (J. El-On).

reported (Jha 2006; Santos et al. 2008). In the present study, the anti-leishmanial activity of *N. lutea* extract against *L. major* promastigotes and intracellular amastigotes was evaluated.

Material and methods

Parasite strains, cell cultures and host animals

Male C3H/HeJ, 10–12 weeks old, were used as the source of peritoneal macrophages for *in vitro* studies. *L. major* (WHO code: MHOM/IL/80/Freidlin) was used in all experiments. The parasites were maintained at 28 °C by biweekly passage in RPMI-1640 plus L-glutamine (20 mM), supplemented with 10% heat inactivated FCS and the antibiotics: 100 µg streptomycin and 100 units of penicillin per ml. Parasites were also maintained as a stablate at –70 °C and *in vivo* in experimentally infected Balb/c mice. Mice were treated according to the NIH guidelines for Animal Care, using a protocol approved by the Animal Ethical Care Committee of Ben-Gurion University.

Plant collection and extracts preparation

N. lutea (mainly leaves) was collected by Prof. Avi Golan, at various seasons from natural habitats in Israel. Additional samples were collected in the Yarkon-Natural Water Reserve in Central Israel. Immediately after harvesting, the plant material was submerged in liquid nitrogen and afterwards at –80 °C until extracted. For the 50% methanol:water (v/v) extraction, samples (1 g) of frozen plant material were ground in a pre-chilled mortar containing liquid nitrogen. Two ml of 50% methanol/water (v/v) were added, and the slurry was mixed and kept on ice for 15 min. After centrifugation at 11,000 rpm for 5 min at room temperature the supernatant was collected and stored at –80 °C for analysis. Larger samples of the plant were further collected, washed with sterile distilled water, freeze dried and afterwards powdered. Concentrations were determined based on dry weight per volume. These samples were further extracted using 2 additional procedures, including: (a) water extraction, boiling at acidic condition, followed by xylene treatment at acidic and basic conditions using a standard procedure. Shortly, grinded powder was first extracted three times with distilled water at 4 °C. The mixture was incubated over 3 days at 4 °C with a gentle mixing using a magnetic stirrer, and then filtered twice through a Whatman-No. 3 filter paper and centrifuged for 10 min at 3000 rpm at 4 °C. The filtrate was acidified to pH 3.0 with 1 M HCl, and after 2 h incubation with gentle mixing at 4 °C it was boiled over 20 min. The resulting precipitate was

filtered out and the filtrate was extracted three times with xylene. The aqueous layer was collected and alkalized (pH-10) with 5 M NaOH and extracted again with xylene. The xylene fraction was collected, acidified with 1 M HCl and after gentle mixing the aqueous layer containing the alkaloids was collected, neutralized by NaOH and concentrated by freeze drying. (b) Methanolic extract was fractionated on a silica gel column using chloroform: ethyl-acetate: diethyl-amine 20:1:1 (v:v:v), as eluant. Column fractions were collected and assayed for anti-leishmanial activity.

Anti-leishmanial efficacy

The anti-leishmanial activity against promastigotes, in culture at 28 °C and amastigotes in C3H mouse macrophages at 37 °C, was determined. Parasites and macrophages were cultivated in RPMI medium containing 10% FCS plus antibiotics. Logarithmic dilutions followed by serial drug dilutions of each extract were tested. RPMI 1640 medium, fetal calf serum (FCS), the antibiotics mixture of gentamycin/streptomycin and glutamine were purchased from Biological Industries (Beit Haemek, Israel). Controls samples included: (a) Growth medium without extracts (100% viability). (b) Appropriate solvent, at the highest concentration to which cells were exposed with the extracts. (c) Positive control, containing the anti-leishmanial drug, paromomycin sulfate (PR) (Farmitalia, Italy).

Effect on *L. major* promastigotes *in vitro*, at 28 °C

Activity of plant extracts against promastigotes was determined by both microscopic counting and the colorimetric cell viability XTT (2,3-bis[2-methoxy-4-nitro-5-sulfophenyl]-2H-tetrazolium-5-carboxanilide) (Sigma Chemical Co; St. Louis, MO) assay (Williams et al. 2003). Promastigotes ($2.5 \times 10^6/100 \mu\text{l}/\text{well}$) from a logarithmic growth phase culture in complete medium were seeded into a flat-bottomed 96-well plastic tissue-cultured microplates, in triplicates, and either logarithmic or serial dilutions of each extract (100 µl) was added. After 3 days incubation at 28 °C, 25 µl of XTT (0.2 mg/ml) were added to each well, followed by an additional 3–5 h incubation at 37 °C. The optimal density (OD) at 450/650 nm was measured using an ELISA plate reader (Dynatech MR5000, USA). The 50% lethal dose (IC₅₀) was evaluated graphically by plotting concentration versus percentage growth inhibition. The anti-leishmanial activity was further determined by microscopic counting of the free living parasites for each well and the percentage growth inhibition for each concentration of extract was calculated in relation to control.

Download English Version:

<https://daneshyari.com/en/article/2497196>

Download Persian Version:

<https://daneshyari.com/article/2497196>

[Daneshyari.com](https://daneshyari.com)