

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.jfda-online.com

Original Article

Clove (*Syzygium aromaticum*) ingredients affect lymphocyte subtypes expansion and cytokine profile responses: An *in vitro* evaluation



Shaghayegh Pishkhan Dibazar, Shirin Fateh, Saeed Daneshmandi*

Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

ARTICLE INFO

Article history:

Received 8 December 2013

Received in revised form

16 April 2014

Accepted 25 April 2014

Available online 28 October 2014

Keywords:

B cells
Clove extract
Cytokine
In vitro
Splenocytes
T cells

ABSTRACT

Clove (*Syzygium aromaticum*) has been used in folk medicine in many disorders. The present work aimed to investigate effects of clove essential oil as eugenol and water soluble ingredients on mouse splenocytes. Clove extracts were harvested and in different concentrations (0.001–1000 $\mu\text{g/mL}$) were affected to splenocytes and also phytohemagglutinin (PHA = 5 $\mu\text{g/mL}$) and lipopolysaccharide (LPS = 10 $\mu\text{g/mL}$) activated splenocytes; then splenocytes proliferation assayed using the MTT ([3-(4, 5-dimethylthiazole-2-yl) -2, 5-diphenyl tetrazolium bromide]) method were done. On the culture supernatant interferon (IFN)- γ , interleukin (IL)-4, IL-10, and transforming growth factor (TGF)- β cytokines were measured. Clove ingredients (100 $\mu\text{g/mL}$ and 1000 $\mu\text{g/mL}$) reduced PHA stimulated splenocytes proliferation and enhanced LPS stimulated cells expansion. Treated splenocytes showed suppression of IFN- γ release and induction of IL-4, IL-10, and TGF- β secretion (in the range of 0.1–1000 $\mu\text{g/mL}$). The results of this study suggest clove extracts could suppress the T cell cellular immunity and enhance humoral immune responses. In clove affection cytokine pattern shifted toward modulatory and Th2 responses and accelerator of humoral immunity cytokines.

Copyright © 2014, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

Herbal medicine is used to treat different diseases in most of the world's cultures. Identification of the active component and mechanisms of action of traditional medicines on the immune system is highly desirable [1]. The immune system is divided into adaptive and innate systems and the adaptive immune system is divided into cellular and humoral responses. T and B lymphocytes are the main cells that are involved in adaptive immune phenomena. Many of the drugs

and mediators in the treatment of the disease perform their functions by the influence of immune cells, especially lymphocytes. These agents could be body mediators, synthetic factors, or natural elements [2,3]. Cloves are dried flower buds derived from an evergreen tree *Syzygium aromaticum* (L.) Merr. & Perry (i.e., *Eugenia aromaticum* or *Eugenia caryophyllata*), a tree 10–20 m high that is indigenous to India, Indonesia, Zanzibar, Mauritius, and Iran [4]. It is commonly used in Africa, Asia, and other parts of the world in the preparation of various spicy rich dishes. It has a deep brown color, intense fragrance, and a burning taste [5]. Clove has several therapeutic properties; it is

* Corresponding author. Department of Immunology, Faculty of Medical Sciences, Jalale Ale-Ahmad Exp.Way, Tarbiat Modares University, Tehran, Iran.

E-mail address: daneshmandi2006@yahoo.com (S. Daneshmandi).

<http://dx.doi.org/10.1016/j.jfda.2014.04.005>

1021-9498/Copyright © 2014, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

a well-known food flavor and a popular remedy for dental and respiratory disorders, headache, and sore throat in traditional medicines of Australia, and in Asian countries [6]. Clove possesses antidiabetic, anti-inflammatory, antithrombotic, anesthetic, pain-relieving, and insect-repellent properties [7,8]. The main ingredients of clove are eugenol (50–87%), eugenyl acetate, tanene, thymol, and β -caryophyllene [9]. These components are responsible for clove extract's effect when used under different conditions. In addition, these components have been shown to modulate some immune responses, including anti-inflammatory effects, although the mechanisms through which these effects are mediated remain unclear [10]. In addition to eugenol, which is extracted using alcoholic method [11], water-soluble ingredients also have an important function. Biological and therapeutic properties of clove have been reported, but its effect on the immune system is poorly investigated. Those who have studied the benefits of clove in traditional medicine propose its effects on the immune cells and immune system. The goal of this study was to investigate *in vitro* effects of the essential oil in eugenol (prepared by alcoholic extraction) and water-soluble elements of clove on splenocytes and mitogen-stimulated splenocytes for analysis of T and B lymphocyte subtypes [phytohemagglutinin (PHA (Sigma, St. Louis, MO)) mitogen for T cells and lipopolysaccharide (LPS (Sigma, St. Louis, MO)) mitogen for B cells] as the main effector cells in cellular and humoral immune responses. Moreover, the effect on cytokines secretion (IFN- γ , IL-4, IL-10, and TGF- β) by BALB/c splenic lymphocytes was assessed.

2. Materials and methods

2.1. Animals

Six- to 8-week-old female inbred Balb/c mice were purchased from the Pasteur Institute of Iran (Tehran, Iran). They were kept in the animal house of Tarbiat Modares University, Tehran, Iran and were given standard mouse chow and sterilized water throughout the study. The study design was approved by the Ethical Committee of the Department (Tarbiat Modares University, Tehran, Iran) for Animal Care and Use.

2.2. Plant material

The flower buds of clove (*Syzygium aromaticum*) were collected from plants cultivated in the Center of Medicinal Plants Research 25 km north of Tehran, Iran, and confirmed by the Center of Agricultural Research, Tehran, Iran.

2.3. Preparation of clove extracts

Extraction of eugenol (in the essential oil) was done by the alcoholic extraction method [11]; in brief, dried powdered flower buds of clove (25 g) were soaked in 100 mL of ethanol to prepare the essential oil and in 100 mL of distilled water to prepare the aqueous extract. They were allowed to mix on a rotary for 24 hours and then filtered using Whatman No. 1 filter paper. The filtrate extract was poured into a special lyophilizing flask. The flask was connected to a vacuum pump and evacuated until

drying. The isolates were recovered, weighed, and representative stock of 100 mg/mL was prepared in 5 mL of, accordingly, water or 50% ethanol for use in the various assays herein. Yields for both materials were routinely 2–3 g (8–12% [w/w] starting material).

2.4. Analyses of extract contents

Previous reports indicated that clove oil contained eugenol (~75%), β -caryophyllene (~5%), eugenyl acetate (~16%), and other components <1% [12]; analyses of the aqueous extracts showed it contained 49% eugenol [13]. The extracts in the current study were assessed using the same gas chromatography-mass spectrometry methods as that of Lee et al [13]; the data showed eugenol made up 74% of material in the ethanolic extract and 43% in aqueous extract. Although we did not assess directly the levels of β -caryophyllene and eugenyl acetate, we assume these are in line with the values reported in both Chaieb et al [12] and Lee et al [13] and thus comprise the bulk of the remaining materials in each extract.

2.5. Preparation and treatment of splenocytes

The mice were decapitated under mild diethyl ether anesthesia and the spleens were rapidly excised under sterile conditions. This tissue was subsequently homogenized in 10 mL cold RPMI 1640 complete medium (Sigma Chemical Company, St. Louis, MO) in a glass homogenizer. Homogenized spleen tissues passed through a fine steel mesh to obtain a homogeneous cell suspension. The erythrocytes were osmotically lysed by 0.75% NH₄Cl in Tris buffer (0.02%, pH = 7.2). After centrifugation (360g at 4°C for 10 minutes), the pelleted cells were washed three times with phosphate buffered saline (PBS) and resuspended in RPMI 1640 complete medium supplemented with 11 mM sodium bicarbonate, 2 mM L-glutamine, 100 U/mL penicillin, 100 μ g/mL streptomycin, and 10% fetal bovine serum (all reagents were purchased from Gibco, Grand Island, NY). The total viable splenocyte cell count was determined using the trypan blue exclusion assay. Cell viability was more than 95%. Splenocyte treatment was assayed as follow: 5×10^6 cell/mL of the cells was seeded into each well of a 96-well flat-bottom microtiter plate (Nunc, Kamstrup, Denmark) in complete medium and PHA (final concentration 5 μ g/mL), LPS (final concentration 10 μ g/mL), or PBS was added to the wells. Clove ingredients (final concentration 0.001–1000 μ g/mL) were added, giving a final volume of 200 μ L (tetraplicate wells) and incubated for 48 hours at 37°C and 5% CO₂.

2.6. Cell proliferation assay

After 48 hours of incubation with various concentrations of clove extracts, cell proliferation was measured based on the MTT [3-(4, 5-dimethylthiazole-2-yl) -2, 5-diphenyl tetrazolium bromide] reduction assay [14]. In brief, after incubation, 20 μ L of MTT (5 mg/mL in PBS) were added to 200 μ L wells (in one tenth of the total volume) and incubated for 4 hours at 37°C and 5% CO₂. Then medium was removed and the formazan blue crystals, which formed by reacting MTT with mitochondrial dehydrogenase in the living cells, were dissolved by 100 μ L of acidic isopropanol (0.04 M HCl in isopropanol). The

Download English Version:

<https://daneshyari.com/en/article/2507357>

Download Persian Version:

<https://daneshyari.com/article/2507357>

[Daneshyari.com](https://daneshyari.com)