

Serum lipid levels and suicidality among male patients with schizoaffective disorder

Darko Marčinko *, Vesna Marčinko, Dalibor Karlović, Anita Marčinko, Marko Martinac,
Dražen Begić, Miro Jakovljević

Department of Psychiatry, University Hospital Zagreb, Kišpatićeva 12, 10000 Zagreb, Croatia

Received 23 April 2007; received in revised form 12 August 2007; accepted 12 August 2007

Available online 19 August 2007

Abstract

Suicidal behavior in schizoaffective disorder is a serious problem and suicide risk during lifetime ranges between 5%–10%. Neurobiology of suicidal behavior has not been studied sufficiently, and a high number of studies are oriented toward lipid investigation. The aim of our study was to investigate whether there were differences in the level of lipids (cholesterol, HDL-cholesterol, LDL-cholesterol, triglycerides) in hospitalized suicidal ($n=20$) and non-suicidal ($n=20$) patients with schizoaffective disorder. The study also included male healthy control subjects ($n=20$). Hamilton Depression Rating Scale (HDRS-17), and Positive and Negative Syndrome Scale (PANSS) were used to confirm the level of psychopathology in patients with schizoaffective disorder. Severity of suicidality was measured by Scale for Suicide Ideation (SSI) at time of admission. Results of the study indicated significantly lower concentrations of cholesterol ($p<0.001$), LDL-cholesterol ($p<0.01$) and HDL-cholesterol ($p<0.01$). There were no differences in the number of previous hospitalization and previous suicide attempts between suicidal and non-suicidal patients ($p>0.05$). Duration of the illness was significantly ($p<0.05$) shorter in suicidal patients. Suicidal patients also had a significantly higher score on HDRS-17 ($p<0.001$) and PANSS ($p<0.01$) compared to non-suicidal patients.

© 2007 Elsevier Inc. All rights reserved.

Keywords: Cholesterol; LDL-cholesterol; Lipids; Men; Schizoaffective disorder; Suicidality

1. Introduction

Changes in serum concentration of lipids, particularly of cholesterol, are among the most studied biological factors in the field of suicidality. In the early 1990s, analyses of mortality rates in prevention-related clinical trials showed a correlation between low cholesterol and non-illness-related mortality (Muldoon et al., 1990; Lindberg et al., 1992; Neaton et al., 1992). Further analyses in extensive studies demonstrated the occurrence of suicide as a secondary complication, and investigations showed association between low cholesterol

and recurrent suicide attempts (Zureik et al., 1996; Partonen et al., 1999). Case–control studies indicated lower cholesterol levels in patients admitted after suicide attempt as compared to admitted non-suicidal patients (Garland et al., 2000; Kim et al., 2002; Diaz-Sastre et al., 2007). Correlation of low cholesterol level with suicidality has also been confirmed in other psychiatric disorders, such as anorexia nervosa (Favaro et al., 2004) and panic disorder (Ozer et al., 2004). Our previous studies confirmed the role of cholesterol in suicidal behavior of patients diagnosed with different form of psychotic disorders (Marčinko et al., 2004, 2005, 2007).

Some studies demonstrated that association of low cholesterol with suicide attempts was more pronounced among violent suicide attempters or those with a history of such attempts (Alvarez et al., 2000; Vevera et al., 2003; Kim and Myint, 2004; Marčinko et al., 2005). One study corroborated causative correlation between low cholesterol and violence in general (Golomb, 1998). There have, however, been some case–control studies that failed to show an association between cholesterol

Abbreviations: BMI, Body mass index; HDL-C, High density lipoprotein cholesterol; HDRS-17, Hamilton Depression Rating Scale; LDL-C, Low density lipoprotein cholesterol; PANSS, Positive and Negative Syndrome Scale; SSI, Scale for Suicide Ideation.

* Corresponding author. Tel.: +385 1 2421 603; fax: +385 1 2388 329.

E-mail address: darko.marcinko@zg.t-com.hr (D. Marčinko).

level and suicide attempts (Almeida-Montes et al., 2000; Deisenhammer et al., 2004). Also, some previous epidemiological and clinical studies have presented conflicting results on possible associations between low cholesterol levels and heightened rates of death from suicide (Tanskanen et al., 2000; Muldoon et al., 2001). Genetic factors have also been investigated as part of the etiology of suicidal behavior (Turecki, 2001). Regarding neurobiological research of suicidal behavior in psychotic disorders, most studies deal with schizophrenia which actually motivated us to conduct this investigation with schizoaffective disorder. The aim of this study was to investigate if there were differences in lipid concentrations (cholesterol, HDL cholesterol, LDL-cholesterol, triglycerides) between suicidal and non-suicidal patients with schizoaffective disorder on hospital treatment, and a healthy, control group.

2. Methods

2.1. Sample

Subjects were male patients ($N=40$) treated at the Department of Psychiatry, University Hospital Zagreb, during the period of 18 months. Within patients, 20 patients were consecutively admitted suicidal men with schizoaffective disorder, and 20 patients were consecutively admitted men with schizoaffective disorder, without suicidal behavior. The diagnosis of schizoaffective disorder was made according to diagnostic criteria of the International Statistical Classification of Diseases and Related Health Problems, 10th Revision (ICD-10) after performing structured diagnostic psychiatric interview (WHO, 1996). All of the patients have depressed type of schizoaffective disorder. Hamilton Depression Rating Scale, HDRS-17 (Hamilton, 1960) was applied to assess depressive symptoms. Positive and negative syndrome scale, PANSS (Kay et al., 1987) was used to estimate psychotic symptoms. The trained psychiatrists performed clinical evaluation. Patients were classified as suicidal at the hospital admission if a suicidal ideation, or a suicide attempt, or both, was present. Suicidality was assessed positive if item 3 of the HDRS-17 scale was scored positively. Severity of suicidality was measured by Scale for Suicide Ideation (SSI) (Beck et al., 1979) at the time of admission for all of the participants.

All participants were free of all medications in the previous 3 months. Healthy male subjects ($N=20$) with no history of psychiatric illness and somatic disorders or diseases were randomly selected as a control group. They were recruited from the general population and were closely matched for age. All subjects gave written informed consent to participate in the study. This study was approved by the Clinical Hospital Center Medical Ethics Committee.

Only patients whose biochemical analyses from the time of admission were available were included in the study. Venous blood samples were collected within 24 h of admission. None of the patients or control subjects included in the study had used any cholesterol-lowering drugs before the sampling. The exclusion criteria were: hypertension, hypothyroidism, diabetes mellitus, disorders of the lipoprotein metabolism, diagnosis of

substance abuse, including alcoholism, eating disorders and organic brain syndrome.

2.2. Assessment

Venipuncture was performed for all subjects between 8 and 9 a.m. after 12 h overnight fast. Immediately after collecting blood samples, serum concentrations of cholesterol, HDL-cholesterol and triglycerides were determined using enzyme method and commercial kits (Olympus Diagnostic, GmbH, Hamburg, Germany) on Olympus AU 600 automated analyzer. Inter-assay laboratory coefficients of variation were 3.2% for cholesterol, 2.5% for triglycerides, and 3.0% for HDL-cholesterol. Reference intervals for the measured parameters were as follows: cholesterol <5.0 mmol/L, LDL <3.0 mmol/L, HDL >1.0 mmol/L, and triglycerides <1.7 mmol/L. Height and weight of each patient were recorded while they were standing barefoot in light clothes on the medical scale that measures height and weight. Body mass index (BMI) was calculated in the way that body weight in kilograms was divided with squared height value in meters.

2.3. Data analysis

The results were expressed as mean values and standard deviations. Kruskal Wallis one way analysis of variance was applied for statistical processing of the obtained data during comparison of continuous variables across the three subject groups (suicidal patients, non-suicidal patients and healthy controls). Student *t*-test was used to compare two groups of patients after normal data distribution in some variables had been demonstrated by Kolmogorov–Smirnov test. Nonparametric Spearman correlation coefficient test was applied in searching relations between variables. In all tests, the criterion of significance was $p<0.05$. Statistical processing of results was done by using commercial statistical package SPSS 11 (SPSS for Windows 11.0, SPSS, Chicago, IL, USA).

3. Results

Results of the study are presented in Table 1. There were no statistically significant differences for BMI and age values between subject groups ($p>0.05$). Analysis showed that total cholesterol values in suicidal patients were statistically significantly lower than in non-suicidal patients and the control group ($p=0.000$). Triglyceride levels in suicidal patients were not statistically significantly different compared to non-suicidal patients and the control group ($p=0.255$). HDL-cholesterol concentrations in suicidal patients were statistically significantly lower than in non-suicidal patients and the control group ($p=0.007$), also the values of LDL-cholesterol ($p=0.001$). SSI values in suicidal patients, were statistically significantly higher than in non-suicidal patients ($p=0.000$; $df=38$). In suicidal patients, PANSS values were statistically significantly higher than in non-suicidal patients ($p=0.004$, $df=38$), and the HDRS-17 score was also statistically significantly higher in suicidal patients than in non-suicidal ones ($p=0.000$, $df=38$). We have found no

Download English Version:

<https://daneshyari.com/en/article/2565718>

Download Persian Version:

<https://daneshyari.com/article/2565718>

[Daneshyari.com](https://daneshyari.com)