



ELSEVIER

Contents lists available at ScienceDirect

## Toxicology Reports

journal homepage: [www.elsevier.com/locate/toxrep](http://www.elsevier.com/locate/toxrep)

## Modulatory effect of fibre-enriched cake on alloxan-induced diabetic toxicity in rat brain tissues



Ochuko L. Erukainure<sup>a,b,\*</sup>, Osaretin A.T. Ebuehi<sup>b</sup>, Folasade O. Adeboyejo<sup>a</sup>, Muhammad Aliyu<sup>c</sup>, Gloria N. Elemo<sup>a</sup>

<sup>a</sup> Department of Food Technology, Federal Institute of Industrial Research, Oshodi, Lagos, Nigeria

<sup>b</sup> Department of Biochemistry, University of Lagos, Lagos, Nigeria

<sup>c</sup> Department of Biochemistry, Ahmadu Bello University, Zaria, Nigeria

### ARTICLE INFO

#### Article history:

Received 8 May 2014

Received in revised form 10 July 2014

Accepted 15 July 2014

Available online 24 July 2014

#### Keywords:

Antioxidant

Diabetes

Food fibre

Oxidative stress

### ABSTRACT

Diabetes is a metabolic disorder characterized by hyperglycaemia and it is fast becoming a scourge in sub-Saharan Africa. The nutritional properties of developed fibre-enriched cake and its protective potential against diabetic induced neurotoxicity in rats were investigated. Fibre-enriched cake was developed from selected fruits and analysed for its nutritional and sensory attributes. Rats were induced with diabetes by a single intraperitoneal injection of alloxan and treated with the formulated cake. After 14 days treatment, the rats were sacrificed by cervical dislocation. Their brain tissues were accessed for reduced glutathione (GSH), catalase, superoxide dismutase (SOD) activities, protein content and lipid peroxidation as well as lipid profiles which cover for total cholesterol, triglycerides, HDL and LDL. Induction of diabetes led to significant reduction ( $p < 0.05$ ) of GSH, catalase, SOD activities and protein content. Feeding on the formulated cake led to their significant increase. Decreased lipid peroxidation, total cholesterol, LDL and triglycerides, and increased concentration of HDL were also observed on feeding with the cake. These results indicate an antioxidant protective potential of the fibre-enriched cake against diabetic-induced brain toxicity. Thus, it can serve as an adjunct to dietary therapy for diabetes.

© 2014 The Authors. Published by Elsevier Ireland Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

### 1. Introduction

Diabetes have been described as a growing scourge which affects more than 12 million people in sub-Saharan Africa (SSA), causing a major drain on her health resources already overburdened by other infectious diseases [19]. It is a metabolic disorder characterized by hyperglycaemia and is associated with long-term vascular complications such as retinopathy, nephropathy, cardiopathy, and neuropathy [20]. Oxidative stress has been reported to play a

major role in these complications. A number of studies have implicated hyperglycaemia-induced oxidative stress in the aetiology of a variety functional and structural disorder in the central and peripheral nervous system [13,20,10].

Over the years, it has been documented that medicinal plants are very effective in the treatment and management of diabetes [22]. The combination of two basic central factors, food and medication has been attributed to their utilization [14]. These plants are major source of fibre with tremendous health benefits [9]. The health benefits of dietary fibre have been reported in several studies. Diabetes prevalence has been shown to correlate with fibre intake among various populations [32,31]. Its consistent consumption has been reported to cause a reduction in blood glucose concentrations [31]. In previous study we

\* Corresponding author at: Department of Food Technology, Federal Institute of Industrial Research, Oshodi, Lagos, Nigeria.  
Tel.: +234 8062179388.

E-mail address: [loreks@yahoo.co.uk](mailto:loreks@yahoo.co.uk) (O.L. Erukainure).

reported the hypoglycemic and antidiabetic effect of fibre-enriched snacks in diabetic rats [8].

This paper is a continuation of our study on the effect fibre-enriched snacks on diabetes. It aims at reporting the nutritional properties of the cake and its protective potential against diabetic induced neurotoxicity in rats.

## 2. Materials and methods

### 2.1. Plant materials

Banana (*Musa species*), oranges (*Citrus sinensis*), watermelon (*Citrullus lanatus*), pineapple (*Ananas cosmosus*) and pawpaw (*Carica papaya*) were purchased from Ketu fruit market, Ketu, Lagos, Nigeria. They were processed into fibre paste as described by Erukainure et al. [10,9].

### 2.2. Production of high fibre cake

High fibre cake was produced as described by Erukainure et al. [9].

### 2.3. Proximate analysis

The proximate nutritional qualities of the cake sample were carried out according to the method of AOAC [3], which covers for total protein, ash, fat, crude fibre, carbohydrates and reducing sugar, respectively.

### 2.4. Sensory attributes

Sensory evaluation was conducted on the developed cake according to the method described by Ihekoronye and Ngoddy [17]. It was compared to readily available commercial cake. They were given the reference codes YLQ and PGE for the developed and commercial cake respectively. The coded samples were presented to a 10-men panellist to evaluate for the attributes: colour, texture, taste, crumb grains, mouth feel, aroma, and overall acceptability. Scores were given to the scales: (9) extremely acceptable, (8) very acceptable, (7) moderately acceptable, (6) slightly acceptable, (5) neither acceptable nor unacceptable, (4) slightly unacceptable, (3) moderately unacceptable, (2) very unacceptable and (1) extremely unacceptable.

### 2.5. Animals

Eighteen male albino rats of Wister strain weighing about 150–200 g were used for the study. They were fed on standard rat pellet diet (Ladoko feeds) and allowed to adapt for one week. They were provided water *ad libitum* and maintained under standard laboratory conditions of natural photoperiod of 12-h light–dark cycle. The animals used in the present study were maintained in accordance with the approval of the Animal Ethical Committee, University of Lagos, Lagos, Nigeria. The approval number from the Animal Institutional Ethical Committee is UL/CMUL/IEC 2011/1003.

### 2.6. Induction of diabetes

Diabetes was induced by a single intraperitoneal injection of 180 mg/kg of alloxan monohydrate in normal saline water in a volume of about 3 mL. After 72 h of alloxan injection, the diabetic rats (glucose level > 250 mg/dl) were separated and used for the study.

### 2.7. Experimental design

The rats were divided into three groups, each consisting of six animals:

- Group 1 – normal rats + pelletized mouse chows.
- Group 2 – diabetic (untreated).
- Group 3 – diabetic + high fibre cake.

Treatment lasted for two weeks. At the end of the feeding trials, the rats were fasted overnight and sacrificed by cervical dislocation.

### 2.8. Preparation of tissue homogenates

The brain tissues were removed, rinsed in ice-cold 1.15% KCl solution to wash off excess blood, blotted dry with filter paper. They were homogenized in four parts of homogenizing buffer and centrifuged at 10,000 rpm for 15 min in an ultracentrifuge at a temperature of  $-2^{\circ}\text{C}$  to get the mitochondrial fraction. The supernatant (post-mitochondrial fraction) was decanted and stored at  $-4^{\circ}\text{C}$  for subsequent analysis. Each time the supernatant was outside the freezer, it was kept in ice bags.

The protein content of the tissue fractions of the organs were determined by Lowry's method using bovine serum albumin (BSA) as standard [24].

### 2.9. Determination of oxidative stress parameters

Lipid peroxidation was determined by measuring malondialdehyde (MDA) formed by thiobarbituric acid reaction (TBAR) [6]. Catalase (CAT) activity was estimated by measuring the rate of decomposition of  $\text{H}_2\text{O}_2$  [2]. The level of superoxide dismutase (SOD) activity was determined by the method of Misra and Fridovich [26]. While the method of Ellman [7] was adopted in estimating the activity of reduced glutathione (GSH).

### 2.10. Determination of hypolipidemic activities

Tissue total cholesterol, triglyceride and high density lipoprotein (HDL) were measured by enzymatic colorimetric method using Randox kits according to manufacturer's protocol. The concentration of low-density lipoprotein (LDL) cholesterol was calculated by the formula of Friedwald et al. [12].

### 2.11. Statistical analysis

To address the biological variability, each set of experiments was repeated at least three times ( $n=3$ ) for proximate analysis and six times for experimental rats

Download English Version:

<https://daneshyari.com/en/article/2572331>

Download Persian Version:

<https://daneshyari.com/article/2572331>

[Daneshyari.com](https://daneshyari.com)