

Rising Threat; Bonsai

Gokhan AKSEL,¹ Oner BOZAN,¹ Mine KAYACI,¹ Ozlem GUNEYSEL,² Seckin Bahar SEZGIN¹

¹Department of Emergency Medicine Clinic, Umraniye Training and Research Hospital, Istanbul

²Department of Emergency Medicine Clinic, Dr. Lutfi Kirdar Kartal Training and Research Hospital, Istanbul, both in Turkey

SUMMARY

Objectives

In recent years, and especially in the past few months, the number of synthetic cannabinoid (bonsai) users has increased in our country. The aim of this study was to draw attention to the consumption of bonsai among young people and reveal the demographic and basic clinical characteristics of these users.

Methods

This was a retrospective study conducted at the Ümraniye Training and Research Hospital. All of the adult patients (≥ 18 year old) with synthetic cannabinoid intoxication who presented to the Emergency Department throughout the two years of the study (July 1st 2012–June 30th 2014) were enrolled. The frequencies were given as the median and inter-quartile range).

Results

197 patients were included in this study, with 190 male patients (96.4%) and 7 (3.6%) female patients. Two of the four hospitalized patients were exitus, 52 left on their own will and a total of 141 patients were discharged after 6-12 hours of observation in the ED.

Conclusions

The use of synthetic cannabinoids (bonsai) in the recent years, especially in the summer months of 2014 was investigated in this study. Although these patients can have a benign clinical course, the process can also be fatal. It should especially be noted that patients with depressed respiration, low GCS scores and high PaCO₂ values are at higher risk for mortality and the necessity of early intubation should be kept in mind.

Key words: Cannabinoid receptor agonists; emergency medicine.

Introduction

Synthetic cannabinoids (SC) first emerged in 2004 and became rapidly popular, especially among adolescents.^[1,2] SC's have various names worldwide, with the most common one being "spice". However, in Turkey they are commonly known as "bonsai".^[3]

In recent years, especially in the last months, the number of bonsai users has increased in our country.

In this study we aim to draw attention to the consumption of bonsai among young people and reveal the demographic and basic clinical characteristics of these users. This study was the sole study, which mentions the role of synthetic can-

Submitted: November 02, 2014 **Accepted:** November 26, 2014 **Published online:** March 02, 2015

Correspondence: Gokhan AKSEL, M.D. Umraniye Egitim ve Arastirma Hastanesi, Acil Tip Klinigi, Elmalikent Mahallesi, Adem Yavuz Caddesi, No: 1, Umraniye, Istanbul, Turkey.

e-mail: aksel@gokhanaksel.com



nabinoids as a public health problem in Turkey.

Materials and Methods

This was a retrospective descriptive cross-sectional study conducted at Ümraniye Training and Research Hospital. All of the adult patients who were over 17 years old with synthetic cannabinoid intoxication and presented to the Emergency Department (ED) throughout two years of the study (July 1st 2012–June 30th 2014) were enrolled. Patient records

were obtained through the hospital information system (HIS). ICD codes F12.2, Z72.2, T40 and X44 were searched and charts that presented with “bonsai intoxication” were included. Other drug intoxications and patients less than 18 years old were excluded from the study. Suspicious abuse, patient data that was lacking and patients with a history of trauma were also excluded from the study. For the statistical analysis, SPSS for Windows ver. 11.0 (Chicago, IL, USA) was used. The frequencies were given as median and inter-quartile range.

Results

There were 197 patients who were included in this study; 190 patients (96.4%) were male and 7 (3.6%) were female. Two of hospitalized 4 patients (all were intubated) died, 52 left refusing treatment, and a total of 141 patients were discharged after 6–12 hours of observation in the ED. Patients' median systolic blood pressure (SBP) SBP was 120 mmHg, diastolic blood pressure (DBP) was 70 mmHg and pulse rate was 89 beats/min. Median SBP of the four patients who were intubated was 110 mmHg, DBP was 69 mmHg and pulse rate was 71 beats/min. Demographic specifications of the patients and clinical characters are shown in Tables 1–3. During this two-year study the number of abuses rose significantly in the second year, especially in June 2014 as shown in Figures 1 and 2.

Table 1. Demographical findings and vital signs of all of the patients

Variable	Median (IQR)
Age (year)	22 (19-27)
Male (n, %)	190 (96.4)
Pulse (beats/min)	89 (78-105)
SBP (mmHg)	120 (110-126)
DBP (mmHg)	70 (62-80)
SaO ₂ (%-%)	98 (97-99)
GCS	15 (15-15)

IQR: Inter quartile range; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; SaO₂: Saturation of oxygen; GCS: Glasgow coma scale score.

Table 2. Demographical findings and vital signs of the patients who were intubated

Variable	Median (IQR)
Age (year)	29 (23-24)
Male (n, %)	4 (100.0)
Pulse (beats/min)	71 (45-125)
SBP (mmHg)	110 (83-125)
DBP (mmHg)	68 (45-77)
SaO ₂ (%-%)	41 (15-41)
GCS	3 (3-4)

IQR: Inter quartile range; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; SaO₂: Saturation of oxygen; GCS: Glasgow coma scale score.

Discussion

Cannabinoids are separated into three main groups: endogenous, natural and synthetic. SC's were natural-synthetic mixtures sold as legalized marijuana in some countries and smoked in the form of cigarettes.^[3] The most common form of natural cannabinoid is 9-tetrahydrocannabinol (THC).^[3,4] Cannabinoids affect the CB1 and CB2 receptors in the body and they show these effects generally based on a CB1 like mechanism of action, impaired consciousness, sleep changes and cardiovascular effects. Whereas the role of CB2 is poorly understood, it is known that THC acts only through CB1. This is different than SC's, which act both through the CB1 and CB2 receptors and are more effective than THC.^[5-7]

The popularity of these drugs is growing since they cannot

Table 3. Outcomes of intubated patients and their blood gas results

Patient	Age/sex	Clinical result	pH	pO ₂ (mmHg)	PCO ₂ (mmHg)	HCO ₃ (mmHg)
Patient 1	34/male	Intubated+Exitus	6.91	15	128	12
Patient 2	26/male	Intubated+Exitus	6.58	65	173	?
Patient 3	19/male	Intubated+Discharged	7.06	42	61	14
Patient 4	24/male	Intubated+Discharged	6.90	40	119	11

Download English Version:

<https://daneshyari.com/en/article/2604058>

Download Persian Version:

<https://daneshyari.com/article/2604058>

[Daneshyari.com](https://daneshyari.com)