



## Original article

## Investigation of anxiety levels of patients with chest pain admitted to emergency department

Gokhan Basara<sup>b</sup>, Ahmet Baydin<sup>a,\*</sup>, Ahmet Yilmaz<sup>c</sup>, Oguzhan Yucel<sup>d</sup>, Ali Kemal Erenler<sup>b</sup>, Naci Murat<sup>e</sup><sup>a</sup> Ondokuz Mayıs University, Emergency Department, Samsun, Turkey<sup>b</sup> Samsun Education and Research Hospital, Emergency Department, Samsun, Turkey<sup>c</sup> Samsun Training and Research Hospital, Department of Neurology, Samsun, Turkey<sup>d</sup> Samsun Education and Research Hospital of Cardiology, Samsun, Turkey<sup>e</sup> Ondokuz Mayıs University, Department of Statistics, Samsun, Turkey

## ARTICLE INFO

## Article history:

Received 30 January 2015

Received in revised form

22 May 2015

Accepted 2 June 2015

Available online 4 March 2016

## Keywords:

Chest pain

Emergency department

Anxiety

## ABSTRACT

**Introduction:** We aimed to investigate the demographical features, anxiety levels and clinical findings of the patients admitted to our Emergency department (ED) due to chest pain.

**Methods:** Patients with chest pain older than 18 years were included into the study. Demographical features such as age, sex and education level, initial diagnosis in the ED, whether they were hospitalized or coronary intervention performed, were recorded. To determine the anxiety levels of the patients, State-trait Anxiety Inventory (STAI) was performed.

**Results:** Two-hundred and eight adult patients with chest pain were included into the study. We could not determine a relationship between STAI levels of patients according to demographical findings, however, STAI scores tended to decrease by age. Considering the education levels of the patients, it was determined that STAI scores of university graduates were higher than others. The STAI scores of patients discharged from the ED were higher than those hospitalized. When patients were compared according to whether coronary intervention (CI) was performed or not, it was determined that patients who did not require CI had higher STAI scores. When coronary lesion localization of the patients hospitalized was investigated, any relationship could not be determined.

**Conclusion:** In this study, we determined that anxiety levels of the patients with chest pain do not correlate with the severity of the disease. Higher anxiety levels of patients discharged from ED when compared to those with ACS is a challenging problem for both ED physicians and cardiologists.

Copyright © 2016 The Emergency Medicine Association of Turkey. Production and hosting by Elsevier B.V. on behalf of the Owner. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Chest pain is one of the most common complaints among Emergency department (ED) admissions.<sup>1</sup> Determination of life-threatening conditions and making accurate diagnosis are the mainstays of management. In 40–60% of patients with chest pain, life-threatening conditions such as acute coronary syndrome (ACS) are not determined and these patients are being discharged with diagnosis of non-specific chest pain.<sup>2,3</sup> It was also reported that in

80% of the patients with chest pain, cardiologists make a diagnosis of non-cardiac chest pain. In group of patients with non-cardiac chest pain, 50–70% suffer continuous chest pain causing loss of labor, inappropriate hospitalization and unnecessary use of cardiac drugs.<sup>4</sup> In patients with atypical chest pain without an organic reason, it is reported that psychiatric disorders such as panic attack and depression may be seen.<sup>5</sup> In the literature, there are studies reporting that frequency of depression, anxiety and alexithymia is higher than normal controls. In the literature, there is study reporting that frequency of depression, anxiety and alexithymia is higher than normal controls. In the study, it was reported that anxiety, depression and neurotic tendency are more frequently seen in patients with non-cardiac chest pain.<sup>6</sup> In this study, we aimed to determine demographical features, anxiety status and clinical findings of patients admitted to our ED due to chest pain.

\* Corresponding author. Ondokuz Mayıs University, Emergency Department, Kurupelit, Atakum, 55139, Samsun, Turkiye. Tel.: +90 5325839677.

E-mail address: [abaydin@omu.edu.tr](mailto:abaydin@omu.edu.tr) (A. Baydin).

Peer review under responsibility of The Emergency Medicine Association of Turkey.

## 2. Material and methods

After ethical approval from Local Ethic Committee, 208 adult patients with chest pain admitted to our ED were involved into the study. A form called the State-Trait Anxiety Inventory (STAI) was filled by the patients in order to determine anxiety levels. The STAI is a validated 20 item self report assessment device which includes separate measures of state and trait anxiety. The original STAI form was constructed by Charles D. Spielberger, Richard L. Gorsuch, and Robert E. Lushene in 1970. According to the scale, higher scores are positively correlated with higher levels of anxiety.<sup>7</sup> The STAI has been adapted in more than 30 languages for cross-cultural research and clinical practice. It was also adapted to Turkish Society by Onver and Le Conte. In our study, the original version of the form was used (Tables 1a and 1b).

Patients with co-morbidities (pneumonia, acute abdomen, urinary tract infection), acute ST elevation, hypertensive crisis and a history of psychoactive drug use were excluded from the study. Data of the patients was recorded on standard forms according to their demographical features (age, sex, education level), findings of coronary artery disease and STAI scores.

All data was analysed by Statistical Package for Social Sciences for Windows (SPSS) 20.0 programme. For normally distributed data, Student's t test was used in comparison of mean values of two different groups. In comparison of more than two groups, ANOVA (Tukey's) test was used. When data was not normally distributed, Kruskal–Wallis test was used. Then groups were compared by Bonferroni corrected Mann–Whitney U test (p value lower than 0.05 was considered statistically significant). In correction analyses, Spearman correlation analysis was used.

## 3. Results

Of 208 patients with chest pain, 131 were male (63%) and 77 were female (37%). It was determined that as the age increases, STAI scores tended to decrease. When patients were compared according to sex, any statistical significance could not be determined in STAI scores. When education status of the patients was investigated, a statistical significance was determined between literates and illiterates (Table 2). It was also determined that 105 of 208 patients were hospitalized in coronary intensive care unit while 103 were discharged from the ED. Of the 105 patients hospitalized,

70 were diagnosed as unstable angina pectoris (UAP) and 35 as non-STEMI. A statistical significance was determined between patients hospitalized and discharged from the ED according to STAI scores. Those who were discharged had higher scores. Coronary angiography was performed in all patients hospitalized. In 65 of these patients, percutaneous coronary intervention (CI) was performed. In addition, there was a significant statistical between the patients who underwent PTCA and who did not. Patients who did not undergo CI had higher STAI scores. In patients who had no confirmed coronary lesion after intervention, the STAI scores were higher (Table 3).

Patients were also divided into subgroups according to the location of the coronary lesion; Left anterior descending (LAD), right coronary artery (RCA), LAD + Circumflexing (Cx), (LAD + RCA, RCA + Cx, LAD + Cx + RCA). It was determined that there was not any statistical significance in STAI scores according to the location of the lesion.

## 4. Discussion

Anxiety is a major health problem that affects life quality and causes loss of labor and unnecessary health spendings. In a study with anxiety patients, it was determined that pain is the most common complaint with a proportion of 81%.<sup>8</sup> It was reported that female patients with chest pain present with anxiety more frequently.<sup>9</sup> Thus, female patients with chest pain are twice more consulted to cardiologists when compared to male patients. This may result in inappropriate medical treatment and poor results.<sup>10</sup> While male patients reveal psychological symptoms of anxiety such as tension, susceptibility and a premonition of a bad event, female patients tend to present with chest pain, palpitation, shortness of breath and nausea. Besides, while anxiety of male patients is commonly related to sexual problems and return to job, female patients worry about their responsibilities such as care of their children.<sup>11</sup> In our study, there was no difference in anxiety levels of two genders.

As the education level increases, anxiety score increases as well.<sup>9</sup> In concordance, our study revealed that more educated people have higher anxiety levels. However, there is no strong evidence in literature to suggest that there is any relationship between the education status and anxiety level.<sup>12</sup> So, it is possible to say that relationship between education status and anxiety level is

**Table 1a**  
State anxiety inventory (STAI 1).

		Not at all	Somewhat	Moderately so	Very much so
1.	I feel calm	(1)	(2)	(3)	(4)
2.	I feel secure	(1)	(2)	(3)	(4)
3.	I am tense	(1)	(2)	(3)	(4)
4.	I feel strained	(1)	(2)	(3)	(4)
5.	I feel at ease	(1)	(2)	(3)	(4)
6.	I feel upset	(1)	(2)	(3)	(4)
7.	I am presently worrying over possible misfortunes	(1)	(2)	(3)	(4)
8.	I feel satisfied	(1)	(2)	(3)	(4)
9.	I feel frightened	(1)	(2)	(3)	(4)
10.	I feel comfortable	(1)	(2)	(3)	(4)
11.	I feel self-confident	(1)	(2)	(3)	(4)
12.	I feel nervous	(1)	(2)	(3)	(4)
13.	I am jittery	(1)	(2)	(3)	(4)
14.	I feel indecisive	(1)	(2)	(3)	(4)
15.	I am relaxed	(1)	(2)	(3)	(4)
16.	I feel content	(1)	(2)	(3)	(4)
17.	I am worried	(1)	(2)	(3)	(4)
18.	I feel confused	(1)	(2)	(3)	(4)
19.	I feel steady	(1)	(2)	(3)	(4)
20.	I feel pleasant	(1)	(2)	(3)	(4)

Download English Version:

<https://daneshyari.com/en/article/2604100>

Download Persian Version:

<https://daneshyari.com/article/2604100>

[Daneshyari.com](https://daneshyari.com)