

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://www.elsevier.com/locate/poamed>

Original Research Article

Acute myocardial infarction due to the unprotected left main coronary artery disease: The power of TIMI 3 flow



Marcin Sadowski^{a,*}, Wojciech Gutkowski^a, Grzegorz Raczyński^a,
Mariusz Gąsior^b, Marek Gierlotka^b, Lech Poloński^b

^aŚwiętokrzyskie Cardiology Center, Kielce, Poland

^bSilesian Center for Heart Diseases, Zabrze, Poland

ARTICLE INFO

Article history:

Received 21 December 2013

Accepted 24 July 2014

Available online 15 August 2014

Keywords:

Myocardial infarction

TIMI

Left main coronary artery

Percutaneous coronary intervention

Unprotected left main coronary artery

ABSTRACT

Introduction: Acute myocardial infarction (MI) due to unprotected left main coronary artery (ULMCA) as an infarct-related artery is a rare disease with a poor in-hospital and long-term outcome. A nearly linear correlation between post-percutaneous coronary intervention (PCI) Thrombolysis in Myocardial Infarction (TIMI) 3 grade and improved outcome is well known, however poorly reported in regard to ULMCA.

Aim: To assess the impact of pre- and post-procedural TIMI flow grade on outcomes of emergency PCI in men and women with acute MI due to the ULMCA disease.

Material and methods: Data were selected from the ongoing, nationwide, multicenter, prospective, observational registry. The study group consisted of 643 consecutive patients hospitalized during one year with acute MI with ULMCA as an infarct-related artery. Data analyzed included information from patients' history, coronary risk factor profile, clinical presentation, therapeutic approach and adjunctive treatment. The primary end-points were in-hospital, 30-day, and 12-month mortality.

Results and discussion: There were 184 women and 459 men (28.6% vs. 71.4%), $P < 0.0001$. PCI was performed in 120 women and in 279 men (65.2% vs. 60.8%; $P = 0.3$) with a high rate of post-procedural TIMI 3 flow (87.6% vs. 82.1%; $P = 0.17$). Successful PCI defined as the restoration of TIMI flow grade 3 was the only factor decreasing in-hospital (OR 0.1, 95% CI (0.05–0.23), $P < 0.0001$) and 12-month (OR 0.2, 95% CI (0.13–0.3), $P < 0.0001$) mortality.

Conclusions: In patients with acute MI survival after ULMCA PCI depended on TIMI grade before and after the procedure.

© 2014 Warmińsko-Mazurska Izba Lekarska w Olsztynie. Published by Elsevier Urban & Partner Sp. z o.o. All rights reserved.

* Correspondence to: Świętokrzyskie Cardiology Center, Grunwaldzka 45, 25-736 Kielce, Poland. Tel.: +48 41 3671580; fax: +48 41 3671456. E-mail address: emsad@o2.pl (M. Sadowski).

1. Introduction

Acute myocardial infarction (MI) due to unprotected left main coronary artery (ULMCA) as an infarct-related artery (IRA) is a rare disease with a poor in-hospital and long-term outcome, especially when complicated by cardiogenic shock.¹ The beneficial role of immediate either pharmacological or mechanical reperfusion is unquestionable. The degree of reperfusion has been usually assessed by means of Thrombolysis in Myocardial Infarction (TIMI) grade.² A nearly linear correlation between post-percutaneous coronary intervention (PCI) TIMI 3 grade and improved outcome is well known,³ however poorly reported in regard to ULMCA. Contrary to stable patients with ULMCA disease a decision whether to perform PCI or coronary artery by-pass grafting (CABG) is not evidence-based as guidelines on ULMCA revascularization in the setting of acute MI are not unequivocal.^{4,5} Nevertheless, PCI is often the only option for those patients due to hemodynamical and electrical instability as ULMCA supplies a large amount of myocardium. Although in-hospital mortality remains very high (35%–44%), results from some observational studies encourage to perform PCI in this subset of patients.^{6–9} Gender-related differences in the management and outcomes of patients with acute coronary syndromes are well described,¹⁰ however data on population with ULMCA disease are scarce.

2. Aim

The aim of our study was to assess the impact of pre- and post-procedural TIMI flow grade on outcomes of emergency PCI in men and women with acute MI due to ULMCA.

3. Material and methods

The principles of our registry (PL-ACS) have been reported elsewhere and are available on-line.^{11,12} Briefly, this is an ongoing, nationwide, multicenter, prospective, observational mandatory registry of all consecutive ACS cases in Poland. So far, there are over 435 000 cases recorded. The study group consisted of 643 consecutive patients hospitalized during one year with acute MI with ULMCA as an IRA. Data analyzed included information from patients' history, coronary risk factor profile, clinical presentation, therapeutic approach and adjunctive treatment. The primary end-points were in-hospital, 30-day, and 12-month mortality. Mortality data were obtained for all the subjects included. The primary results of our LM study from our registry were reported elsewhere.¹³

3.1. Statistical analysis

Variables were expressed as mean \pm standard deviation, counts and percentages or median and interquartile ranges as appropriate. The significance between groups was tested using Student's *t* test, Mann–Whitney *U* test or Kruskal–Wallis ANOVA test depending on normality as well as homogeneity of variances tested by *F* test. Categorical variables were tested by

χ^2 test. Follow-up mortality was analyzed using the Kaplan–Meier method for multiple-group comparisons. A multivariate Cox proportional hazard regression model was performed to adjust the influence of important clinical factors on mortality. Odds ratios (OR), relative risks (RR) and 95% confidence intervals (CI) were calculated. A two-sided *P*-value less than or equal to 0.05 was considered significant. For all calculations, Statistica 7.1 software (StatSoft, Inc., Tulsa, OK, USA) was used.

4. Results

There were 184 women (28.6%) and 459 men (71.4%). Women were generally older than men (69.0 ± 12.3 vs. 64.5 ± 11.0 years; $P < 0.0001$) with a significant proportion of women over 65 (69.6% vs. 51.2%; $P < 0.0001$), with arterial hypertension (73.9% vs. 64.3%; $P = 0.019$), diabetes (32.1% vs. 21.8%; $P = 0.0063$) and obesity defined as BMI more than 30 kg/m^2 (27.2% vs. 10.7%; $P < 0.0001$). Men more frequently smoked (39.0% vs. 19.0%; $P < 0.0001$). Less women than men presented earlier than 2 hours from symptom onset (8.2% vs. 13.9%; $P = 0.043$). Differences in the incidence of dyslipidemia, prior MI and revascularization, cardiogenic shock, infarct size, left ventricular ejection fraction and extent of the disease (single vs. multivessel) were insignificant. Therapeutic strategy was similar in men and women. PCI was performed in 120 women and in 279 men (65.2% vs. 60.8%; $P = 0.3$) with a high rate of post-procedural TIMI 3 flow (87.6% vs. 82.1%; $P = 0.17$). Intraaortic balloon counterpulsation (IABP) was implanted in 13.6% of women and in 10.7% of men ($P = 0.3$). Tables 1 and 2, and Figs. 1–3 show that survival after ULMCA angioplasty depends on TIMI flow before the procedure. Successful PCI defined as the restoration of TIMI flow grade 3 through the left main coronary artery occurred much more frequently in patients with pre-procedure TIMI grade 3 (Table 3). In multivariate regression analysis a successful PCI assessed as the post-procedural TIMI grade 3 flow was the only factor determining better in-hospital (OR 0.1, 95% CI (0.05–0.23), $P < 0.0001$) and 12-month (OR 0.2, 95% CI (0.13–0.3), $P < 0.0001$) survival while advanced age, STEMI, prior MI and Killip class 3 or 4 on admission significantly increased mortality.¹³

Table 1 – Mortality according to the pre-procedural TIMI flow.

	Women	Men	P
TIMI flow 0 or 1; 242 (61%)			
In-hospital	22 (31%)	52 (30%)	0.85
30-Day	24 (34%)	58 (34%)	0.93
12-Month	29 (41%)	67 (39%)	0.72
TIMI flow 2; 70 (17%)			
In-hospital	5 (28%)	8 (15%)	0.30
30-Day	5 (28%)	9 (17%)	0.33
12-Month	7 (39%)	13 (25%)	0.26
TIMI flow 3; 88 (22%)			
In-hospital	4 (13%)	5 (9%)	0.71
30-Day	5 (16%)	7 (12%)	0.75
12-Month	8 (26%)	13 (23%)	0.74

Download English Version:

<https://daneshyari.com/en/article/2680659>

Download Persian Version:

<https://daneshyari.com/article/2680659>

[Daneshyari.com](https://daneshyari.com)