



Original research

Initial symptom burden predicts duration of symptoms after concussion[☆]



William P. Meehan III^{a,b,c,d,*}, Michael J. O'Brien^{a,b,c}, Ellen Geminiani^{a,b},
Rebekah Mannix^{a,c,d}

^a The Micheli Center for Sports Injury Prevention, United States

^b Sports Concussion Clinic, Division of Sports Medicine, Boston Children's Hospital, United States

^c Brain Injury Center, Boston Children's Hospital, United States

^d Division of Emergency Medicine, Boston Children's Hospital, United States

ARTICLE INFO

Article history:

Received 25 June 2014

Received in revised form 5 November 2015

Accepted 6 December 2015

Available online 12 December 2015

Keywords:

Mild traumatic brain injury

Sports

Head trauma

Sports injury

ABSTRACT

Objectives: To determine which variables predict prolonged (>28 days) duration of symptoms after a concussion.

Design: We conducted a prospective cohort study of adult (>18yo) patients cared for in a specialty concussion clinic.

Methods: Symptoms were assessed using the Post-Concussion Symptom Scale (PCSS) developed at the 3rd International Conference on Concussion in Sports. Possible predictors including age, sex, loss of consciousness, amnesia, history of prior concussion, prior treatment for headaches, history of migraines, and family history of concussions, were measured by self-report. We recorded a PCSS score at each clinical visit and defined time to symptom resolution as the number of days between the date of injury and date of last symptoms.

Results: Of 64 adult patients included in the study, 53.3% were male; 20.3% reported experiencing a loss of consciousness at the time of injury while 23.4% reported amnesia. Patients ranged in age from 18 to 27 years (mean 21 ± 2 years). Most concussions (92.2%) occurred during sports. The mean initial PCSS score for those suffering symptoms for longer than 28 days was significantly higher than those who symptoms resolved within 28 days (42.5 vs. 19.2, $p < 0.01$). Of all potential predictor variables, only the initial PCSS score was independently associated with the odds of symptoms lasting longer than 28 days (aOR 1.037; 95% CI 1.011, 1.063).

Conclusions: Among adult patients with concussions, those with a higher symptom burden after injury have an increased odds of suffering from prolonged symptoms. Other potential predictor variables are not associated with the risk of prolonged recovery.

© 2015 Sports Medicine Australia. Published by Elsevier Ltd. All rights reserved.

1. Introduction

While most concussions, especially those sustained during sports, resolve over the course of a few weeks, some patients suffer prolonged symptoms.^{1–3} Previous studies have attempted to identify potential predictors of prolonged recovery after concussion such as age at time of injury, amnesia or loss of consciousness at the time of injury, sex, history of previous concussions, measures of cognitive function, individual symptoms such as headaches

and dizziness, and collective symptom burden.^{4–6} In a study by Chrisman et al, athletes who reported 4 or more symptoms after sustaining a sport-related concussion were twice as likely to have symptoms for greater than one week. Furthermore, history of prior concussions, amnesia at the time of injury, dizziness, nausea, and difficulties with concentration were all associated with symptom duration of greater than 7 days.⁴ In a similar study by McCrea et al., loss of consciousness, amnesia, and higher symptom severity levels all appeared associated with 7 days or more of symptoms.⁷ Other investigators have found that individual symptoms, such as subjective foginess, dizziness at the time of injury, as well as amnesia and neurocognitive test scores, were also associated with symptom duration.^{7–9}

The ability to predict which patients are likely to suffer prolonged symptoms would allow clinicians to better provide

[☆] This work was funded by National Institutes of Health (NIH) T32 HD040128-06A (Meehan) and the NFL Players Association (Meehan).

* Corresponding author.

E-mail address: William.meehan@childrens.harvard.edu (W.P. Meehan III).

anticipatory guidance and prepare for the possibility of offering various therapies. In addition, the ability to predict a prolonged recovery would allow patients to make better informed decisions about various therapies offered by their treating clinicians, put occupational or academic accommodations in place, and alert co-workers of the possibility of a prolonged recovery.

Previously, we assessed a largely pediatric population and showed that score on symptom inventory at the time of initial clinic visit was independently associated with the odds of suffering the symptoms of a concussion for more than 28 days. It is possible, however, that the recovery of pediatric patients differs from that of adults and that the predictors of prolonged recovery, also differ between children and adults. Age itself has been proposed as a risk factor for worse outcomes after a concussion, including prolonged duration of symptoms, with some analyses suggesting that younger athletes take longer to recover, although not all studies support this idea.^{10–12} Furthermore, there are likely differences between pediatric and adult athletes regarding the biomechanics of concussive injuries, the pathophysiological response to concussion, and the effect of concussion on neurobehavioral outcomes.^{11,13,14}

In this study, we sought to determine if symptom burden at the time of the initial clinic visit was also a predictor of prolonged symptoms for adult patients suffering from concussions. To this end, we analyzed clinical data from 64 adult patients suffering from concussions to identify independent predictors of prolonged (>28 days) symptoms.

2. Methods

We conducted a prospective cohort study of adult (>18yo) patients cared for in a specialty concussion clinic at an academic medical center between October 1, 2009 and July 31, 2011. At the initial clinic visit, patients completed an intake form, providing information about their mechanism of injury and past medical history as well as the Post-Concussion Symptom Scale (PCSS). We used the definition of concussion endorsed by the international consensus on concussion in sport.¹⁵ Therefore, patients with a history of a trauma followed by the onset of signs or symptoms of concussion or decreases in neurocognitive function as determined by medical history and physical examination were diagnosed as a concussion. The study was approved by the institutional review board of Boston Children's Hospital.

We monitored symptoms using the PCSS developed by the international consensus on concussion in sports, which patients completed at each visit. The PCSS is a symptom inventory containing a total of 22 symptoms that patients rank on a scale from 0, when not experiencing a given symptom, to 6, when the symptom is "severe." The PCSS is part of the Sport Concussion Assessment Tool 2 (SCAT 2), which was proposed by the 3rd international consensus on concussion in sport.¹⁵ The total score on the PCSS is the sum of the severity score (0–6) for each of the 22 symptoms. Thus, the maximum possible score on the PCSS is 132 (6 × 22). The symptoms included on the PCSS are not specific to concussion, but can be caused by many other etiologies.^{16,17} Therefore, as with other studies, patients were instructed to rate only those symptoms that started at the time of their concussion and that they were still experiencing within the 24 h prior to their clinic visit.^{18,19} Symptom-free was defined as a post-concussion symptom score of 0. The initial PCSS score used for our analyses was obtained at their first clinic visit.

When assessing potential predictors of symptom duration, we included only those patients seen in clinic within 21 days of injury. We chose to exclude those who presented later than 21 days after injury as (1) we expect that most patients who sustain a concussion will be able to obtain an initial clinic appointment within that time frame and (2) we felt we needed at least one week of

time between presentation to clinic and our outcome of interest, symptom duration for 28 days, in order for potential predictor variables to have an association. We included all sport-related concussions as well as patients who sustained injuries by comparable mechanisms and forces similar to those observed in sports, such as falling from a standing position or fist-fighting. We excluded patients with more severe injury mechanisms and forces, such as motor vehicle collisions and falls from higher than ground level. Simple descriptive statistics were used to describe our population and the sports played at the time of injury. As in previous studies, we identified potential predictor variables from the available medical literature, including age, scores on symptom inventory (PCSS) at initial visit, total number of symptom reported at time of initial visit, sex, loss of consciousness or amnesia at the time of injury, history of prior concussion, prior medical treatments for headaches, history of migraines, and family history of concussion.¹⁹ Migraine headaches were analyzed as a separate category, in addition to headaches not otherwise specified, as some previous literature has suggested headaches with migrainous characteristics may be associated with outcomes after concussion.^{6,20,21} Although computerized neurocognitive data have been suggested as possibly predictive of recovery from concussion, pre- and post-injury tests are not routinely available for adult patients outside of those in the military or those who participate in organized sports. Furthermore, in our sample, only 17 of the 64 patients eligible for analysis of predictors of symptom duration had computerized neurocognitive data performed at their initial visit. Therefore, neurocognitive scores were not included in our analyses.

Since previous studies have shown that the majority of athletes will recover within 4 weeks of injury,^{3,22} we sought to identify risk factors for having symptoms that last longer than 4 weeks. Univariate comparisons were made between the 2 groups: those whose symptoms resolved within 28 days and those whose symptoms persisted beyond 28 days, using Student's *t*-test for continuous variables and the Fischer's exact test for dichotomous variables. Any potential predictor variable that appeared different between the 2 groups with a statistical probability of $p < 0.2$ was entered into a logistic regression model in order to generate adjusted odds ratios. This allowed us to assess the independent effect of symptom burden on duration of symptoms after adjusting for other potential confounding variables. As is common when building a logistic regression model, we used a less stringent threshold for significance when deciding which variables to enter into the multivariate model, since the true size and significance of the effect of certain variables may be masked by potential confounding variables, and revealed after adjusting for those variables in the model.²³ No inferences are drawn from these univariate analyses; they are simply a screening to determine which independent variables are included in the model. Any variable with an adjusted odds ratio with a 95% confidence interval that did not cross 1 in the multivariate model was considered statistically significant.

3. Results

There were 131 adult patients cared for in clinic during the study period. The initial visit for 60 patients occurred more than 21 days after injury. For an additional 7, it could not be definitively established whether or not their symptoms resolved within 28 days of injury. Therefore, 64 patients, none of whom were lost to follow up, were included in the study.

The majority (53.3%) were male. More than 1 in 5 (20.3%) patients reported experiencing a loss of consciousness at the time of injury while 23.4% reported amnesia; 6.3% experienced both amnesia and a loss of consciousness. Most concussions (92.2%) occurred during sports. The most common sports played at the time of injury were soccer and ice hockey (Table 1). Nearly half (46.9%) had been

Download English Version:

<https://daneshyari.com/en/article/2703757>

Download Persian Version:

<https://daneshyari.com/article/2703757>

[Daneshyari.com](https://daneshyari.com)