

Subject Index

- Acute ischemic stroke, *See* Ischemic stroke, acute
- Adenosine diphosphate
and collagen-induced platelet aggregation, in patients treated with ticlopidine and/or aspirin, recurrent cerebral infarction with, 319–323
- Adjuvant therapy
donepezil, to standard medical care, enhancing recovery after acute ischemic stroke with, 177–182
- ADMA, *See* Asymmetric dimethylarginine
- Adults
young
factors associated with misdiagnosis of acute stroke in, 523–527
in Thailand, causes of ischemic stroke in, 247–250
- Alendronate
an open-label trial comparing, and alphacalcidol in reducing falls and hip fractures in disabled stroke patients, 41–46
- Alphacalcidol
an open-label trial comparing, and alendronate in reducing falls and hip fractures in disabled stroke patients, 41–46
- Alteplase
0.6 mg/kg, factors predicting outcome in stroke patients treated with, 517–522
- Americans
with history of stroke, subpar utilization of dental care among, 255–259
- Aneurysm
cerebral, classification based on angioarchitecture, 162–167
of common carotid artery, revealing Takayasu's arteritis, 556–558
intracranial, and middle cerebral artery, recanalization in same ischemic territory with intravenous administration of rt-PA, 269–272
ruptured, on dolichoectatic middle cerebral artery, hyperperfusion syndrome after neck clipping of, 260–263
subarachnoid hemorrhage, life satisfaction, and return to work after, 324–329
- Animal models
effect of combined treatment with curcumin and candesartan on ischemic brain damage in, 541–548
phosphorylated eNOS Ser1177 via the PI3K/Akt pathway is depressed in the brain of stroke-prone spontaneously hypertensive, 406–412
- Antiplatelet agents
are risk factors for cerebellar hemorrhage in patients with primary intracerebral hemorrhage, 346–351
- Aortic arch disease
comparison of CTA and TEE for evaluation of, 436–442
- Aortic dissection
intravenous thrombolysis in an elderly patient with acute ischemic stroke masking, 559–561
before thrombolysis in patients with ischemic stroke has been insufficiently advised, 384 (letter), 385 (reply)
- Aphasia
poststroke, recovery assessed with functional MRI and picture identification task, 336–345
- Aspirin
and/or ticlopidine, recurrent cerebral infarction with adenosine diphosphate- and collagen-induced platelet aggregation in patients treated with, 319–323
and clopidogrel, biochemical resistance in patients with ischemic stroke, predictors of, 227–230
- Asymmetric dimethylarginine (ADMA)
in patients in chronic phase of cerebral infarction, effects of EPA on, 474–478
- Basilar artery
MRI to examine the cerebral microbleeds and dilative remodeling of, 429–435
- Bicuspid aortic valve
mildly stenosis, recurrent posterior circulatory emboli from, 562–564
- Blood pressure
1 year after stroke, 16–23
relationship between dipping profile in, and neurologic deficit in early acute ischemic stroke, 10–15
- Brachiocephalic vein occlusion
left, and peripheral hemodialysis shunt, venous cerebral infarction in patient with, 381–383
- Brain ischemia
in mice, effect of combined treatment with curcumin and candesartan on, 541–548
- Brain microbleeds
topographical distribution with particular emphasis on diffuse microbleeds, 214–221
- Candesartan
and curcumin, on ischemic brain damage in mice, effect of combined treatment with, 541–548
- Cardiac papillary fibroelastoma
a source of cerebral embolism treated with intravenous thrombolysis, 485–487
- Carotid artery
common, aneurysm of, revealing Takayasu's arteritis, 556–558
duplex ultrasonography of, predicts outcome in intravenous alteplase therapy for hyperacute stroke, 24–29
external, branches, preoperative embolization as adjunct to superficial temporal artery-to-middle cerebral artery bypass, 168–171
internal, stenosis of, clinical characteristics and outcomes in carotid endarterectomy for, in a Japanese population, 55–61
- Carotid endarterectomy
clinical characteristics and outcomes in, for internal carotid artery stenosis in a Japanese population, 55–61
- Carotid intima-media thickness
in Framingham Offspring Study, segment-specific association between plasma homocysteine level and, 155–161
- Case reports
acute ischemic stroke lasting for 5 days without definite lesions on diffusion-weighted MRI in a patient with severe leukoariosis, 482–484
cardiac papillary fibroelastoma, a source of cerebral embolism treated with intravenous thrombolysis, 485–487
cerebral embolism with metastatic chondrosarcoma in left atrium, 79–81
cerebral infarction in Duchenne muscular dystrophy, 264–265
cerebral thrombosis complicating Crohn's disease: two cases, 565–569

- common carotid artery aneurysm revealing Takayasu's arteritis, 556–558
- hemodynamic assessment of cervical internal carotid artery stenosis during head manipulation, 479–481
- hepatic encephalopathy with reversible focal neurologic signs resembling acute stroke, 377–380
- hyperperfusion syndrome after neck clipping of ruptured aneurysm on dolichoectatic middle cerebral artery, 260–263
- imaging studies conflict with neurologic exam, 82–84
- Intravenous thrombolysis in an elderly patient with acute ischemic stroke masking aortic dissection, 559–561
- moyamoya syndrome associated with Graves' disease: a case report and review of the literature, 528–536
- preoperative embolization of the external carotid artery branches as an adjunct to superficial temporal artery-to-middle cerebral artery bypass, 168–171
- recanalization of middle cerebral artery and intracranial aneurysm in same ischemic territory with intravenous administration of recombinant tissue plasminogen activator, 269–272
- recurrent posterior circulatory emboli from a mildly stenosed bicuspid aortic valve, 562–564
- recurrent unconsciousness due to bilateral middle cerebral artery stenosis, 266–268
- venous cerebral infarction in a patient with peripheral hemodialysis shunt and occlusion of the left brachiocephalic vein, 381–383
- Cerebellar hemorrhage
 - in patients with primary intracerebral hemorrhage, antiplatelet agents are risk factors for, 346–351
- Cerebral autosomal recessive arteriopathy with subcortical infarcts and leukoencephalopathy (CARASIL)
 - from discovery to gene identification, 85–93
- Cerebral embolism
 - with metastatic chondrosarcoma in left atrium, 79–81
 - treated with intravenous thrombolysis, cardiac papillary fibroelastoma, a source of, 485–487
- Cerebral infarction
 - chronic-stage, platelet aggregates detected by conventional hematology analyzer method is a risk factor for stroke or predictive factor in patients with, 275–281
 - in Duchenne muscular dystrophy, 264–265
 - effects of EPA on ADMA in patients in chronic phase of, 474–478
 - hyperacute, significance of clinical–diffusion mismatch in, 62–67
 - recurrent, with adenosine diphosphate- and collagen-induced platelet aggregation in patients treated with ticlopidine and/or aspirin, 319–323
 - venous, in patient with peripheral hemodialysis shunt and occlusion of left brachiocephalic vein, 381–383
- Cerebral microbleeds
 - clinical correlating factors and cognitive function in community-dwelling healthy subjects with, 105–110
 - and dilative remodeling of the basilar artery, MRI to examine the, 429–435
 - in healthy subjects without major cerebrovascular risk factors, age-related changes in, 302–309
- Cerebrolysin
 - in vascular dementia, 310–318
- Cerebrospinal fluid (CSF)
 - ferritin levels, a sensitive diagnostic test in late-presenting subarachnoid hemorrhage, 489–493
- Cerebrovascular disease
 - ischemic, first, TEE screening in subjects with, 503–509
 - risk factors of, age-related changes in white matter lesions, hippocampal atrophy, and cerebral microbleeds in healthy subjects without, 302–309
- Chinese
 - risk factors and prevention of stroke in, 395–400
- Chondrosarcoma
 - metastatic, cerebral embolism with, in left atrium, 79–81
- Clinical–diffusion mismatch
 - significance of, in hyperacute cerebral infarction, 62–67
- Clopidogrel
 - and aspirin, biochemical resistance in patients with ischemic stroke, predictors of, 227–230
- Cocaine
 - related ischemic stroke, clinical profiles, complications, and disability in, 443–449
- Cognitive function
 - in acute phase poststroke, 549–555
 - and clinical correlating factors, in community-dwelling healthy subjects with cerebral microbleeds, 105–110
- Cognitive impairment
 - at 1 year poststroke, relationships between quality of life and, 450–458
- Collagen
 - and adenosine diphosphate, induced platelet aggregation in patients treated with ticlopidine and/or aspirin, recurrent cerebral infarction with, 319–323
- Community-based stroke care
 - in Japan, factors influencing cooperation among healthcare providers in, 413–423
- Computed tomography angiography (CTA)
 - versus TEE, for evaluation of aortic arch disease, 436–442
- Conventional hematology analyzer
 - is a risk factor for stroke or predictive factor in patients with chronic-stage cerebral infarction, platelet aggregates detected by, 275–281
- COOP/WONCA charts
 - for stroke patients in primary care, reliability and validity of, 465–473
- Copper-zinc
 - kinetics of, and manganese superoxide dismutase activity with acute ischemic stroke, 75–78
- Cortical infarction
 - small, risk factors for, on diffusion-weighted magnetic resonance imaging in patients with acute ischemic stroke, 68–74
- COX-2, *See* Cyclooxygenase-2
- Craniotomy
 - and neuroendoscopic surgery, for spontaneous ICH, comparison of clinical outcomes of, 208–213
- Crohn's disease
 - cerebral thrombosis complicating, 565–569
- CSF, *See* Cerebrospinal fluid
- CTA, *See* Computed tomography angiography
- Curcumin
 - and candesartan, on ischemic brain damage in mice, effect of combined treatment with, 541–548
- Cyclooxygenase-2
 - rs5275 and rs20417, polymorphisms of, impact on 90-day functional outcome in patients with ischemic stroke, 134–144
- Dementia
 - vascular, cerebrolysin in, 310–318
- Dental care
 - among Americans with history of stroke, subpar utilization of, 255–259
- Diagnostic testing
 - sensitive, of CSF ferritin levels in late-presenting subarachnoid hemorrhage, 489–493
- Diffuse microbleeds
 - topographical distribution of brain microbleeds with particular emphasis on, 214–221

Download English Version:

<https://daneshyari.com/en/article/2706580>

Download Persian Version:

<https://daneshyari.com/article/2706580>

[Daneshyari.com](https://daneshyari.com)