

EPIDEMIOLOGICAL ANALYSIS ON ACETABULAR FRACTURES

Maurício Silveira Maia¹, Denise Cristina Montecchio Santos², Daniel Magalhães de Queiroga³, Claydson de Oliveira Castro², Rebeca Macedo Fraga e Silva², Aliny Cristine Brito Reis², Aline Cristina Ducatti⁴

ABSTRACT

This aim of this work was to carry out an epidemiological study on acetabular fractures in the city of Campinas and surrounds, in view of the few published papers on this subject. Medical files with a diagnosis of acetabular fracture between the years 2004 and 2008 that were made available by the Medical Archiving Service of Hospital das Clínicas, State University of Campinas (UNICAMP) were analyzed by six observers. Data on patients' ages, sex, side affected by the fracture, mechanism of injury, material used for syn-

thesis, complications of the operation, associated fractures, length of hospitalization before and after the surgery, time of total internment and number of physiotherapy sessions before and after the surgery were gathered. It was observed in this population that the left side was more affected; the mechanism of injury that most often caused this type of fracture was automobile accidents; injuries to the sciatic nerve were the commonest surgical complications; and the synthesis material most used was reconstruction plates.

Keywords – Epidemiology; Acetabulum; Bone Fractures

INTRODUCTION

Fractures of the acetabulum are generally very serious, since they are associated with high speed and high-energy trauma, such as in vehicle accidents, and often occur in multiple trauma patients. Over the years, treatment of this type of fracture has been a challenge to surgeons because they are located in a region that is difficult to access; there is a lack of reduction techniques; few implants are available for fracture fixation; and the patients' clinical condition is highly complex. Hence, conservative treatment is considered

to be the preferred approach, because it is safer.

The techniques for treating acetabular fractures have become more refined through the studies of Judet and Letournel, which have contributed towards surgical advances and have provided for physiotherapeutic action for these patients' rehabilitation. This has enabled early mobilization of this joint after the surgery. Implementing physiotherapy both before and after the operation is of fundamental importance for these patients' respiratory and motor functions, thereby making early hospital discharge possible⁽¹⁾.

1 – Physiotherapist; Specialist in Physiotherapy for Orthopedics and Traumatology from Campinas State University (UNICAMP); Specialist in Sports Physiotherapy from the Federal University of São Paulo (UNIFESP). Postgraduate Specialization Student of Physiotherapy for Orthopedics and Traumatology, State University of Campinas.

2 – Postgraduate Specialization Student of Physiotherapy for Orthopedics and Traumatology, State University of Campinas.

3 – Physiotherapist at CLIFIM Clinic, Jaguariúna, SP. Postgraduate Specialization Student of Physiotherapy for Orthopedics and Traumatology, State University of Campinas.

4 – Physiotherapist at the Municipal Authority of Uberlândia, MG. Postgraduate Specialization Student of Physiotherapy for Orthopedics and Traumatology, State University of Campinas.

Work performed at Hospital das Clínicas, State University of Campinas (UNICAMP).

Correspondence: Rua Irarié Quadra 07 - lote 06 - Parque Acalanto - 74860-140 - Goiânia, GO.

E-mail: mauriciosilveiramaia@gmail.com

Work received for publication: December 4, 2009; accepted for publication: July 19, 2010.

The aim of this study was to make an epidemiological analysis on cases of acetabular fracture attended at Hospital das Clínicas, State University of Campinas (Unicamp) between 2004 and 2008.

METHODOLOGY

A retrospective study was conducted in the city of Campinas, State of São Paulo, among male and female patients hospitalized in Hospital das Clínicas, State University of Campinas (Unicamp), with a diagnosis of acetabular fracture between 2004 and 2008.

For the purposes of this study, data were gathered between December 1 and 20, 2008, through analysis on medical files held at the medical archiving service. Six observers gathered data on patients' ages, sex, side affected by the fracture, injury mechanism, synthesis material used, complications during the operation, associated fractures, length of hospital stay before and after the operation, total duration of hospital stay and numbers of physiotherapy sessions undertaken before and after the surgery.

To analyze the data, the Microsoft Office Excel 2007 software was used. The data were expressed as means and standard deviations, and were then correlated using percentages and graphical comparisons.

RESULTS

The medical files of 69 patients were evaluated. Seventy-one cases of acetabular fracture were found and, among these, 85.5% were in men and only 14.5% were in women (Figure 1). The mean age among the population analyzed was 33 years, with a range from 16 to 66 years.

The side that was more commonly affected by a fracture was the left side, such that 57.7% (41) of the patients presented a fracture in the left-side ace-




Figure 1 – Correlation between acetabular fractures and gender.

tabulum, while 42.3% (30) presented a fracture in the right-side acetabulum (Figure 2). Occurrences of hospitalization at Hospital das Clínicas due to acetabular fractures increased over the years (Figure 3). There was no difference in prevalence between the months of the year.

The injury mechanism that was most responsible for acetabular fractures was car accidents, which accounted for 46.37% (32), followed by motorcycle accidents (31.88%; 22) and falls (8.69%; 6) (Figure 4)^(3,4).

The mean duration of the hospital stay before the operation was 9.7 days, and the mean hospital stay after the operation was 11.42 days.




Figure 2 – Correlation between acetabular fractures and the affected side.




Figure 3 – Incidence of acetabular fractures per year.

Download English Version:

<https://daneshyari.com/en/article/2708433>

Download Persian Version:

<https://daneshyari.com/article/2708433>

[Daneshyari.com](https://daneshyari.com)