



## Research Paper

# Thumb function and appearance following treatment of Wassel type III duplication thumbs

Haifeng Zhang, Zijing Du, Hao Jiang, Dong Han<sup>\*</sup>, Qingfeng Li

Department of Plastic and Reconstructive Surgery, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China

## ARTICLE INFO

## Article history:

Received 6 May 2016

Received in revised form 16 May 2016

Accepted 18 May 2016

Available online 20 May 2016

## Keywords:

Duplicated thumbs

Wassel type III

Polydactyly

Nail

## ABSTRACT

**Objective:** The purpose of our study is to evaluate thumb function and appearance after surgical correction of Wassel type III thumbs polydactyly.

**Methods:** We have reconstructed 28 cases of Wassel type III duplication thumbs, in which the duplicated digits were equal or almost equal in size by ablation of a radial digit. The extra thumb is osteotomized at the bifurcation level and excised except for the distal bone fragment supporting the nail bed and fillet flap. Meanwhile, the nail of the retained thumb should be reserved completely, and if the nail has relatively poor appearance it should be repaired by nail lengthening surgery. Eighteen cases were followed up for more than 3 years and were available for assessment using the Japanese Society for Surgery of the Hand evaluation form. The average age at follow-up was 5 years. The size of the nail and distal phalanx was measured to assess the growth of the thumb.

**Results:** An average functional point was 12 points (maximum 14 points) and the cosmetic score averaged 3.6 (maximum 4 points) after the assessment. Slightly small nails without a central ridge were deemed acceptable. Second revision surgery is seldom. Long-term results after surgical reconstruction for duplication thumbs were excellent, and all patients and parents were satisfied with the cosmetic and functional results.

**Conclusions:** This procedure is a helpful and effective way to provide functional and aesthetic thumb for Wassel type III duplication thumbs.

© 2016 The Authors. Published by Elsevier Ltd on behalf of Surgical Associates Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Thumb polydactyly is a relatively common congenital hand abnormality. In 1969, Wassel provided a simple, effective classification system for the different phenotypic variations in radial polydactyly based on the level of skeleton duplication [1]. Wassel type III duplication thumbs occupies a certain proportion in it and is one of the difficult types for reconstruction surgery, especially in these duplication thumbs that are equal or almost equal in size.

The aim of surgical intervention that addresses duplication thumbs is to obtain a stable, mobile thumb of appropriate shape and adequate size. Dijkman et al. recommended a reliable assessment system—the Japanese Society for Surgery of the Hand (JSSH)—to evaluate functional and aesthetic outcomes after surgery for radial polydactyly by comparing several assessment systems [2]. Based on the Wassel classification and JSSH assessment system, the purpose

of our study is to introduce our technique that can maintain stability of the IP joints and appearance with acceptable results.


## 2. Methods

### 2.1. Surgical technique

Under general anesthesia, a new clinical evaluation, including the size of the nails of the two thumbs, and the passive mobility of the metacarpophalangeal (MCP) and the interphalangeal (IP) joints, was measured by one of the authors. After a zigzag skin incision and subcutaneous dissection (Fig. 1A), we removed the entire nail of the radial thumb and delimited an ungual soft tissue flap harvested from the radial edge. The nail of the ulnar thumb duplicate was therefore totally preserved. If the nail of one of the thumbs is more than 70% of the contralateral normal, thumb nail width, it is retained in its entirety with excision of the other nail [3]. When the nail width is less than 70% of the contralateral normal, a nail lengthening surgery is needed [4]. Depending on the length of the contralateral fingernail, we designed a small rectangle excision whose size ranges between 2 and 3 mm, and the distance from the distal border of the eponychium is about 5 mm (Fig. 1A). At the level of subcutaneous vascular network,

<sup>\*</sup> Corresponding author. Department of Plastic and Reconstructive Surgery, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, No. 639, Zhizaoju Road, Huangpu District, Shanghai 200011, China. Tel.: +86 (0) 021 23271699, fax: +86 (0) 021 63087768.

E-mail address: [handong12000@163.com](mailto:handong12000@163.com) (D. Han).



**Fig. 1.** Schematic depiction of surgical techniques. (A) A zigzag incision on the thumb and a small rectangle excision in nail lengthening surgery if the nail width is less than 70% of the contralateral normal. (B) A wedge osteotomy is performed to correct the angular deformity and the distal bone fragment from supernumerary thumb is supporting the nail bed and making the distal phalange bigger and wider. (C) Skin closure was performed and the Kirschner wire fixed in osteotomy site. (D) Skin closure on the volar.

the rectangle site was de-epithelialized. After a small retractor separated the eponychium from the nail matrix, we could slide the eponychial flap proximally and suture two ends by PDS® 6.0.

The radial ligament and periosteum from supernumerary thumb should be spread apart gently but ended in the distal bone fragment supporting the nail bed and fillet flap to augment the retained ulnar duplicate. The extra thumb is separated and osteotomized at the bifurcation level and the distal bone fragment attached ligament and periosteum is fixed to retained thumb bone whose distal radial bone piece was excised and then a suitable Kirschner wire should be used to fix them. When the angular deformity of the interphalangeal joint is  $>30^\circ$  [5,6], a corrective closing wedge osteotomy is performed at the proximal phalanx of the retained thumb, and the osteotomy site is held by advancing the transarticular Kirschner wire (Fig. 1B). If the angle of the IP joint was greater than  $20^\circ$  but less than  $30^\circ$ , the radial collateral ligament restoration was done, as a small amount of angulation at the IP joint can be corrected by transferring the soft tissues from the radial digit.

We then proceeded to suture the periosteum to the bone using PDS® 5.0 sutures. We concluded with reinsertion of the extensor apparatus and the flexor pollicis longus tendon with stitches at the metaphysis. Skin closure of the radial flap was performed with separate stitches using PDS® 6.0 (Fig. 1C and 1D). The thumb was immobilized and protected within a light plaster support for 4 weeks after surgery. All procedures were performed by the same surgeon and two assistants.

## 2.2. Demographics

Between 2010 and 2012, twenty eight patients presenting type III thumb duplication were operated in our department (Fig. 2A). There was one case who had a family history of thumb polydactyly in first-degree relatives. Five cases with extra thumb obviously

smaller than the other were treated by an ablation of the smaller digit. And four cases did not follow up over 3 years. The above 10 cases were excluded from this study. The remaining 18 cases included 12 duplication of the right thumb and 6 duplications of the left. The median age at surgery was 12 months (range: 6 months–35 years) including two adults surgeries. Ten were male. In all the cases, patients should have 2 digits with equal or almost equal size and anteroposterior x-rays of the thumb were done.

The 18 patients operated with this technique were followed up for a mean of 47 months (range: 36–60 months). The clinical results were assessed functionally and aesthetically using an evaluation form for duplication thumbs provided by the Japanese Society for Surgery of the Hand (Table 1). Functional assessment outcomes, including abnormal alignment, instability, active flexion, and extension lag at the level of the IP and MCP joints, as well as palmar abduction at the level of MCP and CMC joints, were measured and scored. Although aesthetic assessment is subjective, it is also an important measure of the success of the procedure. Cosmetic evaluation was done in terms of surgical scar and nail deformity. The maximum width and the length of the fingernail were measured and compared to the other side. The result was considered excellent if the score was 20, good if the score was greater than 16 points, fair if the score was between 14 and 16 points, and poor if the score was less than 13 points. The study was approved by our hospital and oral patient consent was obtained.

## 3. Results

There were no early postsurgical complications in all cases (Fig. 2B). A detailed outcome is shown in Table 2. There was no secondary surgery in 18 patients until now. In six cases, the oblique wedge osteotomy of the proximal phalanx had been done. Fifteen cases had relocated surgery of flexor pollicis longus (FPL) tendon

Download English Version:

<https://daneshyari.com/en/article/2728496>

Download Persian Version:

<https://daneshyari.com/article/2728496>

[Daneshyari.com](https://daneshyari.com)