



REVISTA BRASILEIRA DE ANESTESIOLOGIA

Official Publication of the Brazilian Society of Anesthesiology
www.sba.com.br



CLINICAL INFORMATION

Intraoperative “Kounis syndrome” that improved electrocardiography changes and hemodynamic situation after administering nitroglycerine

Victoria O. Sánchez*, Luisa C. Roca, Angel del P. Moreno

Hospital Virgen del Castillo, Yecla, Spain

Received 11 April 2013; accepted 10 June 2013

Available online 23 October 2013

KEYWORDS

Kounis syndrome;
Cefazolin;
Rocuronium

Abstract

A 58-year-old female without cardiovascular risk factors, was going to be operated to repair the rotator cuff. Induction and interscalene brachial plexus block were uneventful, but after her placement for surgery the patient started with severe bronchospasm, hypotension, cutaneous allergic reaction and ST elevation on the electrocardiogram. An anaphylactic shock was suspected and treated but until the perfusion of nitroglycerine was started no electrocardiographic changes resolved. After necessary diagnostic test the final diagnosis was variant I of Kounis syndrome due to cefazolin and rocuronium. Ephinephrine is the cornerstone of treatment for anaphylaxis but should we use it if the anaphylactic reaction is also accompanied by myocardial ischemia? The answer is that we should not use it because myocardial ischemia in this syndrome is caused by vasospasm, so it would be more useful drugs such as nitroglycerin. But what if we do not know if it is a Kounis syndrome or not? In this article we report our experience that maybe could help you in a similar situation.

© 2013 Sociedade Brasileira de Anestesiologia. Published by Elsevier Editora Ltda.

Este é um artigo Open Access sob a licença de [CC BY-NC-ND](http://creativecommons.org/licenses/by-nc-nd/4.0/)

PALAVRAS-CHAVE

Síndrome de Kounis;
Cefazolina;
Rocurônio

Síndrome de Kounis intraoperatória com melhoria das alterações eletrocardiográficas e da situação hemodinâmica após a administração de nitroglicerina

Resumo

Paciente do sexo feminino, 58 anos, sem fator de risco cardiovascular, submetida a cirurgia para reparação do manguito rotador. A indução do bloqueio do plexo braquial interesca-lênico foi feita sem intercorrência, mas, após seu posicionamento para a cirurgia, a paciente apresentou broncoespasmo grave, hipotensão, reação alérgica cutânea e elevação do segmento ST ao eletrocardiograma. Houve suspeita de choque anafilático que foi tratado, mas até que a perfusão de nitroglicerina fosse iniciada não houve resolução das alterações eletrocardiográficas. Após teste diagnóstico necessário, o diagnóstico final foi de variante tipo I

* Corresponding author.

E-mail: alsuelovas@hotmail.com (V.O. Sánchez).

da síndrome de Kounis por causa de cefazolina e rocurônio. Epinefrina é a base sólida do tratamento para anafilaxia, mas devemos usá-la se a reação anafilática também for acompanhada de isquemia miocárdica? A resposta é que não devemos usá-la, porque a isquemia miocárdica nessa síndrome é causada por vasoespasmo; portanto, drogas como a nitroglicerina seriam mais úteis. Porém, e quando não sabemos se é ou não uma síndrome de Kounis? Neste artigo relatamos nossa experiência que, talvez, possa ajudar em uma situação similar.

© 2013 Sociedade Brasileira de Anestesiologia. Publicado por Elsevier Editora Ltda. Todos os direitos reservados.

Introduction

Kounis syndrome was first described in 1991 by Nicholas Kounis and Zavras in the study: "Histamine-induced coronary artery spasm: the concept of allergic angina".¹

Kounis syndrome is characterized by the simultaneous appearance of an allergic reaction and an acute coronary syndrome. It is also known as allergic angina or allergic myocardial infarction.² Kounis syndrome is classified into two groups:

Group I: patients without cardiovascular risk factors and nor known coronary artery disease.

Group II: patients with cardiovascular risk factors with or without associated coronary artery disease.

Recently a Group III²⁻⁴ has been proposed which would belong to patients with drug-eluting stents that are complicated by thrombosis and in which the histology observed infiltrates of mast cells and eosinophils.

This unique syndrome is under-diagnosed because it requires an electrocardiography recording during the allergic reaction, normally prioritizing the treatment of anaphylaxis, and when we finally do the electrocardiogram it is no longer altered. Moreover, many of these reactions occur on the street and when patients reach the health center their electrocardiograms are normal, because the electrical changes tend to disappear in a brief period of time.

The question arises at the moment of using epinephrine. Epinephrine is the cornerstone of treatment for anaphylactic shock, but we should think twice before using it in Kounis syndrome because it could aggravate cardiac ischemia produced by vasospasm.

This doubt also occurs with other drugs like aspirin, indicated in the treatment of acute coronary syndrome but which in turn may cause anaphylactic reactions, or in the case of nitro-glycerine that is also used in acute coronary syndrome, but could worsen the hypotension and tachycardia caused by anaphylaxis.⁵

We report a case of intraoperative Kounis syndrome that was resolved mainly due to the administration of nitroglycerine.

Case report

A 58-year-old woman and 65 kg with hypercholesterolemia in dietary treatment and operated of fibro adenoma breast and tonsils. She had no personal or family history of allergy.

The patient was going to be operated to arthroscopically repair the rotator cuff. In the operating room she was monitored by pulse oximetry, non-invasive blood pressure and electrocardiogram. She was hemodynamically stable with a blood pressure of 190/90 mmHg and was anxious.

As premedication she received 2 g cefazolin, 50 mg ranitidine, and 4 mg of ondansetron. In the operating room she was administered midazolam 2 mg, propofol 10 mg and dexamethasone 4 mg.

She was performed an interscalene brachial plexus block with ropivacaine 0.5% 20 mL without complications. It was proceeded to induct fentanyl 300 µg; propofol 100 mg; atropine 0.5 mg and 50 mg rocuronium. She was intubated by orotracheal via and maintained with inhalation anesthesia with sevoflurane. She was administered preemptive analgesia with paracetamol 1 g and dexketoprofen 50 mg and placed in the lateral decubitus position for surgery.

Traumatologists made cutaneous incision use to infiltrate lidocaine 4 mg with epinephrine 25 µg. Suddenly we observed an increase in heart rate of 140 bpm, followed by severe bronchospasm with difficulty in mechanical ventilation, with pressures up to 40 cmH₂O in the airway. Cardiopulmonary auscultation was tachycardia and hypoventilation with decrease in vesicular murmur. Subsequently, end tidal carbon dioxide descended to 9 mmHg, and oxygen saturation to 75%. Finally we observed the ST elevation in leads II and III on the monitor. Until that moment measurement of blood pressure was programmed every 5 min so, in this instant marked 120/70 mmHg.

Surgery was interrupted immediately, the patient was placed in decubitus supine, we started manual ventilation with FiO₂ of 1. Then the measurement of blood pressure revealed a hypotension of 63/39 mmHg, and we stopped the administration of sevoflurane and started with intensive fluid therapy. When we removed the cloth from the surgical field we observed generalized skin rash. It was made a twelve-lead electrocardiogram in which we could see ST segment elevation in the front and bottom of the heart (Fig. 1).

An emergent consultation to cardiologist recommended implementing the hospital protocol of acute coronary syndrome, with perfusion of nitroglycerine at titrated doses, depending on the patient's hemodynamic situation, clopidogrel 300 mg and acetylsalicylic acid 100 mg by nasogastric tube.

Suspecting an anaphylactic reaction, we administered methylprednisolone 80 mg IV, dexchlorpheniramine maleate 5 mg IV and ranitidine 50 mg IV. At first we were not sure whether to use adrenaline or not, but finally it was decided not to administer it given the possibility of the patient

Download English Version:

<https://daneshyari.com/en/article/2750300>

Download Persian Version:

<https://daneshyari.com/article/2750300>

[Daneshyari.com](https://daneshyari.com)