



Original Contribution

Single-injection thoracic paravertebral block and postoperative analgesia after mastectomy: a retrospective cohort study ☆, ☆ ☆, ★



Rishi R. Agarwal MD (Resident)^a, Anne M. Wallace MD (Professor)^b,
Sarah J. Madison MD (Assistant Clinical Professor)^a,
Anya C. Morgan MA, CCRC (Research Coordinator)^a,
Edward J. Mascha PhD (Associate Staff Biostatistician)^c,
Brian M. Ilfeld MD, MS (Professor, In Residence)^{a,*}

^aDepartment of Anesthesiology, University of California San Diego, San Diego, CA

^bDepartment of Surgery, University of California San Diego, San Diego, CA

^cDepartments of Quantitative Health Sciences and Outcomes Research, Cleveland Clinic, Cleveland, OH

Received 2 October 2014; revised 31 March 2015; accepted 16 April 2015

Keywords:

Postoperative analgesia;
Postoperative pain;
Peripheral nerve block;
Regional anesthesia;
Regional analgesia

Abstract

Background: The treatment of postoperative pain after mastectomy is an area of increasing interest, as this treatment option is now considered a standard of care for those affected by breast cancer. Thoracic paravertebral nerve block (tPVB) using local anesthetics administered before mastectomy can theoretically provide postoperative analgesia, thereby facilitating a more comfortable and shorter hospitalization.

Methods: In this retrospective cohort study, we aimed to determine the duration and degree to which tPVB provides postoperative analgesia in patients who underwent either unilateral or bilateral mastectomy (n = 182). We retrospectively examined the numeric rating scale (NRS) for pain scores recorded by nursing staff throughout individual patient hospitalizations, looking specifically at the following time points: arrival from the postanesthesia care unit to the surgical wards, noon on postoperative day 1 (POD1), and discharge. We also examined the number of days until patients were discharged from the hospital.

Results: Our results revealed a statistically significant decrease in NRS in pain scores for patients who had received a tPVB (n = 92) on arrival from the postanesthesia care unit to the surgical wards (mean NRS decrease of 1.9 points; 99% confidence interval [CI], -3.0 to -0.8; $P < .001$) but did not show statistically significant decreases in NRS for pain scores for patients at noon on POD1 (mean NRS decrease of 0.3 points at noon on POD1, $P = .43$) or at discharge (mean NRS decrease of 0.1 point at discharge, $P = .65$). Moreover, use of tPVB did not have an impact on time until discharge (average decrease of 0.5 hours; 95% CI, -6 to +5 hours, $P = .87$).

☆ Financial support: The Department of Anesthesiology, University of California San Diego (San Diego, CA). The contents of this article are solely the responsibility of the authors and do not necessarily represent the official views of the funding entity.

☆☆ Conflict of interest: none.

★ Presented, in part, as a scientific abstract at the annual meeting of the American Society of Anesthesiologists in New Orleans, LA, October 11-15, 2014.

* Corresponding author at: Department of Anesthesiology, 200 West Arbor Drive, MC 8770, San Diego, CA, 92103-8770. Tel.: +1 858 481 8454; fax: +1 858 683 2003.

E-mail address: bilfeld@ucsd.edu (B.M. Ilfeld).

<http://dx.doi.org/10.1016/j.jclinane.2015.04.003>

0952-8180/© 2015 Elsevier Inc. All rights reserved.

Conclusions: Single-injection tPVB appears to provide meaningful postoperative analgesia in the immediate postoperative period after mastectomy but not after the first day of surgery.
© 2015 Elsevier Inc. All rights reserved.

1. Background

The prevalence of breast cancer in women within the United States is more than 3 million, with an annual incidence of approximately 100,000 and more than 35,000 women undergoing mastectomy annually [1,2]. Pain after mastectomy is often severe [3] and described as a “continuous aching pain” the day after surgery [4]. As such, adequate pain control is a primary concern for both patients and health care providers. In addition, postoperative opioid analgesics often induce nausea and vomiting, increasing patient discomfort as well as postanesthesia recovery room and hospitalization durations.

In contrast, there is evidence that a single-injection thoracic paravertebral nerve block (tPVB) may provide potent and safe intraoperative and postoperative analgesia [4,5]. A tPVB involves the percutaneous injection of long-acting local anesthetic adjacent to the peripheral nerves that innervate the breast immediately lateral to the upper thoracic vertebra, either unilaterally or bilaterally. The aim of this retrospective cohort study was to determine if—and to what degree—a single-injection ropivacaine tPVB provides postoperative analgesia after unilateral and bilateral mastectomy.

2. Methods

After institutional review board (University of California San Diego, San Diego, CA) approval, we analyzed the records of 182 patients who underwent either unilateral or bilateral mastectomy with a single surgeon at the University of California San Diego between the years 2009 and 2011. Before 2010, patients undergoing mastectomy received an opioid-based analgesic regimen. During 2010, the Regional Anesthesia and Acute Pain Medicine Division began providing tPVBs for patients having mastectomy. All patients received oral and intravenous opioids as well as oral acetaminophen for analgesia, as necessary.

For subjects who had received a single-injection tPVB: with a low-frequency (5-2 MHz) curved array transducer (C60x; SonoSite MicroMaxx, Bothell, WA), the paravertebral space between the third and fourth thoracic vertebrae was identified in a parasagittal view approximately 3 cm lateral to midline on the side of surgery by a regional anesthesiology attending or fellow. Following sterile preparation, local anesthetic skin wheal was raised caudal to the ultrasound transducer. An 8.9-cm, 17-gauge, Tuohy-tip needle (Teleflex Medical/Arrow International, Research Triangle Park, NC)

was inserted through the skin wheal in plane beneath the ultrasound transducer and directed to the paravertebral space. Ropivacaine 0.5% with epinephrine 5 µg/mL (20 mL) was slowly injected with gentle aspiration every 3 mL.

We hypothesized that use of a single-injection tPVB would be associated with lower pain scores during hospitalization after mastectomy from arrival on the postsurgical ward through noon the day after surgery (postoperative day 1 [POD1]). Pain was recorded by nursing staff using a 0-10 Likert numeric rating scale (NRS) for pain (0, no pain; 10, worst pain imaginable pain). The primary outcome measures of interest included the NRS (1) at the time of arrival to the surgical wards, (2) at noon on POD1, (3) the minimum NRS between those 2 time points; (4) the maximum NRS between those 2 time points, and (5) the NRS at discharge. A secondary outcome measure was the total number of hospitalization days.

2.1. Statistical analysis

Linear regression was used to assess the association between single-injection tPVB vs no injection and postoperative NRS pain scores (at 5 times: arrival to the surgical wards, noon on POD1, minimum between these 2 time points, maximum between these 2 time points, and at discharge). For pain scores, α level was .05 overall, with significance criterion of $P < .05/5 = .01$ (Bonferroni correction) for each time point. Statistical software (SAS, Cary, NC) was used for all analyses.

3. Results

Between 2009 and 2011, 92 received tPVB (unilateral or bilateral) immediately before surgery, and 90 patients did not receive tPVB (Table 1). Use of tPVB was associated with lower pain scores on arrival to the surgical wards (mean NRS

Table 1 Population data

| | Paravertebral Block (n = 92) | No paravertebral Block (n = 90) |
|-----------------------|------------------------------------|---------------------------------------|
| Age (y) | 51 ± 12 | 54 ± 12 |
| Unilateral mastectomy | 66 (72%) | 52 (58%) |
| Bilateral mastectomy | 26 (28%) | 38 (42%) |

Values are reported as mean (SD) indicated or number of subjects (percentage of group), as indicated.

Download English Version:

<https://daneshyari.com/en/article/2762384>

Download Persian Version:

<https://daneshyari.com/article/2762384>

[Daneshyari.com](https://daneshyari.com)