

◆ Case Reports

Delayed Convulsions and Brief Contralateral Hemiparesis After Retrobulbar Block

Elien Pragt, M.D., André A.J. van Zundert, M.D., Ph.D., F.R.C.A.,
and Chandra M. Kumar, M.B.B.S., M.Sc., F.F.A.R.C.S.I., F.R.C.A.

Objective: This case report describes convulsions and hemiparesis after retrobulbar injection with good outcome in a patient undergoing outpatient cryocoagulation of his right eye.

Case Report: We report a young man in which localized convulsions of the ipsilateral face occurred 9 minutes after retrobulbar injection followed shortly by convulsions of the contralateral arm and leg. After the convulsions, the patient experienced left-sided hemiparesis resolving approximately 1 hour after the injection. There was no hemodynamic instability during this period. It was difficult to determine the exact cause of convulsions and hemiparesis.

Conclusions: We believe these complications occurred because of unintentional injection of local anesthetic agent into the subarachnoid space without affecting the brainstem. Possible mechanisms of spread of local anesthetic agent into the central nervous system after retrobulbar block are discussed. *Reg Anesth Pain Med* 2006; 31:275-278.

Key Words: Retrobulbar block, Complications, Subarachnoid injection, Convulsions, Hemiparesis.

Local anesthesia for major ophthalmic surgical procedures includes peribulbar, retrobulbar, and sub-Tenon's block,¹⁻³ but the exact frequency of their use is not known. Although the classical retrobulbar technique⁴ is still in use, the technique has evolved over the years. In current or modern retrobulbar block, a needle ≤ 32 mm long is inserted through the extreme inferolateral quadrant just below the lateral rectus muscle.⁵ Despite its careful use, sight and/or life-threatening complications have occurred.⁶⁻⁸ Numerous case reports of apnea, respiratory difficulties, cardiovascular collapse, convulsions, and paralysis of 1 or more limbs have been reported after local anesthetic spread to unusual neurologic locations.⁹ The exact mechanism of spread is not fully

understood in most cases. A variety of signs and symptoms after injection of local anesthetic agent into the optic nerve sheath have been reported.⁹ A case of contralateral hemiparesis was reported after retrobulbar block in addition to which the patient developed bilateral ophthalmoplegia and loss of consciousness for approximately 2 hours.¹⁰

We present a case in which localized ipsilateral convulsions occurred 9 minutes after retrobulbar injection, and this was followed shortly by convulsions in the contralateral arm and leg. Contralateral hemiparesis occurred, but the patient recovered full muscle power within an hour. To our knowledge, delayed convulsions and contralateral hemiparesis after retrobulbar block have not been reported.

Case Report

A 36-year-old man (height 1.73 m, weight 80 kg, and American Society of Anesthesiologists 2) was scheduled for outpatient cryocoagulation of his right eye. The medical history included congenital blindness in the right eye and non-Hodgkin's lymphoma with leptomeningeal metastasis. He received intrathecal chemotherapy and skull radiotherapy more than 5 years before. Although his medical conditions remained stable, he developed radiation retinitis in both eyes. He was addicted to nicotine and

From the Department of Anesthesiology, ICU, & Pain Therapy, Catharina Hospital, Brabant Medical School, Eindhoven, The Netherlands (E.P., A.A.J.v.Z.); and Department of Anaesthesia, The James Cook University Hospital, Middlesbrough, United Kingdom.

Accepted for publication January 30, 2006.

Reprint requests: André A.J. van Zundert, M.D., Ph.D., F.R.C.A., Department of Anesthesiology, ICU & Pain Therapy, Catharina Hospital, Brabant Medical School, Michelangelolaan 2, NL-5623 EJ Eindhoven, The Netherlands. E-mail: zundert@iae.nl

© 2006 by the American Society of Regional Anesthesia and Pain Medicine.

1098-7339/06/3103-0016\$32.00/0

doi:10.1016/j.rapm.2006.01.004

other contraband drugs. He did not take any other medications. There was no past history of epilepsy.

He received 3 successive laser therapies for retinitis in both eyes under retrobulbar anesthesia without problems. He was sedated with 60 to 80 mg propofol on each occasion. The axial length of each eye was 21.3 mm. On this occasion, he again received 80 mg propofol in divided doses for sedation before instituting the block. The patient was monitored with an electrocardiogram, peripheral oxygen saturation, and noninvasive blood pressure monitoring, and 4 L/min oxygen was administered. Oxybuprocaine 0.4% eye drops were instilled. A retrobulbar technique, similar to previously used technique, was performed in the right eye by the same experienced anesthesiologist. An Atkinson needle, 25-G, 0.6 × 32 mm, was introduced through the conjunctiva in the extreme inferotemporal quadrant, and 5 mL 1% ropivacaine was injected slowly after negative aspiration for blood and cerebrospinal fluid. There was no resistance to injection, and the feel of the injection appeared as usual. There was no chemosis or conjunctival hemorrhage. The retrobulbar block was the only block administered, and the facial nerve was not infiltrated. An ocular-compression device (Honan balloon) was applied by using a pressure of 25 mm Hg to disperse the local anesthetic agent and to reduce the intraocular pressure. The patient was then transferred from the regional anesthesia block area to the operating room.

Nine minutes after the retrobulbar injection, the patient developed a localized twitching on the ipsilateral face followed, within a minute, by convulsions of left arm and leg. The saturation of the patient decreased to 85% as a consequence of an obstructed airway. A chin lift and administration of 100% oxygen via face mask resolved the problem. The saturation rapidly increased to 100% in 1 minute. Diazepam 5 mg was administered intravenously, which stopped convulsions immediately. Cardiovascular and respiratory systems remained stable. No assisted ventilation or vasopressor drugs were required. Surgery was postponed, and the patient was transferred to the recovery room for further observations. The patient slowly recovered from the residual effects of sedatives and could communicate sensibly 20 minutes after the retrobulbar injection. Although a complete neurologic examination by us would have been very valuable, we failed to notice any neurologic deficit because of rapid change in the clinical conditions of the patient. However, subsequently an examination by a neurologist revealed right-sided facial paralysis and left-sided hemiparesis. At this stage, the eye was checked for the residual effects of retrobulbar block,

and there were no movements except minor flickers. The patient recovered from the residual effects of retrobulbar injection and sedation after 1 hour and did not recall the event. Full neurologic examination revealed full recovery from muscle weaknesses. Magnetic resonance imaging, an electroencephalogram, and cerebral spinal fluid analysis revealed no signs of cerebral pathology or recurrence of non-Hodgkin's lymphoma.

Discussion

Our case exhibited 2 separate but brief phenomena: delayed convulsions and temporary hemiparesis. Despite reviewing the literature, we have not been able to understand the mechanism of local anesthetic spread. Several possible mechanisms might account for these events.

Convulsions can occur because of systemic toxicity either because of absorption or intravascular injection^{9,11} and have been reported after both retrobulbar and peribulbar blocks.^{12,13} It is unlikely for late convulsions to occur because of absorption of a local anesthetic agent, especially when a small dose of local anesthetic agent (50 mg ropivacaine) was used for retrobulbar block when used for any other forms of local anesthetic technique. Another possible mechanism is that of unintentional intra-arterial injection and retrograde flow of local anesthetic into the carotid circulation.¹⁴ This mechanism has been suggested in cases in which immediate grand mal seizure activity has been observed during retrobulbar injection.¹⁵ If unintentional intra-arterial injection was the probable mechanism in our case, convulsions should have occurred within seconds rather than minutes.

Stress or psychological states could also lead to convulsions in susceptible patients,^{15,16} but our patient was heavily sedated with propofol (50 mg propofol intravenously before retrobulbar block is a common practice in our unit and has been successfully administered without significant adverse events in more than 15,000 patients). This patient was young and a known drug abuser; hence, a second dose of 30 mg propofol was slowly administered to keep him sedated for the procedure. He did not show features of deep sedation at any stage during propofol sedation. We cannot completely rule out propofol as the causative agent. Although propofol has been incriminated in inducing convulsions,¹⁷ it has also been used in the management of status epilepticus.¹⁸

Neurologic progression of the original disease might be considered, but negative results of subsequent neurologic investigations probably exclude this possibility. Because the patient had intrathecal

Download English Version:

<https://daneshyari.com/en/article/2767485>

Download Persian Version:

<https://daneshyari.com/article/2767485>

[Daneshyari.com](https://daneshyari.com)