

Relationship between level of HbA1C and breast cancer



Fariba Jousheghany^a, Joshua Phelps^a, Tina Crook^a, Reza Hakkak^{a,b,c,*}

^a Departments of Dietetics and Nutrition, University of Arkansas for Medical Sciences, 4301 W. Markham St., Little Rock, AR 72205, USA

^b Departments of Pediatrics, University of Arkansas for Medical Sciences, 4301 W. Markham St., Little Rock, AR 72205, USA

^c Arkansas Children's Hospital Research Institute, 13 Children's Way., Little Rock, AR 72202, USA

ARTICLE INFO

Article history:

Received 23 March 2016

Accepted 18 April 2016

Available online 23 April 2016

Keywords:

Breast cancer
HbA1C
Diabetes
Tumor stage
Obesity

ABSTRACT

Background: Diabetes and cancer are public health issues worldwide; studies have shown that diabetes is related to increased breast cancer mortality. The purpose of this study was to examine associations between HbA1C and obesity with tumor stage and mortality among breast cancer patients.

Methods: Data for 82 patients with breast cancer (36–89 years of age, diagnosed /treated 1999–2009) were provided by the University of Arkansas for Medical Sciences (UAMS) Data Trust Warehouse. Survival time was estimated from start date of service to date of last follow-up or date of death. The Kaplan–Meier method provided analysis of survival curves for two groups of HbA1C (HbA1C < 6.5% vs HbA1C ≥ 6.5%) and two groups of BMI (BMI < 30 vs BMI ≥ 30 kg/m²); survival curves were compared using log-rank tests. Associations between HbA1C and BMI, and between HbA1C and tumor stage were determined by chi-square.

Results: The relationship between tumor stages and HbA1C was not statistically significant ($X^2 = 0.093$, $p = 0.47$, $df = 1$). The relationship between obesity and HbA1C was statistically significant ($X^2 = 6.13$, $p = 0.013$, $df = 1$). Log-rank tests did not show statistically significant differences between survival curves (HbA1C curves, $p = 0.4$; Obesity curves, $p = 0.09$).

Conclusion: While there was a statistically significant association between HbA1C and obesity, there were no significant associations found with this analysis. However, there are clinically meaningful relationships based on observed trends. Future directions for research may involve exploring a larger sample of patients and the role of therapeutic regimens on blood sugar control and BMI of breast cancer patients and influence on cancer prognosis.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Breast cancer is the most common malignant tumor among women, and of all cancers, it is the second leading cause of mortality in women in the US. Estimates for 2016 predicted that 246,660 women were likely to be diagnosed with invasive breast cancer and that 40,450 women were likely to die from breast cancer [1]. The rate of mortality from breast cancer has declined due to improvements in early detection and treatment [1]. Early detection of the cancer and evaluation of comorbidities result in prompt and relevant treatment that improves breast cancer survival and ultimately saves lives.

Type 2 diabetes may increase the risk of cancer [2–5], is considered a common comorbidity present in a considerable number of breast cancer patients and correlates with poor clinical outcomes [6–9]. Furthermore, diabetes may alter the efficacy of treatment in breast cancer patients affecting drug response and survival endpoints [8,10–12]. Therefore understanding the associations between diabetes and cancer development

will contribute toward improving cancer screening and assessment practices. The aim of this study was to examine associations between HbA1C and obesity with tumor stage and mortality among breast cancer patients.

2. Methods

2.1. Data source

Data for this project came from i2b2 (informatics for integrating biology and the bedside), a large computerized data base at UAMS [13]. The dataset consisted of de-identified secondary data and was exempt from review by the UAMS Institutional Review Board (IRB).

2.2. Study population

A cohort was formed including subjects diagnosed with breast during the years of 2001–2009. African-American (AA) and Caucasian women who were ≥18 years of age at diagnosis, and who had at least one HbA1C on file were included. Subjects with documented pathological tumor stage, BMI and dates of first and last clinic visits

* Corresponding author at: Department of Dietetics and Nutrition, 4301 W. Markham St., Mail Slot 627, Little Rock, AR 72205, USA.

E-mail address: RHakkak@uams.edu (R. Hakkak).

were included in the cohort. Subjects were registered at the oncology clinic. The first visit was defined as the date the patient started service at the clinic and the last follow-up was defined as the last visit. Life /death status was defined according to the Social Security Death Index updated in 2013.

2.3. HbA1C, BMI and tumor stage classification

HbA1C was classified in two groups (normal versus high) with a cut-point value of 6.5% [14,15]. BMI was classified into 2 groups; non-obese was defined as BMI < 30 kg/m² and obese was defined as BMI ≥ 30 kg/m². Tumor stage was classified as low or high based on the TNM classification system (low stages 0–I; high stages II–III) [16].

2.4. Data analysis and statistics

Descriptive statistics including frequencies were used to describe study participant characteristics including age, BMI, HbA1C levels, and tumor stage.

The Kaplan–Meier method provided descriptive analysis of survival curves for the two HbA1C groups and for two BMI groups (BMI < 30 versus BMI ≥ 30 kg/m²); survival curves were compared using log-rank tests. Associations between HbA1C and other variables including BMI and tumor stage were determined by chi-square test. All statistical analyses were performed using Excel (Microsoft Corporation, Seattle, WA), Prism 5 (GraphPad Software, Inc., LaJolla, CA), and IBM SPSS Statistics Version 22 (IBM Corp., Armonk, NY) software.

3. Results

The characterization of the studied population is shown in Table 1. The age of the study population ranged from 36 to 89 years (mean = 60). Sixty-two subjects were ≥50 years of age (categorized as post-menopausal) and 20 were <50 years of age (categorized as pre-menopausal). The study population consisted of 56% Caucasian and 41.5% AA women.

BMI of the study population ranged from 18.73 kg/m² to 53.81 kg/m² (mean = 31.72 ± 7.5 kg/m²). Forty-two subjects (51.2%) were classified as obese (BMI ≥ 30 kg/m²), and 40 (48.8%) were non-obese (BMI < 30 kg/m²). The levels of HbA1C varied from 4.8% to 17% with a mean of 6.9%. Forty-six subjects (54.9%) were grouped into the normal

Table 1
Characterization of the studied population.

	Number (%)	Mean (±SD)
Age		
< 50	20 (24)	60
≥ 50	62 (76)	
Total	82 (100)	
Race		
White	46 (56)	
African American	36 (44)	
BMI		
< 30	40 (49)	25.6 (±2.72)
≥ 30	42 (51)	37.3 (±5.7)
Total	82 (100)	31.7 (±7.5)
Hb1Ac		
< 6.5	46 (55)	5.7 (±0.5)
≥ 6.5	36 (45)	8.3 (±2.6)
Total	82 (100)	6.9 (±2.2)
Tumor stage		
Stage 0	13 (16)	
Stage I	28 (34)	
Stage II	27 (33)	
Stage III	14 (17)	
Low stage (Stages 0 & I)	38 (46.3)	
High stage (Stages II & III)	44 (53.7)	

HbA1C category (<6.5%) and 36 (45.1%) were grouped into the high HbA1C category (≥6.5%).

Thirty-eight subjects were categorized as having low stage tumors (46.3%) and 44 were categorized with high stage tumors (53.7%). A Chi-square test indicated no significant association between HbA1C category and tumor stage ($\chi^2 = 0.093$, $p = 0.47$, $df = 1$).

We further examined the association of high HbA1C on tumor survival. The Kaplan–Meier method provided descriptive analysis of survival curves for the two HbA1C groups (Fig. 1) and for the two BMI groups (Fig. 2). Survival curves were compared using log-rank tests, which determined there were no statistically significant differences between survival curves (HbA1C curves, $p = 0.4$; Obesity curves, $p = 0.09$).

Result of chi-square analysis (Table 2) testing the association between obesity and HbA1C was statistically significant ($\chi^2 = 6.126$, $p = 0.013$, $df = 1$). The percentage of obese patients is clearly larger in the high HbA1C group compared to normal HbA1C group; the percentage of non-obese patients is lower in high HbA1C group compared to normal HbA1C group (Fig. 3).

4. Discussion

Diabetes and cancer are two major public health issues worldwide and many studies have shown that diabetes is related to increased cancer mortality [8,17,18]. According to a review completed by Long and colleagues [19], many studies have shown dysregulation in cellular metabolism may associate with resistance to cancer treatment drugs. Various factors, including insulin, insulin-like growth factor (IGF), and inflammatory cytokines, may link diabetes to breast cancer mortality. In the study reported here, we examined the association of HbA1C as an indicator of blood glucose control with tumor stage and patient mortality among 82 patients diagnosed with breast cancer. We used tumor stage as an indicator of aggressiveness and did not find a statistically significant association between HbA1C levels and tumor stage. These data suggest that blood glucose control as measured by HbA1C is not associated with cancer stage in this population. Our results contradict other studies showing diabetes correlates with late stage diagnosis [9,20]. Peairs et al. [9] performed a meta-analysis examining the effect of pre-existing diabetes on outcomes of breast cancer. Among 4 datasets included in their analysis, 3 showed an association between diabetes and diagnosis at late stage. One study did not find an association between diabetes and breast cancer stage. Peairs et al. [9] suggest that a lack of association between diabetes and breast cancer stage could be due to large number of samples with missing stage in the study. However, a study investigating associations between type 2 diabetes and colon cancer found that, despite a negative association between diabetes and survival rate, there was not an association between diabetes and tumor stage [21]. Together, the data suggest that involvement of diabetes with mortality in breast cancer patients may be due to non-breast cancer specific causes.

According to results reported here, there was not a statistically significant difference between survival curves for the different HbA1C




Fig. 1. Survival analysis was performed using Kaplan–Meier method. Log rank test was used to statistically compare the curves and p value is shown.

Download English Version:

<https://daneshyari.com/en/article/2773052>

Download Persian Version:

<https://daneshyari.com/article/2773052>

[Daneshyari.com](https://daneshyari.com)