



Data in Brief

Expression of microRNAs in HPV negative tonsil cancers and their regulation of PDCD4



Samantha Khoury^a, Alireza Ahadi^a, Xiaoying Zhang^{b,c}, Nham Tran^{a,b,*}

^a Centre for Health Technologies, Faculty of Engineering and Information Technology, University of Technology, Sydney, Australia

^b The Sydney Head and Neck Cancer Institute, Sydney Cancer Centre, Royal Prince Alfred Hospital, Australia

^c Department of Infectious Diseases and Immunology, University of Sydney, NSW, Australia

ARTICLE INFO

Article history:

Received 11 April 2016

Accepted 12 April 2016

Available online 18 April 2016

ABSTRACT

Global rates of tonsil cancer have been increasing since the turn of the millennia, however we still have a limited understanding of the genes and pathways which control this disease. This array dataset which is linked to our publication (Zhang et al., 2015) describes the profiling of human miRNAs in tonsil and normal adjacent tissues. With this dataset, we identified a list of microRNA (miRNA) which were highly over represented in tonsil cancers and showed that several miRNAs were able to regulate the tumour suppressor PDCD4 in a temporal manner. The dataset has been deposited into Gene Expression Omnibus (GSE75630).

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Specifications

Organism/cell line/tissue	Human tonsil SCC and normal adjacent tissues
Sex	Male and female
Sequencer or array type	LNA Exiqon array
Data format	Raw: GenePix Results GPR File
Experimental factors	Tumour vs. normal
Experimental features	Extraction of total RNA from tumour and normal tissues followed by miRNA profiling using custom LNA oligonucleotide arrays
Consent	Level of consent allowed for reuse if applicable
Sample source location	Sydney, Australia

1. Direct link to deposited data

<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE75630>.

2. Experimental design, materials and methods

2.1. Patient cohort

The cohort consisted of 43 patients (39 males, 4 females) treated for tonsillar cancer at Royal Prince Alfred Hospital Sydney, Australia between 2002 and 2006. The mean age was 57 years (range 39–80). Seventeen Tonsil Squamous Cell Carcinomas (SCCs) and matched microscopically normal adjacent (2 cm outside the surgical margin) tissues proved suitable for the profiling analyses (see Fig. 1 for experimental overview).

* Corresponding author at: Centre for Health Technologies, Faculty of Engineering and Information Technology, University of Technology, Sydney, Australia.

E-mail address: nham.tran@uts.edu.au (N. Tran).

2.2. Isolation of RNA from fresh tissue or cultured cells

Approximately 100 mg of fresh frozen tissue was diced and homogenized with a scalpel blade and then mixed with 1 mL of Trizol reagent (Invitrogen, USA). Total RNA was then extracted using isopropanol precipitation and quantified using a NanoDrop ND 1000 (Thermo Fisher Scientific, USA). Samples with ratios of 260/280 in the range of 1.71 to 2.1 were used for the downstream arrays.

2.3. Total RNA labeling

MicroRNAs were labeled at the 3'-end with a P-CU-C3-Cy3 RNA linker by RNA ligation as described [2,3]. 60 µL ligation reaction was prepared with 6 µg of total RNA, 0.1 mM ATP, 20 mM MgCl₂, 3.5 mM DTT, 10 mg/mL BSA, 10% DMSO, 50 mM HEPES, pH 7.8, 250 ng of P-CU-C3-Cy3 (GeneLink, USA) and 20 units of T4 RNA ligase (NEB, USA). The reaction was incubated on ice for 2 h followed by precipitation at –70 °C for 20 min with 0.3 M sodium acetate, 0.5 mg/mL glycogen (Life Technologies, USA) and 2 volumes of 100% ethanol to remove any unbound RNA-linkers. Each labeled sample was dissolved in 30 µL of 400-fold diluted ULS labeled reference set, then mixed with 300 µL Church and Gilbert hybridization buffer. This mixture was denatured at 95 °C for 2 min before hybridization. A mixture of 371 synthetic DNA reference oligonucleotides (Sigma-Genosys, Australia) containing complementary sequences to all LNA probes, was randomly labeled using the ULYSIS labeling kit (Invitrogen, USA) and then filtered using a MicroSpin™ G-25 column (Amersham, USA). Aliquots of a 400-fold dilution of labeled reference set were stored at –20 °C until needed.




Fig. 1. Experimental overview of the project. Total RNA was extracted from cancer and normal tissues and the miRNA population was assessed using a custom printed LNA oligonucleotide array.

2.4. LNA Microarrays

A commercial LNA-modified oligonucleotides library (Exiqon, Denmark) based on miRBase release 7.1, covering 371 human and

mouse miRNA was utilized for expression profiling. Features were deposited onto GAPS II slides (Amersham, USA) at a concentration of 10 μ M (Australian Genome Research Facility, Australia). Individual miRNA LNA probes were printed four times on each array. In addition,

Download English Version:

<https://daneshyari.com/en/article/2821930>

Download Persian Version:

<https://daneshyari.com/article/2821930>

[Daneshyari.com](https://daneshyari.com)