



Carotid intima-media thickness and carotid plaque represent different adaptive responses to traditional cardiovascular risk factors

Liz Andréa Villela Baroncini ^{*,1}, Lucimary de Castro Sylvestre ¹, Roberto Pecoits Filho ¹

Pontifícia Universidade Católica do Paraná, Medical School, Health Sciences Postgraduate Program, Rua Imaculada Conceição 1155 – Bloco CCBS, CEP: 80215-901 Curitiba, Brazil

ARTICLE INFO

Article history:

Received 5 March 2015

Received in revised form 30 July 2015

Accepted 6 August 2015

Available online 8 August 2015

Keywords:

Atherosclerosis

Carotid ultrasound

Carotid intima-media thickness

Hypertension

Dyslipidemia

Diabetes

ABSTRACT

Aim: To assess the effects of each traditional cardiovascular risk factor (hypertension, diabetes mellitus, dyslipidemia, and smoking), including the presence of coronary artery disease (CAD), on carotid intima-media thickness (CINT) and to assess the degree of carotid plaque occurrence.

Methods: A total of 553 outpatients (216 men and 337 women; mean age 67.06 ± 12.44 years) who underwent a carotid artery ultrasound were screened for carotid plaque and CINT measurements.

Results: The CINT medians were higher in males ($P < .001$) and in patients with hypertension ($P < .001$). A linear increase occurred in mean CINT of 0.0059 mm for each year of increase in age. The presence of plaque indicated a tendency to correlate with CINT ($P = .067$). The presence of hypertension associated with diabetes ($P = .0061$; estimated difference 0.0494 mm) or dyslipidemia ($P = .0016$; estimated difference 0.0472 mm) or CAD ($P = .0043$; estimated difference 0.0527 mm) increased the mean CINT measurements. The probability of plaque occurrence in carotid arteries is influenced by the age ($P < .001$) and is higher in patients with dyslipidemia ($P = .008$) and CAD ($P < .001$).

Conclusions: Hypertension is the strongest cardiovascular risk factor that increases CINT, followed by age and male sex. Age and dyslipidemia increase the probability of carotid plaque. Increased CINT and plaque could be present in the same patient caused by different risk factors and with independent effects on the artery wall and different clinical prognoses.

© 2015 The Authors. Published by Elsevier Ireland Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Since 1986 when Pignoli et al. [1], in an in vitro study, concluded that B mode imaging represents a useful tool for detection and monitoring of changes in intimal + medial thickness and allows the evaluation of changes in the arterial wall in areas without localized plaques, carotid intima-media thickness (CINT) measurements have been used in several observational and interventions studies [2–3]. The noninvasive nature of this approach became a recommendation for its use in the pre-clinical diagnosis and follow-up of patients with atherosclerosis. CINT is widely used in clinical research for testing the value of new and emerging risk factors and for evaluation of the effects of various drugs on risk factor modifications [2]. CINT is considered a reflection of multiple risk factors; however, primary contributors to intima-media thickening are age and hypertension, which do not necessarily reflect the atherosclerotic process [3–5]. The primary limitation of CINT is its inability to distinguish lesions with a necrotic core considered a key

indicator of significant plaque advancement and a recognized feature of lesion vulnerability [3]. CINT and carotid plaque, although correlated, reflect different stages and aspects of atherosclerosis and have distinct determinants [6]. A hypertensive hypertrophic response of medial cells can be observed in early phases of atherosclerosis, while carotid plaque formation is often seen in later stages of atherosclerosis and may be caused by inflammation, oxidation, and endothelial dysfunction or smooth muscle cell proliferation, or both [7]. The objective of the present study was to assess the different effects of each traditional cardiovascular risk factor (hypertension, diabetes mellitus, dyslipidemia, and smoking), including also the presence of coronary artery disease (CAD) on CINT and the presence of carotid plaque. The correlation between the presence of plaque and CINT was also investigated.

2. Methods

2.1. Patients

The subjects were selected from a private cardiology clinic next to the ultrasound laboratory. All patients were under the care of a private cardiologist, who assessed the results of blood samples, collected from subjects after overnight fasting. The cardiologist used standard techniques

* Corresponding author.

E-mail address: lizandreabaroncini@hotmail.com (L.A.V. Baroncini).

¹ These authors take responsibility for all aspects of the reability and freedom from bias of the data presented and their discussed interpretation.

to determine serum glucose, total cholesterol, high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol, and triglycerides. The decision to perform a carotid artery study was made exclusively by the private cardiologist. A total of 553 consecutive outpatients (216 men and 337 women; mean age 67.06 ± 12.44 years) who underwent a carotid artery ultrasound were screened for carotid plaque and CIMT measurement, during a period of 6 months. There were no exclusion criteria. Before the study, the ultrasonographer collected information on demographic characteristics and risk factors for each patient according to the report from the private cardiologist. Patients were asked about the presence of hypertension, diabetes mellitus, dyslipidemia, coronary artery disease, and current smoking habit. Hypertension was defined as a history of treated hypertension or the presence of systolic BP ≥ 140 mm Hg or diastolic BP ≥ 90 mm Hg as measured by the private cardiologist. Smoking history was coded as never or current smoker. Subjects were classified as having diabetes when treated for insulin-dependent or non-insulin-dependent diabetes or having elevated fasting glucose levels (≥ 126 mg/dL). The use of lipid-lowering drugs or the presence of total cholesterol > 200 mg/dL, HDL-cholesterol < 40 mg/dL, LDL-cholesterol > 100 mg/dL or triglycerides > 150 mg/dL was recorded. No patient at the present study had a history of stroke or transient ischemic attack and no patient complained of intermittent claudication suggesting peripheral arterial disease. A history of myocardial infarction, angioplasty, or coronary artery by-pass surgery was recorded and a positive CAD history was defined as the presence of any of these diseases in conjunction with the evaluation report from the private cardiologist. Local Institutional ethical committee approval was obtained for the study. Informed consent was obtained from each patient and the study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in a priori approval by the institution's human research committee.

2.2. Ultrasound measurements

Measurements were made with a high-resolution B-mode ultrasonography (Philips Medical Systems' HD11 platform) with a broadband width linear array transducer L 3–12 MHz. Sonography and readings were carried out by trained certified sonographers. On longitudinal 2D ultrasound images of the carotid artery, the near wall and the far wall are displayed as 2 echogenic lines (the adventitia and intima) that are separated by the hypoechoic media. The distance between the leading edge of the first bright line of the far wall (lumen-intima interface) and the leading edge of the second bright line (media-adventitia interface) is defined as the CIMT. Measurements of CIMT were taken manually on the distal 10 mm of both right and left common carotid arteries in the far wall. Five determinations were conducted on each side and the average measurement was used for the CIMT, according to Brazilian Cardiovascular Imaging Department Task Force for Carotid Ultrasound [8]. An artery was classified as being affected by plaque if there was a focal wall thickening at least 50% greater than that of the surrounding vessel wall or a localized region with CIMT greater than 1.5 mm that did not uniformly involve the whole left or right carotid artery with or without a flow disturbance. The vascular ultrasonographer identified plaques in common, internal and external carotids at the time of ultrasound measurement. The presence of plaque was coded as yes or no; however, the degree of stenosis has not been taken into account in the present study.

2.3. Statistical analysis

Categorical variables are expressed as percentages and continuous variables are expressed as mean $\pm$ SD, with a 95% confidence interval (CI) and a significance level of 5%. Multiple linear regression analysis was used to assess the effects of each cardiovascular risk factor (hypertension, diabetes mellitus, smoking, dyslipidemia, and CAD) and the presence of plaque on CIMT. Age and sex were used as control variables. The chi-square test with Yates correction was used to correlate each risk

factor and the presence of plaque. The logistic regression model was used to assess all variables with respect to the presence of plaque. Statistical significance was indicated by a value of $P < .05$. Analyses were performed using IBM SPSS v.20.0 computer software.

3. Results

A total of 553 patients were analyzed. Patients' baseline characteristics are provided in Table 1. At least one risk factor was found in 450 (174 men; 67.6 ± 11.7 years) patients; 103 patients (41 men; 64.6 ± 15.1 years) had no risk factors. Carotid plaque was present in 119 (21.5%) patients; 107 (19.3%) patients with at least one risk factor and 12 (2.1%) patients without risk factors (Table 1). The corrected coefficient of determination was equal to 0.34, indicating that 34% of changes in CIMT were explained by the variables included in the model. The CIMT medians were higher in males and in patients with hypertension. There was a linear increase in mean CIMT of 0.0059 mm for each year increase in age. An interesting result indicates that 25.33% of the variations of CIMT are explained by the age of the individual, excluding the effect of sex, hypertension, diabetes, dyslipidemia, smoking, CAD, and the presence of plaque. Diabetes, dyslipidemia, CAD, and smoking did not influence the CIMT measurements in the present study. The presence of plaque indicated a tendency to influence the CIMT measurements (Table 2). However, there was a significant difference between the effects of hypertension on CIMT and the effects of diabetes, dyslipidemia, and CAD. The presence of hypertension associated with diabetes ($P = .0061$; estimated difference 0.0494 mm) or dyslipidemia ($P = .0016$; estimated difference 0.0472 mm) or CAD ($P = .0043$; estimated difference 0.0527 mm) increases the mean CIMT measurements. Hypertension did not influence CIMT when associated with plaque ($P = .2891$) or current smoking ($P = .1103$). The probability of plaque occurrence on carotid arteries is influenced by age and is higher in patients with dyslipidemia and CAD and had a tendency to be higher in patients who were current smokers. Hypertension and diabetes did not cause an increase in the presence of plaque in the present study (Table 3).

4. Discussion

The main finding of the present study is that CIMT and carotid plaque are influenced differently by traditional cardiovascular risk factors. Hypertension is the risk factor that most increases the CIMT measurements alone and when associated with diabetes, dyslipidemia, and CAD, while dyslipidemia and CAD increase the probability of carotid plaque occurrence. Age influences both CIMT and plaque, and male sex increases CIMT values, but not plaque occurrence. Although intima-media thickness is considered a marker of hypertrophy of intima and media layers of the artery wall, it is not synonymous with atherosclerosis [9–10]. Actually, CIMT and atherosclerosis share common underlying mechanisms for both disease initiation and progression. Pathological intimal thickness is characterized as the formation of lipid pools in the

Table 1
Patient's baseline characteristics.

Patients with risk factors (N/%)	450 (81.3%)
Sex (M/F)	174/276
Age (y $\pm$ SD)	67.6 ± 11.7
Carotid plaque (N/%)	107 (23.8)
History of hypertension (N/%)	333 (74)
History of dyslipidemia (N/%)	249 (55.3)
History of diabetes mellitus (N/%)	100 (22.2)
Cardiovascular history (N/%)	71 (15.7)
Current smoking (N/%)	65 (14.4)
Patients without risk factors (N/%)	103 (18.62)
Sex (M/F)	41/62
Age (y $\pm$ SD)	64.6 ± 15.1
Carotid plaque (N/%)	12 (11.6)

Download English Version:

<https://daneshyari.com/en/article/2926963>

Download Persian Version:

<https://daneshyari.com/article/2926963>

[Daneshyari.com](https://daneshyari.com)