

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: [www.elsevier.com/locate/ihj](http://www.elsevier.com/locate/ihj)

## Original Article

# Cardiac syndrome X: Clinical characteristics revisited



Babu Ezhumalai <sup>a,\*</sup>, Ajith Ananthakrishnapillai <sup>b</sup>, Raja J. Selvaraj <sup>b</sup>,  
Santhosh Satheesh <sup>b</sup>, Balachander Jayaraman <sup>b</sup>

<sup>a</sup> Department of Cardiology, Sri Venkateswara Medical College Hospital and Research Centre, Puducherry, India

<sup>b</sup> Department of Cardiology, Jawaharlal Institute of Postgraduate Medical Education and Research, Puducherry, India

## ARTICLE INFO

## Article history:

Received 7 February 2014

Accepted 25 April 2015

Available online 4 June 2015

## Keywords:

Cardiac syndrome X

Syndrome X

Normal epicardial coronaries

Non-obstructive coronaries

Coronary angiography

## ABSTRACT

**Background:** Cardiac syndrome X includes a heterogeneous group of patients with angina but normal epicardial coronaries in angiography.

**Objective:** Our objective was to study the clinical characteristics of patients with cardiac syndrome X.

**Methods:** Data of patients who underwent coronary angiography over a period of one year was retrospectively analyzed. Those with normal or non-obstructive coronaries in angiography with chest pain were included in this study.

**Results:** 1203 patients underwent coronary angiography during the study period. 105 (8.7%) patients fulfilled the inclusion criteria. There were 52 (49.5%) males and 53 (50.5%) females including 31 (29.5%) postmenopausal women. Many patients had atherosclerotic risk factors. Typical angina and atypical chest pain were reported by 63 (60%) and 42 (40%) patients, respectively. ECG was normal in 46 (43.8%) and abnormal in 59 (56.2%) patients. The most common abnormal finding in ECG was ST-T changes seen in 49 (46.7%) patients. Regional wall motion abnormality with mild left ventricular systolic dysfunction was seen in 4 (3.8%) patients while 101 (96.2%) patients had normal ventricular function in echocardiography. TMT was positive for inducible ischemia in 35 (33.3%) patients and inconclusive in 10 (9.5%) patients. Angiography showed normal epicardial coronaries in 85 (80.9%) patients.

**Conclusions:** Cardiac syndrome X constitutes a significant subset of patients undergoing coronary angiography. It is essential to identify and treat them specifically for microvascular angina. Many of them have atherosclerotic risk factors but their presentation is different from those with obstructive coronaries.

Copyright © 2015, Cardiological Society of India. All rights reserved.

\* Corresponding author. #5, Old Market Street, Mudaliarpet, Puducherry 605004, India. Tel.: +91 9443799346 (mobile).

E-mail address: [drebabu@gmail.com](mailto:drebabu@gmail.com) (B. Ezhumalai).

<http://dx.doi.org/10.1016/j.ihj.2015.04.022>

0019-4832/Copyright © 2015, Cardiological Society of India. All rights reserved.

## 1. Introduction

It is not infrequent to encounter patients with angina or angina-like chest pain having normal or near-normal epicardial coronaries in angiography.<sup>1,2</sup> The term “Syndrome X” was first introduced to define this diagnostic combination by Kemp et al in 1973.<sup>3</sup> Now this entity is known as “Cardiac Syndrome X” (CSX) and it must be differentiated from “Metabolic Syndrome X” and Coronary Artery Disease (CAD). The former is characterized by abdominal obesity, hypertension, reduced high density lipoprotein cholesterol, hypertriglyceridemia and insulin resistance,<sup>4</sup> while the latter is characterized by atherosclerotic obstructive coronaries.

The etiology of CSX is heterogeneous.<sup>5,6</sup> The proposed etiologies are (i) endothelial dysfunction, (ii) microvascular dysfunction or spasm and (iii) abnormal pain perception.<sup>7,8</sup> Patients with CSX have abnormal endothelium-dependent vasoreactivity and impaired vasodilator reserve of coronaries. Microvascular dysfunction or spasm is caused by proinflammatory cytokines released from the dysfunctional endothelium. Autonomic imbalance decreases pain threshold and leads to hypersensitivity to changes in heart rate or contractility.<sup>7</sup> True myocardial ischemia occurs rarely in CSX.<sup>9</sup>

Despite the absence of angiographic abnormalities, many patients with CSX have marked intimal thickening and atheromatous plaque in coronaries on intra-vascular ultrasound imaging.<sup>10</sup> Moreover, multislice computed tomography scanning has shown that the incidence of coronary calcification in CSX (53%) is significantly higher than normal controls (20%) but lower than those with obstructive CAD (96%).<sup>11</sup>

According to prior studies, the prognosis of patients with CSX is generally more favorable than those with obstructive CAD.<sup>6,12,13</sup> On the contrary, recent studies have reported adverse cardiovascular outcomes in patients with non-obstructive coronaries.<sup>14–16</sup> We intended to understand whether such adverse cardiovascular outcomes could be secondary to any change in the clinical characteristics of patients with CSX in the current era. Hence we undertook this study.

## 2. Aim

Our objective was to study the clinical characteristics of patients with cardiac syndrome X.

## 3. Materials and methods

This retrospective, observational study was carried out over a period of one year in a tertiary care hospital. Coronary angiograms of patients who underwent coronary angiography, for suspected ischemic heart disease, during the study periods were reviewed. Patients with normal or non-obstructive coronaries (less than 50% stenosis) in angiography with chest pain were included in this study. The following were the exclusion criteria for our study: coronary angiogram done after acute myocardial infarction, post-revascularisation status, structural heart disease, congenital heart disease and

preoperative indications. Clinical profile and details of investigation such as electrocardiography, echocardiography and treadmill test were analyzed in all the patients included in this study.

Statistical analysis was performed using SPSS software (SPSS Inc. Released 2007. SPSS for Windows, Version 16.0. Chicago: SPSS Inc) and two-tailed *p* value <0.05 was considered significant for all analyses. Categorical variables are expressed as numbers and percentages displayed within parentheses while continuous variables with normal distribution are expressed as ‘Mean ± Standard deviation’.

## 4. Results

A total of 1203 patients underwent coronary angiography during the study period. Of these, 105 (8.7%) patients fulfilled the inclusion criteria. The clinical profile of these subjects included in the study is depicted in Table 1. There were 52 (49.5%) males and 53 (50.5%) females (*p* = 0.89, Chi Square test) including 31 (29.5%) postmenopausal women with overall mean age of 52.9 ± 8.9 years. There were 56 (53.3%) patients with hypertension, 31 (29.5%) patients with type 2 diabetes, 17 (16.2%) patients with body mass index greater than 30 kg/m<sup>2</sup>, 32 (30.5%) patients with dyslipidemia and 19 (18.1%) patients with smoking habit. Analysis of symptoms revealed typical angina more often than atypical chest pain present in 63 (60%) and 42 (40%) patients, respectively (*p* = 0.007, Fisher's exact test).

Resting electrocardiogram was normal in 46 (43.8%) patients and abnormal in the remaining 59 (56.2%) patients. Abnormal findings in ECG included ST-T changes in 49 (46.7%) patients, Q wave in 5 (4.8%) patients and left bundle branch block in 5 (4.8%) patients (Fig. 1). There was no statistically significant difference between normal & abnormal findings in ECG (*p* = 0.09). Echocardiography revealed regional wall motion abnormality with mild left ventricular systolic dysfunction only in 4 (3.8%) patients while the majority of 101 (96.2%) patients had normal study. Symptom-limited TMT performed based on Bruce protocol was positive for inducible ischemia in 35 (33.3%) patients and inconclusive in 10 (9.5%) patients.

**Table 1 – Clinical profile.**

Parameter	n (%)
Age (mean ± SD) in years	52.9 ± 8.9
<b>Gender</b>	
Males	52 (49.5%)
Females	53 (50.5%)
Postmenopausal women	31 (29.5%)
<b>Atherosclerotic risk factors</b>	
Hypertension	56 (53.3%)
Type 2 Diabetes mellitus	31 (29.5%)
BMI > 30 kg/m <sup>2</sup>	17 (16.2%)
Dyslipidemia	32 (30.5%)
Smoking	19 (18.1%)
<b>Symptoms</b>	
Typical angina	63 (60%)
Atypical chest pain	42 (40%)

Note: n – number of patients; BMI – body mass index; SD – standard deviation.

Download English Version:

<https://daneshyari.com/en/article/2927429>

Download Persian Version:

<https://daneshyari.com/article/2927429>

[Daneshyari.com](https://daneshyari.com)