



## Case report

## Myocardial reinfarction with simultaneous occlusions of two major coronary arteries one of which is due to the early stent thrombosis

Celal Kilit <sup>a,\*</sup>, Taner Şen <sup>b</sup>, Adnan Doğan <sup>a</sup>, Basri Amasyalı <sup>a</sup>, Mehmet Özgeyik <sup>a</sup><sup>a</sup> Department of Cardiology, Dumlupınar University, Faculty of Medicine, Kütahya, Turkey<sup>b</sup> Department of Cardiology, Dumlupınar University, Kütahya Evliya Çelebi Training and Research Hospital, Kütahya, Turkey

## ARTICLE INFO

## Article history:

Received 23 November 2015

Received in revised form 20 January 2016

Accepted 22 January 2016

Available online 4 March 2016

## Keywords:

Acute myocardial infarction

Simultaneous occlusion

Stent thrombosis

## ABSTRACT

Coronary artery stent thrombosis is a rare but often fatal complication associated with percutaneous coronary intervention. Although strict adherence to dual anti-platelet therapy minimizes this risk, stent thrombosis will still occur in rare patients, leading to acute, subacute, or late life-threatening coronary syndromes. Intracoronary thrombosis is a usual finding in acute coronary syndrome but simultaneous formation of the thrombi in two different coronary arteries is very rare. Acute coronary syndrome is not simply the result of isolated local plaque disruption and thrombosis, but rather global coronary vessel inflammation, leading to weakening of atherosclerotic plaques in multiple sites nearly simultaneously. We report a case of myocardial reinfarction with simultaneous occlusion of two major coronary branches one of which is due to the early stent thrombosis.

© 2016 The Society of Cardiovascular Academy. Production and hosting by Elsevier B.V. All rights reserved. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Coronary atherosclerosis and thrombosis are major factors contributing to myocardial infarction. However, simultaneous formation of the thrombi in two different coronary arteries is very rare <sup>1</sup>. Patients with ST segment elevation myocardial infarction (STEMI) frequently have obstructive non-culprit lesions <sup>2</sup>. Stenoses in noninfarct arteries may cause serious adverse cardiac events that could be avoided by performing preventive percutaneous coronary intervention (PCI) during the initial procedure. In patients with STEMI and multivessel coronary artery disease undergoing infarct artery PCI, preventive PCI in noninfarct coronary arteries with major stenoses significantly reduced the risk of adverse cardiovascular events, as compared with PCI limited to the infarct artery <sup>3</sup>. We report a case of myocardial reinfarction with simultaneous occlusion of two major coronary branches one of which is due to the early stent thrombosis. In our case, prior primary PCI for inferior STEMI has been limited to the right coronary artery (RCA) which is infarct artery, and preventive PCI has not been performed to significant stenosis in left circumflex artery (LCX).


## Case report

A 69-year-old woman was admitted to emergency department with a sensation of tightness in the chest for 3 h. She had a risk factor of hypercholesterolemia and no other coronary risk factors of smoking, diabetes, hypertension, or positive family history were documented. An electrocardiogram showed ST-segment elevation in leads DII, DIII and aVF (Fig. 1). She underwent cardiac catheterization and emergency angiography revealed significant stenosis in LCX and a total occlusion of RCA after the right ventricular branch (Fig. 2A, B). Left anterior descending artery showed no stenosis. The results of biochemistry laboratory at admission are as follows: Troponin I: 0.56 µg/l (normal range 0–0.023), glucose: 177 mg/dl, creatinine: 0.9 mg/dl, HbA1c: 6.3%, total cholesterol: 236 mg/dl, triglyceride: 106 mg/dl, low density lipoprotein cholesterol: 164 mg/dl, high density lipoprotein cholesterol: 51 mg/dl, hemoglobin: 12.2 g/dl, hematocrit: 36.9%, and platelet:  $333 \times 10^3$ /ml. Balloon angioplasty of the RCA lesion was performed and bare-metal stent was then employed to treat the lesion (Fig. 2C). For anticoagulation, unfractionated heparin was used during the procedure and the patient was loaded with clopidogrel 600 mg orally. There were no complications and the patient tolerated the procedure well. Preventive PCI has not been performed to significant stenosis in LCX. After an uncomplicated 74-hours hospital course, she was discharged with acetylsalicylic acid, clopidogrel, atorvastatin, perindopril, and metoprolol treatment. Four days after discharge, the patient was admitted to the emergency department with chest pain similar to previous. The initial electrocardiogram demonstrated minimal ST elevation and T wave inversion in leads DII, DII and aVF (Fig. 3). The results of

\* Corresponding author at: Dumlupınar University, Kütahya Evliya Çelebi Training and Research Hospital, 43050 Kütahya, Turkey. Tel.: +90 274 2316660 4001; fax: +90 274 2316673.

E-mail addresses: [ckilit@hotmail.com](mailto:ckilit@hotmail.com) (C. Kilit), [medicineman\\_tr@hotmail.com](mailto:medicineman_tr@hotmail.com) (T. Şen), [doganadnan01@gmail.com](mailto:doganadnan01@gmail.com) (A. Doğan), [dramasyali@yahoo.com](mailto:dramasyali@yahoo.com) (B. Amasyalı), [mehmetozgeyik@gmail.com](mailto:mehmetozgeyik@gmail.com) (M. Özgeyik).


Peer review under responsibility of The Society of Cardiovascular Academy.



**Fig. 1.** ECG of the patient at first admission.

biochemistry laboratory at admission are as follows: Troponin I:  $>80 \mu\text{g/l}$  (normal range 0–0.023), glucose: 132 mg/dl, creatinine: 0.79 mg/dl, hemoglobin: 11.6 g/dl, hematocrit: 35.2%, and platelet:  $370 \times 10^3/\text{ml}$ . She was adherent to dual antiplatelet therapy. The patient was diagnosed with myocardial reinfarction probably due to stent thrombosis. A slightly reduced response to clopidogrel (260 P2Y<sub>12</sub> resistance units) was shown in platelet function test (target range for sufficient platelet inhibition,  $<240$  P2Y<sub>12</sub> resistance units). The platelet inhibition response to aspirin was sufficient. Stent thrombosis – at least partly – was thought to be due to slightly reduced response to clopidogrel. Therefore, the patient was loaded with ticagrelor 180 mg orally before the procedure. Emergency angiography revealed a simultaneous total

occlusion of the LCX and an in-stent total thrombotic occlusion of RCA (Fig. 4A, B). Stent in RCA showed no evidence of migration or malposition. The stent occlusion was treated with balloon angioplasty followed by a drug-eluting stent placed distal to the initial stent due to intimal dissection. The LCX occlusion was also treated with balloon angioplasty followed by an adjacent bare-metal stent. For anticoagulation, unfractionated heparin was used during the procedure. She was observed as an inpatient for 3 days and discharged with acetylsalicylic acid, ticagrelor, atorvastatin, perindopril and metoprolol treatment. Recent 1-year follow-up also revealed no significant symptoms and she has been compliant with her dual antiplatelet medication regimen.



**Fig. 2.** (A) Significant stenosis in left circumflex artery. (B) Total occlusion of right coronary artery after the right ventricular branch. (C) Right coronary artery after the bare-metal stent implantation.

Download English Version:

<https://daneshyari.com/en/article/2937094>

Download Persian Version:

<https://daneshyari.com/article/2937094>

[Daneshyari.com](https://daneshyari.com)