EDITORIAL COMMENT

Oxidation, Inflammation, and Aortic Valve Calcification

Peroxide Paves an Osteogenic Path*

Dwight A. Towler, MD, PHD

St. Louis, Missouri

Arterial calcification increasingly afflicts our aging populace (1). Approximately 2% of individuals over age 65 years will require aortic valve replacement (AVR) for calcific aortic stenosis (1). On the basis of recent epidemiologic studies (2), the increasing prevalence of metabolic syndrome and type II diabetes mellitus (T2DM) will further increase the need for AVR—unless strategies are identified and implemented that prevent or reverse valve calcification. Similar concerns exist for 2 other types of vascular mineral deposition: atherosclerotic intimal calcification and medial artery calcification (3). Medial calcification is a strong predictor of lower extremity amputation in T2DM (4), a debilitating and costly outcome. Perturbed Windkessel physiology and altered vascular autonomic responses lead to tissue ischemia (5). Microcalcifications of cholesterol-laden or fibrous

See page 843

components of coronary atherosclerotic plaques attend outward vascular remodeling (6)—harbingers of acute coronary syndrome (7). A better understanding of arterial calcification and vascular mineral metabolism is needed. Once considered only a passive process of dead and dying cells, data from laboratories worldwide have shown that vascular calcification is an actively regulated form of tissue biomineralization (3). In response to metabolic, mechanical, and inflammatory insults, vascular mesenchymal cells elaborate matrix vesicles and gene regulatory programs that: 1) drive osteogenic vascular matrix remodeling (8); and 2) locally neutralize paracrine and systemic inhibitors of calcium deposition (9).

In this issue of the Journal, Miller et al. (10) present an enlightening study that not only reveals the mechanistic underpinnings of human aortic valve calcification but also highlights the critical role of reactive oxygen species (ROS) in the pathobiology of most forms of arterial mineralization. With dihydroethidium staining and lucigenin chemiluminescence, the authors identified increased superoxide levels in stenotic calcified valves versus normal human heart valves. Dihydroethidium staining spatially resolved a gradient of oxidative stress within calcifying aortic valves, with highest levels localizing to regions possessing extensive calcium deposition (10). Dichlorodihydrofluorescein staining for hydrogen peroxide-the more durable ROS product of dismutation that propagates intracellular signals and ironcatalyzed oxidative damage (Fig. 1)-is also increased in regions of valve calcification, notably at the leaflet base (10). This was not due to increased superoxide dismutase (SOD) expression, because SOD isoforms and activities were down-regulated. More importantly, for reasons to be discussed, catalase expression was reduced in both calcified and noncalcified segments of diseased valves compared with normal valves. Thus, increases in ROS "tone" in aortic valves undergoing calcification are accompanied by reductions in defenses that remove several ROS-including hydrogen peroxide, a second messenger (11).

Reduced nicotinamide adenine dinucleotide phosphate (NADPH) oxidases: the road not taken. NADPH oxidase/nitric oxide scavenger (Nox) activities (12) figure prominently in arterial oxidative stress, arising from nonlaminar flow, inflammatory cytokine signaling, and activation of the renin-angiotensin-aldosterone system (13,14). Nox1 and Nox2 play critical roles in the aortic remodeling entrained to angiotensin (13,14). Thus, Miller et al. (10) evaluated whether Nox subunits were increased at venues of aortic valve calcification and oxidative stress. Surprisingly, Nox isoforms were uniformly decreased in calcifying valve segments, and no significant differences in Nox-dependent superoxide generation were measured between normal and diseased valves (10). This was completely unexpected because of the contributions of Nox signaling to atherosclerosis and vascular remodeling (11). Diphenyliodonium-an inhibitor of flavoenzymes such as Nox, xanthine oxidase, and nitric oxide synthase (NOS) (12)-did inhibit superoxide elaborated by calcifying valvular cells, confirming an enzymatic contribution to the generation of valve ROS.

When uncoupled by tetrahydrobiopterin deficiency or inflammation that precludes homodimer formation, NOS monomers use molecular oxygen—rather than arginine—as the terminal electron recipient in the NOS NADPH/flavin/ iron relay (15) (Fig. 1). Therefore, the authors astutely examined the impact of selective NOS inhibition on valve superoxide, implementing the antagonistic arginine analog, nitro-L-arginine methyl ester (L-NAME). L-NAME reduced superoxide production, indicating the contribution of NOS uncoupling to calcified aortic valve ROS generation

^{*}Editorials published in the *Journal of the American College of Cardiology* reflect the views of the authors and do not necessarily represent the views of *JACC* or the American College of Cardiology.

From the Center for Cardiovascular Research, Internal Medicine—Bone & Mineral Diseases, Washington University in St. Louis, St. Louis, Missouri. This work was supported by National Institutes of Health Grants HL69229, HL81138, HL88651, and AR43731 and the Barnes-Jewish Hospital Foundation. Dr. Towler has served as a paid consultant on musculoskeletal disease therapeutics for Wyeth, GlaxoSmithKline, and Eli Lilly and Co. He receives funding for his research from the National Institutes of Health and from the Barnes-Jewish Hospital Foundation.



(10). Had valvular NOS been coupled, L-NAME treatment would have increased superoxide accrual, because NOSdependent nitric oxide production scavenges superoxide via peroxynitrite formation (15) (Fig. 1). Thus, Miller et al. (10) demonstrate that calcifying aortic valves generate a surfeit of superoxide and peroxide via uncoupled NOS activity in the setting of impaired antioxidant defenses—namely, valvular catalase deficiency and reduced nitric oxide production (Fig. 1, asterisks).

Vascular oxidative stress presages osteochondrogenic programming. Miller et al. (10) then related spatial patterns of aortic valve oxidative stress to the elaboration of osteochondrogenic transcription factors known to program biomineralization (16). Runx2/Cbfa1, Msx1, and Msx2 play critical roles in osteogenic mineralization (17). Runx2/ Cbfa1 and Msx2 had been previously identified in calcifying human arteries (18); moreover, in a model of diabetic aortic calcification, Msx2 participates in a signaling relay that entrains osteogenic Wnt/\beta-catenin signaling to vascular inflammation (19,20). Miller et al. (10) identified that Runx2/Cbfa1 and Msx2 were indeed expressed in calcifying human aortic valves, confirming the contribution of active osteochondrogenic regulatory programs to valve calcium accrual (1). Once again, however, another surprise emerged. Whereas the expression of Msx2 was tightly entrained to regions of valve biomineralization, Runx2/Cbfa1 expression was visualized most robustly in adjacent diseased valve

segments-confirmed by reverse transcriptase-polymerase chain reaction analysis (10). The segregation of Msx2 and Runx2/Cbfa1 expression into distinct domains within diseased valves might reflect the actions of the paracrine Wnt signaling milieu that programs osteogenesis (17). Via the cell-surface receptors LRP5 and LPR6, canonical Wnt ligands induce dimerization with coreceptors that activate nuclear β -catenin signals necessary for osteogenic differentiation (reviewed in Johnson et al. [17]). Conversely, these pathways are inhibited by antagonistic ligands such as Dkk1 (17,20). Because Msx2-positive cells elaborate canonical Wnt ligands (Wnt3a and Wnt7a)—but express very little if any Dkk1 (20)—cells in the adjacent vicinity might upregulate Runx2/Cbfa1 expression, a target of Wnt signaling in bone (17). However, the relationship of nuclear β -catenin accumulation to the spatial patterns of Msx2 and Runx2/Cbfa1 expression in calcifying valves has yet to be assessed. Of note, Rajamannan et al. (1) have clearly shown that Wnt3a, LRP5, and β -catenin are up-regulated in calcifying human aortic valves as compared with noncalcifying specimens.

Peroxide paves the path of vascular osteogenesis. Why is this study so significant? In addition to identifying that it is NOS uncoupling—not Nox activation—that generates ROS in calcifying human aortic valves, the authors demonstrate increased accumulation of hydrogen peroxide (H_2O_2) in calcifying valve segments (10). The H_2O_2 is a pro-

Download English Version:

https://daneshyari.com/en/article/2949844

Download Persian Version:

https://daneshyari.com/article/2949844

Daneshyari.com