

Coronary artery ectasia – A sample from Saudi Arabia



Mohammed Atiq Almansori*, Hisham Ali Elsayed

Department of Medicine, Division of Cardiology, University of Dammam

Saudi Arabia

Introduction: Coronary artery ectasia is an uncommon disease that has been increasingly noticed as the increase in utilization of coronary angiograms. The aim of this study is to characterize coronary artery ectasia in the population of Saudi Arabia.

Methods: This is a retrospective study involved all patient with invasive coronary angiogram that was done at a tertiary hospital in Saudi Arabia from January 2011 to December 2013.

Results: A total of 1115 coronary angiograms were reviewed. Coronary artery ectasia was found in 67 patients (6% of all coronary angiograms). The right coronary artery was involved in 73% of cases. And 43% of the cases had severe ectasia.

Conclusion: The prevalence of coronary artery ectasia in Saudi Arabia among patient who went for coronary angiography is higher than what has been published in previous studies and a significant number of patients have severe disease.

© 2015 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Coronary, Ectasia, Atherosclerosis

Introduction

Coronary artery ectasia is an uncommon entity that has been seen with increasing frequency as the utilization of coronary angiograms has expanded, facilitating the diagnosis and management of coronary artery disease. It is defined as dilatation of a coronary artery segment to a diameter at least 1.5 times that of the adjacent normal coronary artery [1–3]. Its first angiographic

description was by Munkner et al. in 1958 [4], but the original description of the disease by Morgagni goes back to 1761 [5]. It has an estimated incidence of 1–5% of all invasive coronary angiograms [3,6,7]. Coronary artery ectasia has been attributed to atherosclerosis in about 50% of cases, related to inflammatory and connective tissue disease in 10–20% of cases, and thought to be congenital in origin in 20–30% of ectasia cases [2,8,9]. The aim of this study is to characterize coronary artery ectasia in the population of Saudi Arabia.

Disclosure: Authors have nothing to disclose with regard to commercial support.

Received 11 August 2014; revised 8 February 2015; accepted 21 March 2015.

Available online 30 March 2015

* Corresponding author at: University of Dammam, PO Box 12875, Dammam 31483, Saudi Arabia. Tel.: +966 545 30 7447; fax: +966 3 896 6741.

E-mail address: malmansori@gmail.com (M.A. Almansori).



P.O. Box 2925 Riyadh – 11461KSA
Tel: +966 1 2520088 ext 40151
Fax: +966 1 2520718
Email: sha@sha.org.sa
URL: www.sha.org.sa



1016–7315 © 2015 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer review under responsibility of King Saud University.
URL: www.ksu.edu.sa
<http://dx.doi.org/10.1016/j.jsha.2015.03.005>



Production and hosting by Elsevier

Methods

This is a retrospective study including all patients who underwent invasive coronary angiograms in a tertiary hospital in Saudi Arabia from January 2011 to December 2013. The data was collected from medical records and hospital databases. An interventional cardiologist reviewed all angiograms, and ectasia was recognized when a dilated segment of a coronary artery had a diameter at least 1.5 times that of adjacent coronary segment. Coronary ectasia was classified based on the extent of the disease, according to the proposal by Markis et al [10]. Diffuse ectasia involving two or three vessels was classified as type I; diffuse disease in one vessel and localized disease in another vessel as type II; diffuse ectasia in one vessel only as type III; and localized ectasia as type IV. Obstructive coronary artery disease was defined as 50% or more diameter stenosis in the left main coronary artery or 70% or more diameter stenosis in a major coronary artery [11]. Slow flow was defined as sluggish clearing of the coronary artery, that is, when the vessel fills completely but either fills or empties slower than uninvolved vessels in the absence of coronary obstruction [12].

The hospital institutional review board approved the study protocol. The data was analyzed and the baseline characteristics were computed using mean and standard deviation for continuous variables and percentage for dichotomous variables.

Results

A total of 1115 coronary angiograms were reviewed. Coronary artery ectasia was found in 67 patients (6% of all coronary angiograms). The baseline characteristics and the biochemical profiles of the patients are shown in Table 1. The mean age of our patients was 56 ± 12 years and most patients were males, constituting 82% of our sample. About 5% of our patients had history of strokes; 57% had history of previous myocardial infarction; 48% had previous history of angioplasty; 15% had history of coronary artery bypass surgery; and no patient had history of connective tissue disease. Fig. 1 shows that coronary artery ectasia was seen more frequently in the right coronary artery, which has ectasia in 73% of the cases, while ectasia of the left main coronary artery was seen in only 8% of the cases. Fig. 2 shows that severe coronary ectasia was seen in a significant portion of our sample, 21% and 22% for coronary ectasia type I and type II respectively, while localized ectasia was seen in 42% of cases. Fig. 3 shows

Table 1. Baseline characteristics and biochemical profile of patients.

Patient characteristics	N = 67
Age (years)	56 ± 12
Sex (male %)	82%
Nationality (Saudi %)	54%
Diabetes mellitus	59%
Hypertension	64%
Dyslipidemia	59%
Smoking	31%
Chronic renal failure	11%
Hemoglobin (g/dL)	13.2 ± 2.2
Glycosylated hemoglobin (%)	7.5 ± 2.1
Total cholesterol (mg/dL)	168 ± 46.8
LDL cholesterol (mg/dL)	100.5 ± 41.7
HDL cholesterol (mg/dL)	35.4 ± 12.2
Triglyceride (mg/dL)	152.9 ± 95.3
Serum creatinine (mg/dL)	1.3 ± 1.5
Ejection fraction (%)	53.4 ± 15.5




Figure 1. Distribution of ectasia in the coronary arteries. LM: left main, LAD: left anterior descending artery, LCx: left circumflex artery, RCA: right coronary artery.




Figure 2. Distribution of different types of coronary ectasia.

that obstructive coronary artery disease was seen in 66% of cases and slow flow in 61%. Regional wall motion abnormalities on echocardiography were seen in 54% of the cases. The mean ejection fraction (EF) in class I was $55.7 \pm 10.5\%$; in class II,

Download English Version:

<https://daneshyari.com/en/article/2977748>

Download Persian Version:

<https://daneshyari.com/article/2977748>

[Daneshyari.com](https://daneshyari.com)